

## A Novel Genetic Variant of *BMP2K* Contributes to High Myopia

Hsin-Ping Liu,<sup>1</sup> Ying-Ju Lin,<sup>2,3</sup> Wei-Yong Lin,<sup>2,5</sup> Lei Wan,<sup>2,3</sup> Jim Jinn-Chyuan Sheu,<sup>2,3</sup> Hui-Ju Lin,<sup>6</sup> Yuhsin Tsai,<sup>3</sup> Chang-Hai Tsai,<sup>4</sup> and Fuu-Jen Tsai<sup>2–4\*</sup>

<sup>1</sup>Graduate Institute of Acupuncture Science, China Medical University, Taichung, Taiwan

<sup>2</sup>Department of Medical Research, China Medical University Hospital, Taichung, Taiwan

<sup>3</sup>School of Chinese Medicine, China Medical University, Taichung, Taiwan

<sup>4</sup>Department of Biotechnology and Bioinformatics, Asia University, Taichung, Taiwan

<sup>5</sup>Graduate Institute of Integrated Medicine, China Medical University, Taichung, Taiwan

<sup>6</sup>Department of Ophthalmology, China Medical University Hospital, Taichung, Taiwan

Loss of eye growth regulation may cause myopia, because modulation of optic globe size is essential for the generation of normal optic power. Evidence has implied variations of *BMP2* gene expression mediate ocular development and retinal tissue remodeling. Given *BMP2* as a potential regulator involved in myopia development, we investigate whether gene *BMP2*-inducible kinase (*BMP2K*, *BIKe*), whose expression is up-regulated during *BMP2*-induced osteoblast differentiation, contributes to susceptibility of high myopia. Participants grouped into high myopia had a spherical equivalent greater than  $-6.00$  D, compared with a control group of spherical equivalent less than  $-0.5$  D. Genotyping of polymorphisms 1379 G/A (rs2288255) and 3171 C/G (rs12507099), corresponding with 405 Gly/Ser and 1002 Thr/Ser variation

in the *BMP2K* gene were determined by PCR-restriction fragment length polymorphism and associative study performed by comparing high myopic subjects and healthy controls. The frequency of A allele in the *BMP2K* gene 1379 G/A polymorphism showed a significant difference between cases and controls ( $P < 0.001$ , OR = 2.99, 95% CI = 1.62–5.54) and subjects with either AA or AG genotype show higher risk than GG genotype ( $P < 0.001$ , OR = 3.07, 95% CI = 1.59–5.92), while 3171 C/G polymorphism was not significant from this survey. These data suggest that *BMP2K* gene 1379 G/A variant is strongly correlated with high myopia and may contribute to a genetic risk factor for high degrees of myopic pathogenesis. *J. Clin. Lab. Anal.* 23:362–367, 2009. © 2009 Wiley-Liss, Inc.

**Key words:** high myopia; bone morphogenetic protein; single nucleotide polymorphism

### INTRODUCTION

Despite occurrence of myopia differs from either country or ethnic groups, myopia is the most common optic disorder in Asian populations with high prevalence up to 70%, based on epidemiological survey (1–3). High predisposition of myopia not only causes aberrant eyesight but also greater risk of blindness and visual impairment, including cataract, retinal detachment, glaucoma, and macular degeneration (4–7), and thus may induce profound impact on public health and social problems. Many animals, as well as human, have ability to modulate axial eye length and focus optical images on the retina to generate normal optical power, known as emmetropisation (8), but patients with myopic pathology show refractive errors and/or loss of adjustment,

leading to concentrate their visual image prior to the retina. Occurrences of high myopia may be complicated and multiple factors. Adjusting the level of hard study or stressful life in Asia, life pressure has been considered an environmental factor in high

Grant sponsor: China Medical University; Grant numbers: CMU95-295; CMU95-138; Grant sponsor: China Medical University Hospital; Grant numbers: DMR-92-008.

Hsin-Ping Liu and Ying-Ju Lin contributed equally to this work.

\*Correspondence to: Dr. Fuu-Jen Tsai, Department of Medical Research, China Medical University Hospital, No. 2 Yuh Der Road, Taichung 404, Taiwan. E-mail: d0704@www.cmuh.org.tw

Received 31 August 2008; Accepted 15 July 2009

DOI 10.1002/jcla.20344

Published online in Wiley InterScience (www.interscience.wiley.com).

prevalence of myopia (9–11). Additional evidence reveals that familial inheritance and genetic basis are also highly responsible for susceptibility to myopic developmental processes (12,13).

Progress of myopia remains unclear. In biological terms, regulation of eye size is likely a key influence on pathological myopia: elongation of ocular globe to result in increased axial length is a feature obtained in patients with myopia and refractive defect (14). Studies on some biological factors in the regulation of this process or on ocular globe remodeling may provide possible clues to identify candidate genes of eye size control and understand more about the causes of pathological forms and high-risk incidence of myopia. Sclera is a specific connective tissue composed of extracellular matrix elements and collagen fibrils lying at the exterior of ocular globe to serve as framework for eye structure and shape maintenance (15,16). During the myopia progression, sclera undergoes dramatic structural change in response to environmental pressure or genetic inheritance. Compared with persons showing normal optic power, high myopic patients tend to have thinner scleral layer and longer-than-average ocular globe, extending axial dimension of the eye and result in refractive error (17–19).

Retina is another fascinating tissue to mediate axial length of eye size (20). Exploration of retinal development uncovered bone morphogenetic proteins (BMPs) and their signaling molecule cascades having an essential influence on the co-operation of eye development through BMP elements and receptors (21–24), despite its original discovery was required for osteogenesis and bone morphogenesis (25,26). Study of chick model of myopia to search for biological factors of eye size control has revealed that *BMP2* gene expression is down regulated at an early form-deprivation stage (27), and in the presence of BMP receptors and secreted BMP2 molecule in chicken embryonic retina have profound effects on eye development, stimulating lens fiber cell differentiation and avoiding cell death in lens epithelium (28), indicating the variation of *BMP2* gene expression as potentially involved in the disturbance of ocular globe development and contribution to myopia development.

In search for regulated factors accounting for variation of eye size, BMP2 is a crucial candidate for retinal development and patterning (29,30). Our study evaluates whether a novel protein BMP2-inducible kinase (BMP2K, BIKE), whose expression is up-regulated during BMP2 induction (31), is correlated with high myopia. To achieve this aim, during performing PCR-restriction fragment length polymorphism (PCR-RFLP) to map the *BMP2K* gene polymorphisms at 1379 G/A and 3171 C/G, we observed that *BMP2K* gene variant

may contribute to genetic background of high myopia pathogenesis in Taiwanese population.

## MATERIALS AND METHODS

### Subjects

All participants enrolled in this study were based on the tenets of the Declaration of Helsinki for research and recruited at China Medical University Hospital in Taiwan. Institutional ethics committee approved this project, with informed consent obtained from all subjects. Their ages ranged from 16 to 25 years, with male-to-female ratio of 1.8:1.0, and all were medical school students. Subjects also met the following criteria: Taiwanese population, no kinship, visual acuity with distance correct of 0.2 logMAR (20/32) or better. Patients with historical diagnosis of any known ocular diseases, intraocular surgery, orthokeratology, retinopathy, prematurity, neonatal problems, and genetic disease and/or connective tissue disorders associated with myopia, such as Stickler or Marfan syndromes, were excluded. The cohort composed of 201 high myopia cases and 86 normal controls collected from February to November 2004. All were diagnosed by measuring refractive error in diopter (D) plus mean spherical equivalent (SE) of the two eyes after one drop of cycloplegic drug (1% mydricle, Alcon, Berlin) application. Those with myopia greater than or equal to  $-6.00$  D in both eyes were included in the case group; subjects had no or mild myopia, with less than or equal to  $-0.5$  D collected as the control group.


### Determination of *BMP2K* Gene Variant by PCR-RFLP

Genomic DNA was extracted from peripheral blood samples by a standard protocol including proteinase K digestion and phenol–chloroform precipitation. Two nonsynonymous single nucleotide polymorphisms (SNP), 1379 G/A (rs2288255) and 3171 C/G (rs12507099), across exonic *BMP2K* locus were selected through public dbSNP database. These polymorphisms of 1379 G/A and 3171 C/G cause codon change at 405 Gly/Ser and 1002 Thr/Ser, respectively. To identify allele preference for the SNPs, PCR-RFLP was designed for genotyping. In brief, PCR reactions were carried out in a total volume of 25  $\mu$ l, containing 5 ng genomic DNA, and specific primers (1379 G/A: forward 5'-CCC AAC CTC AAA CCC TAT TAG TTG-3' and reverse 5'-GGA GAA AGA TGA GGA CTA ATG CG-3'; 3171 C/G: forward 5'-GTA GAT GTA TTT GGC TCC AC-3' and reverse 5'-TTG GAG TCT GAG ATG GTT AA-3'). PCR amplification protocol was set as 95°C for 5 min, followed by 40 cycles of 95°C for 30 sec, 50°C for

30 sec, 72°C for 1 min, and last elongation step at 72°C for 7 min. A 593 bp size of PCR product from 1379 G/A polymorphism was generated and genotyping was performed by restriction enzyme *BstEII* (New England Biolabs, Mississauga, Ontario, Canada) in a total volume of 20 µl at 37°C overnight. Genotyping 3171 C/G polymorphism was determined by a 341 bp size PCR products using restriction enzyme *BfaI*. The digestion fragments were ascertained by agarose gel electrophoresis, and stained with ethidium bromide.

### Statistical Analysis

Distributions of allele and genotype frequencies were performed from each genotyping data by either  $\chi^2$  test or Fisher's exact test (one cell has an expected count smaller than 1) evaluation. Statistical analysis of the odds ratios (OR) and 95% confidence interval (CI) were carried out with SPSS version 10.0 software (Chicago, IL) based on the presence of reference allele and genotype frequencies. Statistical analysis considered only *P* values below 0.05 as significant. Adherence to Hardy–Weinberg equilibrium constant was tested using the  $\chi^2$  test with one degree of freedom.



**Fig. 1.** Genotyping of *BMP2K* gene 1379 G/A polymorphism using PCR followed by restriction enzyme *BstEII* digestion to generate two fragments with 310 and 283 bp length. Lane 1: DNA marker; lanes 2 and 4: homozygous GG and AA genotypes, respectively; lane 3: heterozygous GA genotype.

## RESULTS

### Genotypes for *BMP2K* Gene Polymorphisms

*BMP2K* gene 1379 G/A polymorphism, causing codon change 405 Gly/Ser, was assayed by PCR-RFLP. Genomic DNA samples from high myopia cases and healthy controls were prepared for PCR reaction and obtained a fragment with 593-bp length. Theoretically, two smaller fragments of 310 and 283 bp were obtained in the presence of restriction enzyme *BstEII* treatment. Figure 1 reveals PCR products as digestible, indigestible homozygous “GG” and “AA” genotypes, as well as heterozygous “GA” genotype for *BMP2K* gene. Genotyping 3171 C/G *BMP2K* gene variant was determined by restriction enzyme *BfaI* application. The C allele sequence was separated into two fragments, 120 and 221 bp by *BfaI*, whereas the G allele remained intact (341 bp).

### Different Frequencies of *BMP2K* Gene Polymorphisms in High Myopia Cases and Controls

Table 1 shows distributions of allele and genotype frequencies for *BMP2K* gene 1379 G/A polymorphism in high myopia cases and controls. There was statistical difference in allele frequency for genotyping distribution between high myopia cases and controls ( $P = 0.00029$ ): allele frequency of G:A 80.3:19.7% in high myopia group vs. 92.4:7.6% in control group. Odds ratio of “A” allele presented higher risk of 2.99 folds (95% CI = 1.62–5.54) than “G” allele. Also, marked difference appeared among the distributions of AA, AG, and GG genotype frequencies ( $P < 0.001$ ). Combining AA and AG genotypes, odds ratio in high myopic susceptibility was 3.07 higher risk (95% CI = 1.59–5.92) than GG homozygote, and AA homozygous genotype heightened probability of high myopia. Thus, subjects with “A” allele in the *BMP2K* gene 1379 G/A polymorphism tend to have a higher incidence of pathogenic high myopia.

**TABLE 1. Comparison of the Allele and Genotype Frequencies of *BMP2K* Gene 1379 G/A Polymorphism Between Cases with High Myopia and Healthy Controls**

| Polymorphism |    | Cases  |      | Controls |      | <i>P</i> -value      | Odds ratio (95% CI) | Remarks      |
|--------------|----|--------|------|----------|------|----------------------|---------------------|--------------|
|              |    | Number | (%)  | Number   | (%)  |                      |                     |              |
| 1379 G/A     | G  | 323    | 80.3 | 159      | 92.4 | 0.00029 <sup>a</sup> | 1                   | A vs. G      |
|              | A  | 79     | 19.7 | 13       | 7.6  |                      | 2.99 (1.62–5.54)    |              |
|              | GG | 130    | 64.7 | 73       | 84.9 | 0.00098 <sup>b</sup> | 1                   | AA+GA vs. GG |
|              | GA | 63     | 31.3 | 13       | 15.1 |                      | 3.07 (1.59–5.92)    |              |
|              | AA | 8      | 4.0  | 0        | 0.0  |                      |                     |              |

CI, confidence interval. Hardy–Weinberg equilibrium test:  $\chi^2 = 0.0113$  and 0.5749 for cases and controls, respectively ( $P > 0.5$ ).

<sup>a</sup>*P*-values were calculated by  $\chi^2$  test.

<sup>b</sup>Fisher's exact test (2 × 3).

**TABLE 2. Comparison of the Alleles and Genotypes 3171C/G Polymorphism in BMP2K Gene Between High Myopia Patients and Healthy Controls**

| Polymorphism |    | Cases  |      | Controls |      | P-value <sup>a</sup> | Odds ratio (95% CI) | Remarks   |
|--------------|----|--------|------|----------|------|----------------------|---------------------|-----------|
|              |    | Number | (%)  | Number   | (%)  |                      |                     |           |
| 3171 C/G     | C  | 148    | 72.5 | 124      | 75.6 | 0.5063               | 0.85 (0.53–1.37)    | C vs. G   |
|              | G  | 56     | 27.5 | 40       | 24.4 |                      |                     |           |
|              | CC | 54     | 52.9 | 45       | 54.9 | 0.4916               | 0.45 (0.11–1.80)    | CC vs. GG |
|              | CG | 40     | 39.2 | 34       | 41.5 |                      |                     |           |
|              | GG | 8      | 7.8  | 3        | 3.6  |                      |                     |           |

CI, confidence interval. Hardy–Weinberg equilibrium test:  $\chi^2 = 0.0243$  and 1.2648 for cases and controls, respectively ( $P > 0.5$ ).

<sup>a</sup>P-values were calculated by  $\chi^2$  test.

With 3171 C/G *BMP2K* gene polymorphism, distribution of allele frequency for C:G was 72.5:27.5% in high myopia group vs. 75.6:24.4% in control group (Table 2). Neither C/G allele distribution nor CC/CG/GG genotype frequency showed significant association between high myopia and control groups ( $P > 0.05$ ). In both 1379 G/A and 3171 C/G genotype distributions, they were in accordance with the Hardy–Weinberg equilibrium.

## DISCUSSION

We tested a hypothesis of whether polymorphism within *BMP2K* gene influences susceptibility to high myopia in Taiwanese population. All subjects were young medical school students engaged in intense professional studies, making risk factors of education or near-viewing pervasive from our survey. While genotyping in *BMP2K* gene 1379 G/A (rs2288255), which causes protein codon Gly to Ser alteration at 405 position, we found that frequency of either “AA” or “AG” genotypes revealing higher distribution in high myopia cases than “GG” genotype, as tabulated in statistical analysis. Moreover, subjects with “A” allele had higher risk for high myopia than “G” allele carriers, since there was significant difference in our cohort of case–control participants. Therefore, survey of genetic factor associated with high myopia reveals that polymorphism in *BMP2K* codon 405 is a probable candidate responsible for this disease progression.

Incidence of myopia is rising according to epidemiological evidence, especially in Asia, motivating us to uncover genetic factors involved in pathological myopia for the purposes of decreasing social health costs and preventing this disease. Development of myopia predominantly results from dysfunction of eye growth cooperating by visual signals and outer coat of ocular globe membranes, sclera, choroids, and retina (20,30,32). In particular, it increases axial elongation and reduces tolerance for the intraocular pressure of the

eye (33,34), which may cause irreversible injury to these membranes, increasing risk of high degrees of myopia. Other pathologic diseases such as glaucoma, cataract, and retinal detachment also reveal strong correlation between mechanical stresses and excessive increase in eye size (7).

Exploring cellular factors essential for the control of eye size may help us understand myopic pathology. Changes of TGF $\beta$  expression and activities of TGF $\beta$ -associated signal transduction have been linked with scleral changes and remodeling. Sclera is a connective tissue consisting mainly of collagen (35). In a previous study of a tree shrew model of high myopia, collagen production fell 28% (36). This drop in TGF $\beta$  expression seems to mediate collagen level down-synthesis in sclera (37), and these disturbances may alter scleral extracellular matrix organization. In the longer term, they may lead to enlargement of eye size and thinness of the sclera, a crucial process for the pathogenesis of high myopia (38).

Besides profound influences of TGF $\beta$  on scleral remodeling and myopia, expression of BMPs, including BMP2 molecule and BMPs associated components, has also been responsible for ocular globe patterning and remodeling (29,30). From RT-PCR and Northern blot analysis, BMP2 appeared in mouse retina neurons, retinal pigmented epithelium, and human optic nerve head (22,39). To culture retinal ganglion cells with BMP2 shows BMP2 increasing cell survival, promoting neurite development (40) and BMPs released by retina, suggesting a possible role in the survival and differentiation of lens fiber cells (28). Other research also discovered that adult human cornea can express BMP2 protein, hinting that BMP2 might involve in the proliferation and modulation of keratocyte chemotaxis of human corneal fibroblasts (41,42). BMPs molecules are members of TGF $\beta$  superfamily and originally identified to induce cartilage and bone development (25,26,43). BMPs functions include control of a broad range of biological processes: e.g., regulation of cell

growth, differentiation, and promoting the synthesis of extracellular matrix proteins via a set of serine/threonine kinase receptors (44). After stimulation, activated BMPs transduce their signal to a series of family proteins known as Smads, similar downstream molecular cascades in TGF $\beta$  activation (28,45). Smad4 is probably an important target after BMP stimulation, and activated Smad4 proteins might translocate to nuclei so as to regulate transcription activity of some specific proteins (46,47). Regulation of BMPs signaling cascades may be significant for ocular development, so we try to test correlation between *BMP2K* gene polymorphism and high myopia, a progressive disease linked with eye structure patterning and remodeling.

According to prior data, BMP2K is a serine/threonine kinase protein and contains nuclear localization signal to direct protein to nuclei and may affect transcription activities of target genes (31). Study on the effect of shock wave-medicated changes to human osteoblast highlights stimulated cell proliferation and differentiation, along with induced up-regulation of *BMP2K* gene expression (48). Likewise, protein level of BMP2K increased during BMP2-induced osteoblast differentiation, and its function acts mostly as a regulator of cell differentiation process, suggesting that maintenance of skeletal homeostasis is important for BMP2K protein (31). With BMP2 highly responsible for eye size control and *BMP2* gene down expression at form-deprivation myopia model (27), BMP2-induced protein kinase, BMP2K, may be significant for myopia, but not about BMP2K's association with this progression. We assayed two nonsynonymous SNPs, 1379 G/A and 3171 C/G, in *BMP2K* gene for genotyping in cases vs. control study. Both variants obeyed Hardy-Weinberg equilibrium ( $P > 0.05$ ). Lack of association was found between 3171 C/G polymorphism and high myopia, but the variant at 1379 G/A in *BMP2K* gene showed significant difference between patients with high myopia and healthy controls ( $P = 0.00029$ ). Even though after the Bonferroni correction at 0.05  $\alpha$  level,  $P$  value below 0.025 (0.05/2) is still considered as significant. In our results, allele "A" in 1379 G/A polymorphism is a likely risk factor for myopia due to a higher prevalence in high myopic groups than healthy controls. This polymorphism in 1379 G/A, corresponding with 405 Gly/Ser variation, may cause changes of BMP2K protein activity participating in ocular globe development and leading to myopic progression.

Although this traditional uninformative 1379 G/A polymorphism has been shown to carry important information about the responsiveness of high degrees of myopia, the possible action of BMP2K-mediated eye development still needs to be determined. Present data supply important insight into BMP family

as potential signaling molecules in high myopic pathogenesis.

## REFERENCES

1. Saw SM, Katz J, Schein OD, Chew SJ, Chan TK. Epidemiology of myopia. *Epidemiol Rev* 1996;18:175-187.
2. Seet B, Wong TY, Tan DT, et al. Myopia in Singapore: Taking a public health approach. *Br J Ophthalmol* 2001;85:521-526.
3. Lin LL, Shih YF, Hsiao CK, Chen CJ. Prevalence of myopia in Taiwanese schoolchildren: 1983 to 2000. *Ann Acad Med* 2004;33:27-33.
4. Steidl SM, Pruett RC. Macular complications associated with posterior staphyloma. *Am J Ophthalmol* 1997;123:181-187.
5. Rabb MF, Garoon I, LaFranco FP. Myopic macular degeneration. *Int Ophthalmol Clin* 1981;21:51-69.
6. Hayasaka S, Uchida M, Setogawa T. Subretinal hemorrhages with or without choroidal neovascularization in the maculas of patients with pathologic myopia. Graefe's archive for clinical and experimental ophthalmology. *Albrecht von Graefes Archiv fur klinische und experimentelle Ophthalmologie* 1990;228:277-280.
7. Saw SM, Gazzard G, Shih-Yen EC, Chua WH. Myopia and associated pathological complications. *Ophthalmic Physiol Opt* 2005;25:381-391.
8. Saunders KJ, McCulloch DL, Shepherd AJ, Wilkinson AG. Emmetropisation following preterm birth. *Br J Ophthalmol* 2002;86:1035-1040.
9. Au Eong KG, Tay TH, Lim MK. Education and myopia in 110,236 young Singaporean males. *Singapore Med J* 1993;34:489-492.
10. Feldkamper M, Schaeffel F. Interactions of genes and environment in myopia. *Dev Ophthalmol* 2003;37:34-49.
11. Goss DA. Nearwork and myopia. *Lancet* 2000;356:1456-1457.
12. Wu MM, Edwards MH. The effect of having myopic parents: An analysis of myopia in three generations. *Optom Vis Sci* 1999;76:387-392.
13. Mutti DO, Mitchell GL, Moeschberger ML, Jones LA, Zadnik K. Parental myopia, near work, school achievement, and children's refractive error. *Invest Ophthalmol Vis Sci* 2002;43:3633-3640.
14. Scott R, Grosvenor T. Structural model for emmetropic and myopic eyes. *Ophthalmic Physiol Opt* 1993;13:41-47.
15. Rada JA, Shelton S, Norton TT. The sclera and myopia. *Exp Eye Res* 2006;82:185-200.
16. Muir H. Proteoglycans as organizers of the intercellular matrix. *Biochem Soc Trans* 1983;11:613-622.
17. Rada JA, Achen VR, Penugonda S, Schmidt RW, Mount BA. Proteoglycan composition in the human sclera during growth and aging. *Invest Ophthalmol Vis Sci* 2000;41:1639-1648.
18. Norton TT, Rada JA. Reduced extracellular matrix in mammalian sclera with induced myopia. *Vis Res* 1995;35:1271-1281.
19. Avetisov ES, Savitskaya NF, Vinetskaya MI, Iomdina EN. A study of biochemical and biomechanical qualities of normal and myopic eye sclera in humans of different age groups. *Metab Pediatr Syst Ophthalmol* 1983;7:183-188.
20. Wallman J, Winawer J. Homeostasis of eye growth and the question of myopia. *Neuron* 2004;43:447-468.
21. Belecky-Adams T, Adler R. Developmental expression patterns of bone morphogenetic proteins, receptors, and binding proteins in the chick retina. *J Comp Neurol* 2001;430:562-572.
22. Mathura Jr JR, Jafari N, Chang JT, et al. Bone morphogenetic proteins-2 and -4: Negative growth regulators in adult retinal pigmented epithelium. *Invest Ophthalmol Vis Sci* 2000;41:592-600.

23. Solursh M, Langille RM, Wood J, Sampath TK. Osteogenic protein-1 is required for mammalian eye development. *Biochem Biophys Res Commun* 1996;218:438–443.
24. Bakrania P, Efthymiou M, Klein JC, et al. Mutations in BMP4 cause eye, brain, and digit developmental anomalies: Overlap between the BMP4 and hedgehog signaling pathways. *Am J Hum Genet* 2008;82:304–319.
25. Reddi AH. Role of morphogenetic proteins in skeletal tissue engineering and regeneration. *Nat Biotechnol* 1998;16:247–252.
26. Hogan BL. Bone morphogenetic proteins: Multifunctional regulators of vertebrate development. *Genes Dev* 1996;10:1580–1594.
27. McGlenn AM, Baldwin DA, Tobias JW, Budak MT, Khurana TS, Stone RA. Form-deprivation myopia in chick induces limited changes in retinal gene expression. *Invest Ophthalmol Vis Sci* 2007;48:3430–3436.
28. Belecky-Adams TL, Adler R, Beebe DC. Bone morphogenetic protein signaling and the initiation of lens fiber cell differentiation. *Development (Cambridge, England)* 2002;129:3795–3802.
29. Sakuta H, Takahashi H, Shintani T, Etani K, Aoshima A, Noda M. Role of bone morphogenetic protein 2 in retinal patterning and retinotectal projection. *J Neurosci* 2006;26:10868–10878.
30. Jacobi FK, Zrenner E, Broghammer M, Pusch CM. A genetic perspective on myopia. *Cell Mol Life Sci* 2005;62:800–808.
31. Kearns AE, Donohue MM, Sanyal B, Demay MB. Cloning and characterization of a novel protein kinase that impairs osteoblast differentiation in vitro. *J Biol Chem* 2001;276:42213–42218.
32. Christensen AM, Wallman J. Evidence that increased scleral growth underlies visual deprivation myopia in chicks. *Invest Ophthalmol Vis Sci* 1991;32:2143–2150.
33. Phillips JR, Khalaj M, McBrien NA. Induced myopia associated with increased scleral creep in chick and tree shrew eyes. *Invest Ophthalmol Vis Sci* 2000;41:2028–2034.
34. Siegwart Jr JT, Norton TT. Regulation of the mechanical properties of tree shrew sclera by the visual environment. *Vis Res* 1999;39:387–407.
35. Marshall GE, Konstas AG, Lee WR. Collagens in ocular tissues. *Br J Ophthalmol* 1993;77:515–524.
36. Gentle A, Liu Y, Martin JE, Conti GL, McBrien NA. Collagen gene expression and the altered accumulation of scleral collagen during the development of high myopia. *J Biol Chem* 2003;278:16587–16594.
37. Jobling AI, Nguyen M, Gentle A, McBrien NA. Isoform-specific changes in scleral transforming growth factor-beta expression and the regulation of collagen synthesis during myopia progression. *J Biol Chem* 2004;279:18121–18126.
38. Honda S, Fujii S, Sekiya Y, Yamamoto M. Retinal control on the axial length mediated by transforming growth factor-beta in chick eye. *Invest Ophthalmol Vis Sci* 1996;37:2519–2526.
39. Wordinger RJ, Agarwal R, Talati M, Fuller J, Lambert W, Clark AF. Expression of bone morphogenetic proteins (BMP), BMP receptors, and BMP associated proteins in human trabecular meshwork and optic nerve head cells and tissues. *Mol Vis* 2002;8:241–250.
40. Kerrison JB, Lewis RN, Otteson DC, Zack DJ. Bone morphogenetic proteins promote neurite outgrowth in retinal ganglion cells. *Mol Vis* 2005;11:208–215.
41. Kim WJ, Mohan RR, Mohan RR, Wilson SE. Effect of PDGF, IL-1alpha, and BMP2/4 on corneal fibroblast chemotaxis: Expression of the platelet-derived growth factor system in the cornea. *Invest Ophthalmol Vis Sci* 1999;40:1364–1372.
42. Mohan RR, Kim WJ, Mohan RR, Chen L, Wilson SE. Bone morphogenetic proteins 2 and 4 and their receptors in the adult human cornea. *Invest Ophthalmol Vis Sci* 1998;39:2626–2636.
43. Wozney JM, Rosen V, Celeste AJ, et al. Novel regulators of bone formation: Molecular clones and activities. *Science* 1988;242:1528–1534.
44. Kingsley DM. The TGF-beta superfamily: New members, new receptors, and new genetic tests of function in different organisms. *Genes Dev* 1994;8:133–146.
45. Goumans MJ, Mummery C. Functional analysis of the TGFbeta receptor/Smad pathway through gene ablation in mice. *Int J Dev Biol* 2000;44:253–265.
46. ten Dijke P, Miyazono K, Heldin CH. Signaling inputs converge on nuclear effectors in TGF-beta signaling. *Trends Biochem Sci* 2000;25:64–70.
47. Verschueren K, Huylebroeck D. Remarkable versatility of Smad proteins in the nucleus of transforming growth factor-beta activated cells. *Cytokine Growth Factor Rev* 1999;10:187–199.
48. Hofmann A, Ritz U, Hessmann MH, Alini M, Rommens PM, Rompe JD. Extracorporeal shock wave-mediated changes in proliferation, differentiation, and gene expression of human osteoblasts. *J Trauma* 2008;65:1402–1410.