



Review

# Global and regional estimates of prevalence of refractive errors: Systematic review and meta-analysis

Hassan Hashemi <sup>a</sup>, Akbar Fotouhi <sup>b</sup>, Abbasali Yekta <sup>c</sup>, Reza Pakzad <sup>a</sup>, Hadi Ostadimoghaddam <sup>d</sup>, Mehdi Khabazkhoob <sup>e,\*</sup>

<sup>a</sup> Noor Research Center for Ophthalmic Epidemiology, Noor Eye Hospital, Tehran, Iran

<sup>b</sup> Department of Epidemiology and Biostatistics, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

<sup>c</sup> Department of Optometry, School of Paramedical Sciences, Mashhad University of Medical Sciences, Mashhad, Iran

<sup>d</sup> Refractive Errors Research Center, Mashhad University of Medical Sciences, Mashhad, Iran

<sup>e</sup> Department of Medical Surgical Nursing, School of Nursing and Midwifery, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Received 29 January 2017; revised 1 August 2017; accepted 24 August 2017

Available online 27 September 2017

## Abstract

**Purpose:** The aim of the study was a systematic review of refractive errors across the world according to the WHO regions.

**Methods:** To extract articles on the prevalence of refractive errors for this meta-analysis, international databases were searched from 1990 to 2016. The results of the retrieved studies were merged using a random effect model and reported as estimated pool prevalence (EPP) with 95% confidence interval (CI).

**Results:** In children, the EPP of myopia, hyperopia, and astigmatism was 11.7% (95% CI: 10.5–13.0), 4.6% (95% CI: 3.9–5.2), and 14.9% (95% CI: 12.7–17.1), respectively. The EPP of myopia ranged from 4.9% (95% CI: 1.6–8.1) in South-East Asia to 18.2% (95% CI: 10.9–25.5) in the Western Pacific region, the EPP of hyperopia ranged from 2.2% (95% CI: 1.2–3.3) in South-East Asia to 14.3% (95% CI: 13.4–15.2) in the Americas, and the EPP of astigmatism ranged from 9.8% in South-East Asia to 27.2% in the Americas. In adults, the EPP of myopia, hyperopia, and astigmatism was 26.5% (95% CI: 23.4–29.6), 30.9% (95% CI: 26.2–35.6), and 40.4% (95% CI: 34.3–46.6), respectively. The EPP of myopia ranged from 16.2% (95% CI: 15.6–16.8) in the Americas to 32.9% (95% CI: 25.1–40.7) in South-East Asia, the EPP of hyperopia ranged from 23.1% (95% CI: 6.1%–40.2%) in Europe to 38.6% (95% CI: 22.4–54.8) in Africa and 37.2% (95% CI: 25.3–49) in the Americas, and the EPP of astigmatism ranged from 11.4% (95% CI: 2.1–20.7) in Africa to 45.6% (95% CI: 44.1–47.1) in the Americas and 44.8% (95% CI: 36.6–53.1) in South-East Asia. The results of meta-regression showed that the prevalence of myopia increased from 1993 (10.4%) to 2016 (34.2%) ( $P = 0.097$ ).

**Conclusion:** This report showed that astigmatism was the most common refractive errors in children and adults followed by hyperopia and myopia. The highest prevalence of myopia and astigmatism was seen in South-East Asian adults. The highest prevalence of hyperopia in children and adults was seen in the Americas.

Copyright © 2018, Iranian Society of Ophthalmology. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

**Keywords:** Myopia; Hyperopia; Astigmatism; Meta-analysis

## Introduction

Refractive errors are the most common ocular problem affecting all age groups. They are considered a public health challenge. Recent studies and WHO reports indicate that refractive errors are the first cause of visual impairment and the second cause of visual loss worldwide as 43% of visual

**Declaration of Conflicting Interests:** The authors declare that there is no conflict of interest.

\* Corresponding author.

E-mail address: [khabazkhoob@yahoo.com](mailto:khabazkhoob@yahoo.com) (M. Khabazkhoob).

Peer review under responsibility of the Iranian Society of Ophthalmology.

impairments are attributed to refractive errors.<sup>1</sup> In a review study, Naidoo et al.<sup>2</sup> showed that uncorrected refractive errors were responsible for visual impairment in 101.2 million people and blindness in 6.8 million people in 2010.

Refractive errors also affect the economy of different societies.<sup>3,4</sup> According to a study by Smith et al.,<sup>4</sup> uncorrected refractive errors result in an annual economy loss of \$269 billion worldwide. According to this report,<sup>4</sup> this index was \$121.4 billion in individuals above 50 years.

A review of the literature and medical databases reveals that many studies have been conducted on the epidemiology of refractive errors across the world since 1990.<sup>5,6</sup> Although numerous studies report the prevalence of refractive errors every year, many new articles are published on the epidemiology of these errors annually due to their importance and prevalence.

Although recent studies<sup>7,8</sup> suggest an increase in the prevalence of myopia due to lifestyles changes, differences in ethnic groups, measurement methods, definitions of refractive errors, and age groups of the participants hinder a definite conclusion regarding the pattern of the distribution of refractive errors worldwide.

The distribution of refractive errors is not equal in different countries. A high prevalence of myopia in East Asian countries is a common finding in most previous studies.<sup>7</sup> However, there are some controversies regarding hyperopia. Although some studies have shown a high prevalence of hyperopia in Europe and western countries, it is difficult to make a conclusion since most of these studies were conducted on the elderly, and the high prevalence of hyperopia in this age group is a normal finding due to lens changes. Considering the diversity of the results and use of different definitions and measurement techniques, we decided to evaluate the prevalence of refractive errors across the world in this meta-analysis. Moreover, the status of refractive errors in the world is presented according to the WHO regions in this report.

## Methods

The present meta-analysis was conducted according to the Preferred Reporting Item for Systematic Reviews and Meta-Analysis (PRISMA) guidelines.<sup>9</sup>

### Search strategy

To extract articles from 1990 to 2016 on the prevalence of refractive errors for this meta-analysis, international databases including Medline, Scopus, Web of Sciences, Embase, CABI, CINAHL, DOAJ, and Index Medicus for Eastern Mediterranean Region-IMEMR were searched. The literature was reviewed using a combination of words like population (children, student, adult, and related MeSH terms), outcome [refractive error, myopia, hyperopia, astigmatism, spherical equivalent (SE), cylinder power], and study design (prevalence, ratio, cross-sectional, survey, descriptive, and epidemiology). A search strategy was developed for MEDLINE which then used for other databases. Table 1 presents the

Table 1

Search strategy for MEDLINE (MeSH, Medical Subject Headings).

---

1: Refractive errors [Text Word] OR Refractive errors [MeSH Terms]
2: Myopia [Text Word] OR Myopia [MeSH Terms]
3: Hyperopia [Text Word] OR Hyperopia [MeSH Terms]
3: Astigmatism [Text Word] OR Astigmatism [MeSH Terms]
4: Spherical equivalent [Text Word] OR Spherical equivalent [MeSH Terms]
5: Cylinder power [Text Word] OR Cylinder power [MeSH Terms]
6: 1 OR 2 OR 3 OR 4 OR 5 OR 6
7: Pediatric [Text Word] OR pediatric [MeSH Terms]
8: Children [Text Word] OR children [MeSH Terms]
9: Student [Text Word] OR Student [MeSH Terms]
10: Adolescent [Text Word] OR Adolescent [MeSH Terms]
11: Adult [Text Word] OR Adult [MeSH Terms]
12: 7 OR 8 OR 9 OR 10 OR 11
13: Prevalence [Text Word] OR Prevalence [MeSH Terms]
14: Frequency [Text Word] OR Frequency [MeSH Terms]
15: Cross-Sectional [Text Word] OR Cross-Sectional [MeSH Terms]
16: Descriptive [Text Word] OR Descriptive [MeSH Terms]
17: Survey [Text Word] OR Survey [MeSH Terms]
18: 13 OR 14 OR 15 OR 16 OR 17
19: 6 AND 12 AND 18

---

details of the search strategy. In addition, the reference lists of all searched studies and reviews were evaluated to find similar studies.

### Study selection

After an extensive search, all studies were entered into EndNote X6. Duplicate articles were identified and removed using the duplicates command. Relevant articles were selected in three phases. In phases 1 and 2, the titles and abstracts of the studies were screened, and irrelevant articles were excluded. In phase 3, the full texts of the studies were carefully evaluated. All three phases were conducted by two interviewers independently (S.M. and F.J.). It should be noted that the reviewers were blind to the process of article selection.

The two reviewers had 81% agreement in finding similar studies and 88.7% agreement in data collection. In the remaining 11.3%, the results were evaluated by a third reviewer (M.P.), and the required data were extracted.

### Data extraction and assessment of study quality

The title and abstract of each article was carefully evaluated by 2 reviewers, and data such as the first author's name, publication date, study location (country), study design and characteristics, participants' characteristics (age, sex, sample volume), definitions used for the prevalence of refractive errors, and the prevalence of refractive errors (myopia, hyperopia, and astigmatism) were extracted. The quality of the selected articles was evaluated by the 2 reviewers using the STROBE checklist that contains 22 questions on the methodologic aspects of descriptive studies including the sampling method, study variables, and statistical analysis. The quality assessment results were classified into low quality (less than 15.5), moderate quality (15.5–29.5) and high quality (32–46). Low quality studies were excluded from the meta-analysis.

### Eligibility criteria to select articles for meta-analysis

For studies on children under 20 years of age, only the studies that used cycloplegic refraction were selected for the meta-analysis. For studies on adults, the results of age groups above 30 years were included in the meta-analysis. For studies that were conducted on all age groups, if cycloplegic refraction was used, the first author was contacted by email to obtain the results of cycloplegic refraction in participants below 20 years of age and the results of non-cycloplegic refraction in participants above 30 years of age.

### Statistical analysis

To compare the prevalence of refractive errors in the six WHO regions, we estimated the prevalence of myopia, hyperopia, and astigmatism in each region based on studies with a similar methodology and definition of refractive errors.

Statistical analysis was performed on all studies that were entered into the meta-analysis. The binomial distribution formula was used to calculate the variance and estimated pooled prevalence. The Q statistic with a significance level of 10% was used to evaluate the presence of heterogeneity, and  $I^2$  was used to determine the amount of heterogeneity among studies. To merge the studies, the random effect model was used if there was heterogeneity, and the fix model was used if there was no heterogeneity. The estimated pool prevalence (EPP) was reported for children and adults separately according to WHO regions.

In this study, the WHO regions according to the most recent classification were African Region, Region of the Americas, South-East Asia, Europe, Eastern Mediterranean region, and Western Pacific region.

The forest plot was used to show the total and specific prevalence of refractive errors. Finally, meta-regression analysis was used to evaluate the trend of the prevalence of refractive error with the study year and sample size. It should be mentioned that all analyses were performed with the STATA software version 11.2.

## Results

A total of 9334 articles were identified in this study. After excluding duplicate studies, the titles or abstracts of 4629 articles were reviewed. Then 4326 articles were excluded after reading their abstracts with regards to the inclusion criteria of the study, and 140 articles were excluded after reading their full texts because the required data could not be extracted. Finally, 163 articles were used for the final analysis. However, the number of articles was different for the meta-analysis of myopia, hyperopia, and astigmatism, which is explained in detail in the following sections. Fig. 1 shows the phases of article selection.

Table 2 presents a summary of the results of the studies according to the WHO regions. It should be noted that not all the studies were used in our study, and only the studies that met the criteria were included in the meta-analysis. Table 3




Fig. 1. Flow of information through the different phases of the systematic review.

shows the results of meta-analysis for different refractive errors according to the age group and WHO region.

### Prevalence of myopia

We evaluated 157 studies for myopia. A review of the literature showed different definitions of myopia. Of 157 articles, 130 defined myopia based on a cut point of  $SE \leq -0.5$  D or  $SE < -0.5$  D, of which 67 were conducted on children, and 63 were conducted on adults. Of 67 articles on children, 49 (73.1%) used the cut point of  $SE \leq -0.5$  D and of 63 articles on adults, 50 (79.4%) used the cut point of  $SE < -0.5$  D, which showed a significant difference ( $P < 0.001$ ). Therefore, we used the cut point of  $SE \leq -0.5$  D in children and  $SE < -0.5$  D in adults for myopia in our meta-analysis.

The total sample size of the 49 articles on children that were included in the meta-analysis was 606,155 children. As shown in Fig. 2 and Table 3, the EPP of myopia was 11.7% [95% confidence interval (CI): 10.5–13.0] in all children based on  $SE \leq -0.5$  D. As seen in Fig. 2, according to the WHO regions, the EPP of myopia in children ranged from 4.9% in South-East Asia to 18.2% in the Western Pacific region.

The total sample size of the 50 studies on adults that were included in the meta-analysis was 233,025 participants. The results of meta-analysis based on  $SE < -0.5$  D showed that the EPP of myopia was 26.5% (95% CI: 23.4–29.6) in adults. Myanmar had the highest prevalence (51.0%), and India had the lowest prevalence (4.4%). According to Fig. 3 and Table 3, South-East Asia and the Americas had the highest and lowest EPP of myopia, respectively (32.9% vs. 16.2%). Fig. 4 shows the trend of myopia from 1993 to 2016. The results of meta-regression showed that the prevalence of myopia increased from 1993 (10.4% 95% CI: 7.5–13.6) to 2016 (34.2%: 27.6–40.7) (coefficient = 0.004, 95% CI: –0.001–0.009,  $P = 0.097$ ).

### Prevalence of hyperopia

The prevalence of hyperopia was reported in 146 articles. Although there were different cut points to define hyperopia, a

Table 2  
Summary of studies according refractive errors in worldwide.

Country	Size	Place	Age	Refraction	Myopia		Hyperopia			Astigmatism	
					<-0.5	≤-0.5	≥2	>0.5	≥0.5	≥0.75	≥0.5
USA <sup>10</sup>	11,260	Los Angeles	3–5	Non-cycloplegic		21%			58%		
China <sup>11</sup>	1839	Anyang of Henan	12.9–17.6	Cycloplegic		82.7%			7.5%		
Norway <sup>12</sup>	224	Trondheim	Mean 20.6	Cycloplegic					47%		
China <sup>13</sup>	1565	Inner Mongolia	6–21 Y	Non-cycloplegic	54.1%			15.5%			
USA <sup>14</sup>	4144	Monterey Park	>50	Non-cycloplegic	35.1%			40.2%		45.6%	
Korea <sup>15</sup>	33,355	Seoul	≥5 Y	Non-cycloplegic	51.9%			13.4%			
China <sup>16</sup>	1415	Harbin	≥40 Y	Non-cycloplegic	38.5%			19.9%			
New York <sup>17</sup>	4709	New York	40–84 Y	Non-cycloplegic	21.9%			46.9%			
Puerto Rico <sup>18</sup>	784	Puerto Rico	≥40 Y	Non-cycloplegic	14.7%			51.5%			
Netherlands <sup>19</sup>	520	Dutch	11–13 Y	Non-cycloplegic	28%			8%			
Netherlands <sup>19</sup>	444	Dutch	17–60 Y	Non-cycloplegic	30%			10%			
Bangladesh <sup>20</sup>	11,624	National	≥30 Y	Non-cycloplegic	22.1			20.6%		32.4	
India <sup>21</sup>	1414	Tamil Nadu	>40 Y	Non-cycloplegic	19.4%			39.7%			
Australia <sup>22</sup>	148	Adelaide	44.8 ± 14.5 Y	Non-cycloplegic	31.1%			33.1%			
India <sup>23</sup>	11,786	Hyderabad	≤15 Y	Cycloplegic	3.19%			62.62%			
India <sup>24</sup>	3509	Chennai	>39 Y	Non-cycloplegic		27%		18.7%			54.8%
India <sup>24</sup>	3513	Chennai	>39 Y	Non-cycloplegic		16.8%		52.3%			53%
China <sup>25</sup>	8398	Shanghai	3–10 Y	Cycloplegic		20.1%	17.8%				
California <sup>26</sup>	1501 NHW	Los Angeles and Riverside	6–72 M	Cycloplegic	1.2% <sup>a</sup>		25.65%				
California <sup>26</sup>	1507 Asian	Los Angeles and Riverside	6–72 M	Cycloplegic	3.98% <sup>a</sup>		13.47%				
California <sup>27</sup>	2994	Los Angeles	6–72 M	Cycloplegic			20.8%				
California <sup>27</sup>	3030	Los Angeles	6–72 M	Cycloplegic			26.9%				
Australia <sup>28</sup>	1765	Sydney	6	Cycloplegic			13.2%				
Australia <sup>28</sup>	2353	Sydney	12	Cycloplegic			5.0%				
Brazil <sup>29</sup>	1032	Pelotas	7–15	Non-cycloplegic			13.4%				
India <sup>30</sup>	4074	Hyderabad	7–15 Y	Cycloplegic		4.1%	0.8%			6.30%	
China <sup>31</sup>	5884	Beijing	5–15	Cycloplegic		14.9%	2.6%				
Malaysia <sup>32</sup>	4634	Selangor	7–15 Y	Non-cycloplegic	20.7%					21.3%	
China <sup>33</sup>	2749	Anyang	7.1 Y	Cycloplegic	3.9%		23.3%			25.6	
China <sup>33</sup>	2112	Anyang	12.7 Y	Non-cycloplegic	67.3%		1.2%			28.3	
India <sup>34</sup>	1789	Hyderabad	7–15 Y	Cycloplegic	51.4%		3.3%				
India <sup>34</sup>	1525	Hyderabad	7–15 Y	Cycloplegic	16.7%		3.1%				
Australia <sup>35</sup>	1816	Sydney	6–72 M	Cycloplegic	10.5%		28.9%				
South Africa <sup>36</sup>	1939	Durban	35–90 Y	Non-cycloplegic				37.7%			25.7%
Equatorial Guinea <sup>37</sup>	425	Malabo	6–16 Y	Cycloplegic		10.4%	U(3.1%)			U(32.5%)	
Rwanda <sup>38</sup>	634	Nyarugenge	11–37 Y	Cycloplegic		10.2%		4.3%			4.4%
Ethiopia <sup>39</sup>	4238	Butajira	7–18 Y	Non-cycloplegic		6.0%	0.33%				2.17
Ghana <sup>40</sup>	2435	Ashanti region	12–15 Y	Cycloplegic		3.2%	0.3%				
Kenya <sup>41</sup>	4414	Nakuru	≥50 Y	Non-cycloplegic				27.4%			
Nigeria <sup>42</sup>	13,599	Across the country	≥40 Y	Non-cycloplegic	16.2%			50.7%		63.0%	
Morocco <sup>43</sup>	545	Morocco	6–16 Y	Cycloplegic		6.1%	18.3%			23.5%	
Benin <sup>44</sup>	1057	Cotonou	4–16 Y	Non-cycloplegic						91.9%	
South Africa <sup>45</sup>	4890	Durban	5–15 Y	Cycloplegic		4.0%	2.6%			9.6	
Uganda <sup>46</sup>	623	Kampala	6 and 9	Cycloplegic		11%	37%			52%	
Ethiopia <sup>47</sup>	811	Gondar	6–16 Y	Non-cycloplegic		4.8%	1.6%				0.4%
South Africa <sup>36</sup>	520 (male)	Durban	20–75 Y	Non-cycloplegic		1.9%					5.8%
Ethiopia <sup>48</sup>	420	Debre Markos	7–15 Y	Non-cycloplegic		5.47%	1.4%				1.9%
Ethiopia <sup>49</sup>	1852	Gondar	4–24 Y	Non-cycloplegic		2.3%				1.3%	
Brazil <sup>50</sup>	7654	Sao Paulo	>1 Y	Cycloplegic		25.3%		33.8%			59.7%
Brazil <sup>51</sup>	2454	Botucatu	1–91 Y	Non-cycloplegic				33.8%			59.7%
Brazil <sup>52</sup>	1608	Rio Grande do Sul	7–10 Y	Non-cycloplegic							
Mexico <sup>53</sup>	317	Toluca	6–12 Y	Cycloplegic		9.7%			5.4%		
Brazil <sup>54</sup>	1024	Natal	5–46 Y	Cycloplegic							
Chile <sup>55</sup>	5303	La Florida	5–15 Y	Cycloplegic		5.8%	14.5%			27.2%	
Wisconsin <sup>56</sup>	4275	Beaver Dam	43–84 Y	Non-cycloplegic				49.0%			
California <sup>57</sup>	431	Los Angeles	>55 Y	Non-cycloplegic		10.4%			24.8%		31.8%
China <sup>58</sup>	3070	Yongchuan	6–15 Y	Cycloplegic		13.75%					3.75%
Malaysia <sup>59</sup>	705	Kota Bharu	6–12 Y	Non-cycloplegic		5.4%	1.0%			0.6%	
Singapore <sup>60</sup>	946	Singapore	15–19 Y	Non-cycloplegic		73.9%			1.5%		58.7%

Table 2 (continued)

Country	Size	Place	Age	Refraction	Myopia		Hyperopia			Astigmatism	
					<−0.5	≤−0.5	≥2	>0.5	≥0.5	≥0.75	≥0.5
China <sup>61</sup>	1892	Xichang	11.4–17.1 Y	Cycloplegic			0.2%				1.7%
China <sup>62</sup>	2480	Guangzhou	3–6 Y	Cycloplegic		2.5%	20%				
Nepal <sup>63</sup>	440	Kathmandu	7–15 Y	Cycloplegic						31.0%	
India <sup>64</sup>	Urban: 5021	Maharashtra	6–15	Cycloplegic	3.16%		1.06				0.16
India <sup>64</sup>	Rural: 7401	Maharashtra	6–15	Cycloplegic	1.45%		0.39				0.21
Cambodia <sup>65</sup>	5527	Phnom Penh	12–14 Y	Cycloplegic		5.8%				0.7%	3.76%
Singapore <sup>66</sup>	1232	Tanjong Pagar district	40–79 Y	Non-cycloplegic	38.7%			28.4%			
Myanmar <sup>67</sup>	2076	Meiktila district	≥40 Y	Non-cycloplegic							
Indonesia <sup>68</sup>	1043	Sumatra	≥21 Y	Non-cycloplegic						13.9%	
Japan <sup>69</sup>	3021	Tajimi	>40 Y	Non-cycloplegic	51%			27.9%			
India <sup>70</sup>	2522	Andhra Pradesh	40–92 Y	Non-cycloplegic				18.4%			
South Korea <sup>71</sup>	22,562	Knhanes	>20 Y	Non-cycloplegic	41.8%			24.2			
South Korea <sup>72</sup>	1079	Jeolla	8–13 Y	Non-cycloplegic		46.5%				6.2%	
China <sup>73</sup>	2255	Xuzhou	24–80 M	Cycloplegic	48.1						
Vietnam <sup>74</sup>	2238	Ba Ria – Vung Tau	12–15 Y	Cycloplegic		20.4%	0.4%				0.7%
China <sup>75</sup>	1675	Heilongjiang	5–18 Y	Cycloplegic		5.0%				1.6%	
South Korea <sup>76</sup>	1532	Namil-myeon	≥40 Y	Non-cycloplegic			41.8%				
Nepal <sup>77</sup>	2000	Kathmandu	5–16Y	Cycloplegic	6.85						
India <sup>78</sup>	4711	Not-available	30–100 Y	Non-cycloplegic	20.5%			18.0&			
Laos <sup>79</sup>	2899	Vientiane	6–11 Y	Cycloplegic		0.8%	2.8%				9%
China, <sup>80</sup>	2422	Bai nationality	6–15 Y	Non-cycloplegic	38.1%			22.8%			
Singapore <sup>81</sup>	2804	Southeast district of Singapore	55–89 Y	Non-cycloplegic	30.1		41.5 <sup>a</sup>				
Singapore <sup>82</sup>	2805	Southwestern Singapore	Over 40 Y	Non-cycloplegic	22.8%			35.9%			
Thailand <sup>83</sup>	1100	Bangkok and Nakhonpathom	6–12 Y	Cycloplegic		11.1%	1.4%				0.3%
China <sup>84</sup>	4979	Harbin	≥50 Y	Non-cycloplegic	28.0%			8.9%			
Singapore <sup>85</sup>	2974	Malay	40–80 Y	Non-cycloplegic				27.4%			
China <sup>86</sup>	4364	Guangzhou	5–15 Y	cycloplegic	9.5%		5.8%				33.6%
China <sup>87</sup>	2256	Lanzhou	15–19 Y	Non-cycloplegic	35.1%		0.2%				40.8%
China <sup>88</sup>	4439	Beijing	>40 Y	Cycloplegic	62.3%			19.5%			
China <sup>89</sup>	2515	Yangxi	13–17 Y	Cycloplegic	86.5%		1.20%				25.3%
India <sup>90</sup>	1062	Kanchipuram	6–16 Y	Cycloplegic	21.4%		0.56				
India <sup>91</sup>	2508	Tamil Nadu	>39 Y	Non-cycloplegic	42.4%			18.70%			
India <sup>92</sup>	6447	New Delhi	5–15 Y	Cycloplegic		7.4%	7.7%				10.19%
Nepal <sup>93</sup>	5067	Mechi zone	5–15 Y	Cycloplegic		1.2%	2.1%				3.5%
Poland <sup>94</sup>	5724	Szczecin	6–18 Y	Cycloplegic		13%					4.0%
Poland <sup>95</sup>	4422	Szczecin	6–18 Y	Cycloplegic		13.3%					
Sweden <sup>96</sup>	143	Gothenburg	4–15	Cycloplegic		6%	9%				22%
England <sup>97</sup>	2495	Not available	44–46 Y	Non-cycloplegic	47.8		8.8 <sup>a</sup>				
England <sup>98</sup>	7444	Not available	48–92 Y	Non-cycloplegic	23		39.4 <sup>a</sup>				
Northfolk											
Norway <sup>97</sup>	5792	Not available	38–87 Y	Non-cycloplegic		19.4	33.7 <sup>a</sup>				
Greece <sup>97</sup>	1952	Not available	60–94 Y	Non-cycloplegic		14.2	39.4 <sup>a</sup>				
France <sup>97</sup>	618	Not available	73–93 Y	Non-cycloplegic		16.7	53.6 <sup>a</sup>				
Netherlands <sup>97</sup>	2662	Not available	14–87 Y	Non-cycloplegic		21.2	27.4 <sup>a</sup>				
Germany <sup>97</sup>	14,069	Not available	35–74 Y	Non-cycloplegic		31.9	23.9 <sup>a</sup>				
France <sup>97</sup>	576	Not available	76–92 Y	Non-cycloplegic		19.1	51.1 <sup>a</sup>				
France <sup>97</sup>	2315	Not available	60–93 Y	Non-cycloplegic		16.2	53 <sup>a</sup>				
Netherlands <sup>97</sup>	6566	Not available	55–106 Y	Non-cycloplegic		16.4	52.3 <sup>a</sup>				
Netherlands <sup>97</sup>	2579	Not available	55–99 Y	Non-cycloplegic		21.9	45.7 <sup>a</sup>				
Netherlands <sup>97</sup>	3530	Not available	46–97 Y	Non-cycloplegic		32.5	28.8 <sup>a</sup>				
UK <sup>97</sup>	6095	Not available	16–85 Y	Non-cycloplegic		31.4	26 <sup>a</sup>				
Germany <sup>97</sup>	2372	Not available	35–84 Y	Non-cycloplegic		36.1	24 <sup>a</sup>				
England <sup>98</sup>	4488	Not available	48–89 Y	Non-cycloplegic		27.8%			49.4%		
Germany <sup>99</sup>	13,959	Gutenberg	35–74 Y	Non-cycloplegic	35.1			31.8%		32.3	
Spain <sup>100</sup>	417	Segovia	40–79 Y	Non-cycloplegic		25.4%			43.6%	53.5	
Greece <sup>101</sup>	1500	Athens	40–77 Y	Non-cycloplegic	35.1%			14.40%			
Sweden <sup>102</sup>	1045	Goteborg	12–13 Y	cycloplegic		49.7%					
Turkey <sup>103</sup>	21,062	Diyarbakir	6–14 Y	Cycloplegic		3.2%					14.3%
Pakistan <sup>104</sup>	45,122	Rawalpindi	5–16 Y	Cycloplegic		1.89%					0.76%

(continued on next page)



Table 2 (continued)

Country	Size	Place	Age	Refraction	Myopia		Hyperopia		Astigmatism	
					<−0.5	≤−0.5	≥2	>0.5	≥0.5	≥0.75
Turkey <sup>105</sup>	709	Eskisehir	7–8 Y	cycloplegic		22.6%				11.0%
Iran <sup>106</sup>	1367	Mashhad	>54	Non-cycloplegic	27.2%			51.6%		37.5%
Iran <sup>107</sup>	1854	Shiraz	7–15	Cycloplegic		4.35%	5.04%			11.27%
Iran <sup>108</sup>	201	Khaf	19–90	Non-cycloplegic	28			19.2%		14.3%
Iran <sup>109</sup>	1551	Bojnord	6–17	Cycloplegic		4.3%		5.4%		11.5%
Iran <sup>110</sup>	937	Sari	55–87	Non-cycloplegic				39.5%		
Iran <sup>111</sup>	2098	Yazd	40–80	Non-cycloplegic	36.5			20.6%		53.8
Jordan <sup>112</sup>	1647	Tafila	12–17 Y	Non-cycloplegic		63.5%			11.2%	
Jordan <sup>113</sup>	1093	Amman	17–40 Y	Non-cycloplegic	36.3%			5.67%		36.8%
Saudi Arabia <sup>114</sup>	1319	Riyadh	4–6 Y	Non-cycloplegic				2.1%		
Saudi Arabia <sup>115</sup>	1536	Riyadh	12–13 Y	Non-cycloplegic	53.71%					
Pakistan <sup>116</sup>	917	Khyber Pakhtunkhwa	>30 Y	Non-cycloplegic	2.5%					
Iran <sup>117</sup>	4072	8 Cities	7 Y	Cycloplegic	4.5%	3.04%	6.20%			
Pakistan <sup>118</sup>	1644	Kohat	5–15 Y	Non-cycloplegic						
Iran <sup>119</sup>	2410	Tehran	7–12 Y	Cycloplegic		4.9%	3.5%			22.6%
Iran <sup>120</sup>	1109	Dezful	6–15 Y	Cycloplegic	3.4%		12.9%			
Iran <sup>121</sup>	3675	Mashhad	4–6 Y	Non-cycloplegic						
Pakistan <sup>122</sup>	300	Haripur	5–20 Y	Cycloplegic	14.9%	52.6%				28.4%
Pakistan <sup>123</sup>	533	Lahore	9–18 Y	Non-cycloplegic						
Iran <sup>124</sup>	2124	Khaf	16–65 Y	Non-cycloplegic						
Iran <sup>125</sup>	434	Aligoudarz	14–21 Y	Non-cycloplegic		29.3%				21.7%
Iran <sup>126</sup>	1431	Mashhad	18–32 Y	Non-cycloplegic				7.8%		
Iran <sup>127</sup>	5020	Shahrud	40–64 Y	Non-cycloplegic						
Iran <sup>128</sup>	2048	Mashhad	>15 Y	Non-cycloplegic		22.36%			34.21%	25.64%
Iran <sup>128</sup>	765	Mashhad	≤15 Y	Cycloplegic		3.64%	27.4%			
Iran <sup>129</sup>	5903	Qazvin	7–15 Y	Cycloplegic		65%,	12.46%			16.1%
China <sup>130</sup>	1269	Liwan	≥50 Y	Subjective	32.3%			40.0%		
Pakistan <sup>131</sup>	14,490	Nationally	>30 Y	Non-cycloplegic				27.1%		
Iran <sup>132</sup>	5544	Dezful	7–15 Y	Cycloplegic		3.4%	16.6%			18.7%
Pakistan <sup>133</sup>	2317	Kolkata	5–10 Y	Non-cycloplegic	36.5%					
Australia <sup>134</sup>	1936	Sydney	4–12 Y	Non-cycloplegic	14.02%		8.4%		38.4%	
Australia <sup>135</sup>	2535	Sydney	4–12 Y	Non-cycloplegic	3.8%	6.5%				39.25%
Australia <sup>136</sup>	3654	Sydney	49–97 Y	Non-cycloplegic				57%		37%
Singapore <sup>137</sup>	10,033	Singapore	>40 Y	Non-cycloplegic		38.9%		31.5%		
Iran <sup>138</sup>	4864	Shahrud	40–65 Y	Cycloplegic	30.2			35.6		
Argentina <sup>139</sup>	1518	Buenos Aires	25–65 Y	Non-cycloplegic				18.1%		
China <sup>140</sup>	6491	Handan	≥30 Y	Non-cycloplegic				15.9%		
Iran <sup>141</sup>	4354	Tehran	≥5 Y	Non-cycloplegic		21.8%		26%		29.6%
Iran <sup>141</sup>	4354	Tehran	≥5 Y	Cycloplegic		17.2%		56.6%		30.3%
China <sup>142</sup>	4319	Beijing	40–90 Y	Non-cycloplegic				20%		
Taiwan <sup>143</sup>	2045	Taipei	≥65 Y	Non-cycloplegic	19.4%			59%		
China <sup>140</sup>	6491	Handan	40–79 Y	Non-cycloplegic	22.9%			1.6%		
Mongolia <sup>144</sup>	1617	Hövsögöl and Omnögobi	≥40 Y	Non-cycloplegic	19.4%			32.9%		
Australia <sup>145</sup>	4744	Victoria	≥40 Y	Non-cycloplegic	19.4%			18%		
India <sup>146</sup>	5885	Central Maharashtra	≥30 Y	Non-cycloplegic	17.2%			18%		
Australia <sup>147</sup>	1884	Central Australia	>20 Y	Non-cycloplegic	17%					
Norfolk Island <sup>148</sup>	677	Norfolk Island	≥15 Y	Non-cycloplegic	17%					
Maryland <sup>149</sup>	6000	10 Cities	45–84 Y	Non-cycloplegic	11.1%					
Iran <sup>150</sup>	815	Shahrood	6 Y	Cycloplegic		20.5%	1.7%			
Mongolia <sup>151</sup>	1057	Khovd	7–17 Y	Non-cycloplegic						
India <sup>152</sup>	1378	Bangalore	7–15 Y	Cycloplegic		4.4%				
Mexico <sup>153</sup>	1035	Monterrey	12–13 Y	Cycloplegic		44%				
Iran <sup>132</sup>	3490	Dezful	7–15	Cycloplegic	3.4%		16.6			18.7

Y: Year, M: Month.

<sup>a</sup> Spherical equivalent (SE) worse than >0.75 diopter (D).

common point in children who underwent cycloplegic refraction was the use of SE  $\geq +2$  D as the cut point. We also considered this cut point for children who underwent cycloplegic refraction. As for adults, since about 74% of the studies used SE  $> +0.5$  D to define hyperopia, we also adopted this cut point for the meta-analysis of hyperopia.

A total of 91 articles were included in the meta-analysis of hyperopia, 45 of which were conducted on children (cycloplegic refraction, SE  $\geq +2$  D) and 46 on adults (non-cycloplegic refraction, SE  $> +0.5$  D).

The total sample size of the 45 articles analyzed for children was 200,995 participants. The results of meta-analysis of

Table 3  
Estimated pool prevalence (EPP) of myopia, hyperopia, and astigmatism in children and adult by WHO regions.

Astigmatism	Astigmatism	Hyperopia	Myopia
	%EPP(95%CI); weight	%EPP(95%CI); weight	%EPP(95%CI); weight
<b>Children</b>			
Africa	14.2 (9.9–18.5); 10.33	3.0 (1.8–4.3); 10.57	6.2 (4.8–7.6); 16.48
Americas	27.2 (26–28.4); 2.11	14.3 (13.4–15.2); 4.14	8.4 (4.9–12); 6.09
South-East Asia	9.8 (6.3–13.2); 16.47	2.2 (1.2–3.3); 20.89	4.9 (1.6–8.1); 8.52
Europe	12.9 (4.1–21.8); 6	9 (4.3–13.7); 1.04	14.3 (10.5–18.2); 16.04
Eastern Mediterranean	20.4 (14.5–26.3); 29.11	6.8 (4.9–8.6); 30.75	9.2 (8.1–10.4); 26.69
Western Pacific	12.1 (8.4–15.8); 35.98	3.1 (1.9–4.3); 32.59	18.2 (10.9–25.5); 26.18
All	14.9 (12.7–17.1); 100	4.6 (3.9–5.2); 100	11.7 (10.5–13.0); 100
<b>Adult</b>			
Africa	11.4 (2.1–20.7); 8.85	38.6 (22.4–54.8); 6.54	16.2 (15.6–16.8); 2.01
Americas	45.6 (44.1–47.1); 2.95	37.2 (25.3–49); 13.05	22 (16.4–27.7); 7.98
South-East Asia	44.8 (36.6–53.1); 17.58	28 (23.4–32.7); 21.79	32.9 (25.1–40.7); 18.02
Europe	39.7 (34.5–44.9); 8.82	23.1 (6.1–40.2); 4.36	27 (22.4–31.6); 29.99
Eastern Mediterranean	41.9 (33.6–50.2); 29.39	33 (26.9–39); 19.54	24.1 (14.2–34); 13.98
Western Pacific	44.2 (30.6–57.7); 32.41	28.5 (20.1–37); 34.73	25 (20–30.1); 28.01
All	40.4 (34.3–46.6); 100	30.9 (26.2–35.6); 100	26.5 (23.4–29.6); 100

EPP: Estimated pool prevalence.  
CI: Confidence interval.

hyperopia in children are presented in Table 3 and Fig. 5. The EPP of hyperopia was 4.6% (95% CI: 3.9–5.2) in children. According to the WHO regions, the lowest and highest EPP was seen in South-East Asia (2.2%, 95% CI: 1.2–3.3) and the Americas (14.3%, 95% CI: 13.4–15.2), respectively.

The total sample size of the 46 articles analyzed for adults was 199,691 participants. The results of meta-analysis of hyperopia in adults are presented in Table 3 and Fig. 6.

The EPP of hyperopia was 30.6% (95% CI: 26.1–35.2) in adults. Based on the results of meta-analysis, Africa had the highest EPP of hyperopia (38.6%, 95% CI: 22.4–54.8) followed by the Americas (37.2%, 95% CI: 25.3–49) while Europe had the lowest EPP (23.1%, 95% CI: 6.1–40.2). The trend of hyperopia was not significant in the past three decades (coefficient:  $-0.005$ ; 95% CI:  $-0.012$  to  $0.002$ ,  $P = 0.196$ ) (Fig. 7).

### Prevalence of astigmatism

The definition of astigmatism in epidemiologic studies has less variation. The results of 135 studies on astigmatism were collected which used different cut points to define astigmatism. A cylinder power  $\geq 0.5$  D and a cylinder power  $> 0.5$  were more common definitions in epidemiologic studies. The most common cut point was a cylinder power  $> 0.5$  D according to which 82 out of 135 articles on astigmatism were included in the meta-analysis. Considering the changes of astigmatism with age, the articles were divided to those conducted on children and adults. For studies that evaluated age groups above 1 year of age, the data of adults and children were analyzed separately.

48 articles were included in the meta-analysis for children with a total sample size of 152,570 participants. According to Table 3 and Fig. 8, the EPP of astigmatism was 14.9% (95% CI: 12.7–17.1) in children. According to WHO regions, the lowest EPP was seen in South-East Asia (9.8%) while the

highest EPP was seen in the Americas (27.2%) followed by the Eastern Mediterranean region (20.4%).

For adults, 34 articles with a total sample size of 122,436 participants were included in the meta-analysis. The results showed that 40.4% (95% CI: 34.3–46.6) of adults had astigmatism (Fig. 9). However, astigmatism showed a lot of variation in different WHO regions; the highest EPP of astigmatism was seen in the Americas, and the lowest EPP was seen in Africa (11.4% vs. 45.6). However, it should be noted only one study was conducted in the Americas. After the Americas, South-East Asia had the highest EPP of astigmatism (44.8%, 95% CI: 36.6–53.1). The trend of astigmatism was not significant in the past three decades (coefficient:  $0.003$ ; 95% CI:  $-0.006$  to  $0.011$ ,  $P = 0.559$ ).

### Discussion

Refractive errors are the most common visual problems.<sup>1</sup> Due to their importance, many studies have evaluated their epidemiology, etiology, and treatment methods. Numerous studies across the world have reported the prevalence of refractive errors as an index of descriptive epidemiology, and it may be the only field in refractive errors which includes reports from almost every corner of the world.<sup>2–4, 8,12,14,17–52,54–71,73–76,78–103,105–117,119–130,132–169</sup>

The distribution of refractive errors is clear in some parts of the world according to previous studies; for example, we already know that myopia is prevalent in East Asian countries. However, despite the considerable number of studies on the prevalence of refractive errors, few studies have reviewed the epidemiology of refractive errors systematically to show the status of refractive errors across the world. Due to the importance of refractive errors and scarcity of review and meta-analysis studies in this regard, we evaluated the prevalence of refractive errors systematically in this meta-analysis.




Fig. 2. Forest plot of estimated pool prevalence (EPP) of myopia [spherical equivalent (SE) ≤ -0.5] in children by WHO regions.

The results of different studies in different age groups showed that prevalence of myopia ranged from 0.8% in children aged 6–11 years in Laos<sup>79</sup> to 86.5% in 15–19-year-old Chinese<sup>89</sup> children. However, defining myopia as SE < -0.5 D in adults and SE ≤ -0.5 D in children and considering the

results of cycloplegic refraction in children limited this range. The EPP of myopia was about 11.7% in children, ranging from 0.8% in Laos<sup>79</sup> to 47.3% in China. As mentioned earlier, the lowest prevalence of myopia was seen in South-East Asia, and the highest prevalence was seen in the Western Pacific region.






Fig. 3. Forest plot of estimated pool prevalence (EPP) of myopia [spherical equivalent (SE) ≤ -0.5] in adults by WHO regions.




Fig. 4. Trend of myopia from 1990 to 2016.

Previous studies showed myopia aggregation in South-East Asian countries while according to this meta-analysis, myopia aggregation in children is seen in the Western Pacific region.<sup>7</sup> However, it is rather difficult to explain the low prevalence of myopia in South-East Asian children, but it seems that one of the reports from Nepal<sup>63,77,93,170</sup> with a very large sample size decreased the estimated prevalence of myopia in this region.

In adults, the prevalence of myopia ranged from 4% to 51%, and the EPP of myopia was 26.5%. The highest prevalence of myopia in adults was seen in South-East Asia, and the lowest prevalence was seen in the Africa. A comparison of the results of myopia in children and adults suggests different questions and hypotheses as to why children have the lowest and adults have the highest prevalence of myopia in South-East Asia. It seems that the limited number of studies on children in South-East Asia is one of the reasons for this finding while there is more variation in adults. On the other hand, in South-East Asia, only studies on children and adults from India were included in our meta-analysis; therefore, it may be rather difficult to make a comparison and the finding may be influenced by the Indians' race.

It seems that in countries like South-East Asian countries where the prevalence of myopia is low in children and high in adults, environmental factor have a more prominent role than genetic and ethnic factors, or the genes responsible for myopia in these individuals are expressed at higher ages.

It has been previously shown that some genes are responsible for myopia; however, it is well documented that the genes cannot cause myopia per se.<sup>7</sup> In 1969, a study<sup>171</sup> was conducted on Eskimos in northern Alaska whose living conditions were about to change. Only 2 out of 131 adults who grew up in isolated communities had myopia whereas more than half of their children and grandchildren were myopic.

Regarding this meta-analysis, we believe that countries like China and Singapore that are categorized under the Western Pacific region have genetic differences with the current South-East Asian countries because the distribution of myopia in childhood and adulthood is similar in these countries. With regards to the high prevalence of myopia in children and adults

in Europe, we believe that the role of genetic and ethnic factors is more important than environmental factors.

As mentioned earlier, children in South-East Asia had the lowest and children in the Americas had the highest prevalence of hyperopia. In adults, Africans and Americans had the highest and Europeans had the lowest prevalence of hyperopia. It is a little perplexing to explain the results; however, the results of meta-analysis showed a high prevalence of hyperopia in American children and adults. Moreover, although the prevalence of hyperopia in African adults was a little higher than American adults, its prevalence was higher in American children. Emmetropization may play a role in this regard, and in addition to ethnic and genetic factors, differences in computerization and lifestyle changes may have contributed to increased prevalence of hyperopia in African and American regions as compared to other parts of the world. The role of myopization in hyperopia becomes more prominent when we consider the results of Europe where the prevalence of hyperopia is the lowest and the prevalence of myopia ranks second.

The results of our meta-analysis showed that about 15% of children and 40% of adults had astigmatism. However, the prevalence of astigmatism has a great variation in different studies, ranging from 0.3% in Thailand<sup>83</sup> to 91.9% in Benin.<sup>44</sup> The use of a cylinder power >0.5 D as the cut point in our study limited this range. Although part of the variation can be due to differences in age groups, we observed this variation in both adults and children.

As mentioned earlier, the lowest and the highest prevalence of astigmatism in children was seen in South-East Asia and the Americas, respectively. However, according to Table 3, the Eastern Mediterranean and Western Pacific regions have the highest variation in the prevalence of astigmatism. One of the limitations of the studies conducted in the Eastern Mediterranean region is that most of them are from Iran,<sup>106–111,117,119–121,124–129,132,138,141,150,155,156,172–175</sup> which makes conclusion difficult, although a range of 6.6–51.4% for astigmatism in Iran is also noticeable.

The highest and the lowest prevalence of astigmatism was seen in American and African adults, respectively. However, the details of the results presented in tables and figures reject this finding. After the Americas, South-East Asia followed closely by the Western Pacific region had a high prevalence of astigmatism. The only eligible study for astigmatism analysis in the Americas was conducted on Chinese people living in the USA; therefore, it is in fact related to the Western Pacific region. Ethnic and racial differences may have a more prominent role in astigmatism in comparison with myopia and hyperopia.<sup>176</sup>

It seems that the eyelid and palpebral fissure shape in South-East Asian and some Western Pacific countries is the major cause of high astigmatism in these people.<sup>176</sup> A great part of the high prevalence of astigmatism in Western Pacific countries is due to the high prevalence of astigmatism in Chinese people.

The findings of this meta-analysis provide a new perspective of the status of refractive errors across the world based on the WHO classification.






Fig. 5. Forest plot of estimated pool prevalence (EPP) of hyperopia [spherical equivalent (SE) ≥ +2] in children by WHO regions.




Fig. 6. Forest plot of estimated pool prevalence (EPP) of hyperopia [spherical equivalent (SE) > +0.5] in adults by WHO regions.

As mentioned earlier, the prevalence of myopia, hyperopia, and astigmatism in children was lower in South-East Asian countries in comparison with other WHO regions while in adults, the highest prevalence of myopia was seen in South-

East Asia. On the other hand, the prevalence of hyperopia was high in both children and adults in American countries. Therefore, it seems that environmental factors may have a more important role in myopia since children are not myopic,






Fig. 7. Trend of hyperopia from 1990 to 2016.

and its prevalence is higher in adults in regions where near work is more common.<sup>7</sup> On the other hand, ethnic and genetic factors could have a more prominent role in hyperopia since the highest prevalence of hyperopia was seen in American children and adults.

The results of myopia and astigmatism in children and adults are interesting. The lowest and highest prevalence of myopia and astigmatism was seen in South-East Asian children and adults, respectively. It should be noted that common factors can cause myopia or astigmatism. The relationship between near work and myopia has been shown in different studies.<sup>177,178</sup> Some studies have reported that near work causes astigmatism due to incyclotorsion.<sup>179,180</sup> On the other hand, there are reports that 15-year-old children in some Asian countries spend more time on near work than their counterparts in some countries like UK and USA.<sup>7</sup> Therefore, it is possible that near work in this age group has caused astigmatism in non-astigmatic children due to incyclotorsion, manifesting the problems of astigmatism and myopia in adulthood. However, the role of ethnic, genetic, and environmental factors should be taken into account, as well.

Squinting can cause astigmatism, especially with the rule (WTR) astigmatism, in myopic patients and myopia in astigmatic patients.<sup>181</sup> Previous studies have shown the relationship between astigmatism and myopia.<sup>181–183</sup> However, the use of the SE in epidemiologic studies should not be overlooked. Considering the fact that astigmatism is part of the SE with a minus sign, this index is considered myopia in a spherically emmetropic individual due to a negative cylinder power. Therefore, part of this relationship can be due to the use of SE.

Our results showed that the prevalence of myopia had an increasing trend in the past three decades. Many studies have reported that myopia is becoming an epidemic, especially in East Asian countries, but few meta-analyses have confirmed this finding. The results of our meta-regression confirmed the increasing trend of myopia. Different reasons can be mentioned for the increase in the prevalence of myopia worldwide, including lifestyle changes and ever-increasing use of the computer and computer-related systems resulting in

increased near work. Different studies have evaluated the mechanism of developing myopia following near work. Increased lens thickness and the pressure of the ciliary muscle on the globe wall increase the axial length (AL) during accommodation. Some researchers believe that optical changes during accommodation (increased accommodative lag or increased higher order aberration) can change the choroidal thickness, resulting in AL changes during near work.<sup>184</sup> However, it should also be noted that myopic patients are more interested in near work. In addition to the effect of near work, more and more people use computers for their daily activities as a result of computerization, and are not therefore engaged in outdoor activities.<sup>177,181,185</sup> The hypothesis of the effect of outdoor activity on myopia has been tested in many studies.<sup>178,185</sup> A recent clinical trial showed that the incidence of myopia was about 10% lower in children who were engaged in outdoor activities.<sup>185</sup> Other studies have confirmed this finding as well.<sup>178,186</sup> According to this hypothesis, the factor that prevents myopia in outdoor activity is light. Some studies<sup>187,188</sup> have shown the role of intense light in the prevention of myopia formation. The mechanism is that light stimulates the secretion of dopamine in the retina which in turn prevents ocular elongation during the process of ocular development and prevents myopia. Finally, the age cohort effect should not be forgotten.

According to our results, astigmatism and hyperopia did not have a significant trend in the past three decades. As mentioned earlier, although the trend of hyperopia was not significant, the prevalence of hyperopia had a decreasing trend from 1990 to 2016 with a regression coefficient similar to myopia. First, we believe that the non-significant trend of hyperopia during these three decades may be due to the lower number of studies in hyperopia analysis. Second, more variation in the results of hyperopia versus myopia during the three decades may play a role the non-significant trend of hyperopia. However, the results of this meta-analysis propose the hypothesis that the decrease in the prevalence of hyperopia may be due to the increase in the prevalence of myopia in these years.

Considering the stability of the trend of astigmatism, although an increase was expected in its trend as in myopia, it seems that the role of outdoor activity in myopia is more prominent than near work because near work was expected to have a similar effect on the trend of astigmatism as well.

Lack of studies in many countries and lack of studies in each year in many countries were among the limitations of our study. Many studies were not included in the final analysis because they used different criteria for the detection of refractive errors or because we only analyzed the studies published in English. An important limitation of many studies was that they did not use cycloplegic refraction in children which caused us limitations in the analysis of refractive errors in individuals under 20 years of age. Although we tried to include studies with similar criteria in the analysis, these exclusion criteria may have biased the results. We did not evaluate different categories of refractive errors as low, moderate, or high myopia or hyperopia. Although there was great




Fig. 8. Forest plot of estimated pool prevalence (EPP) of astigmatism (cylinder power >0.5) in children by WHO regions.




Fig. 9. Forest plot of estimated pool prevalence (EPP) of astigmatism (cylinder power >0.5) in adults by WHO regions.

heterogeneity in the results of the studies, we tried to address the differences among studies through subgroup analysis and using a random effects model. Despite the above limitations, this is the first study to show the overall prevalence of refractive errors according to WHO regions regardless of any categorization, which can be considered the most important advantage of the study.

In conclusion, this meta-analysis showed the prevalence of myopia, hyperopia, and astigmatism in children and adults separately according to WHO regions for the first time. The results showed that astigmatism, hyperopia, and myopia were the most common refractive errors in children and adults in the mentioned order. Children in South-East Asia had the lowest prevalence of astigmatism, hyperopia, and myopia as compared to other WHO regions, while the highest prevalence of myopia and astigmatism was seen in South-East Asian adults. The highest prevalence of hyperopia in

children and adults was seen in the Americas. A direct correlation was found between the prevalence of myopia and astigmatism in most WHO regions. The trend of myopia has increased linearly in the past three decades, maybe as a result of increased indoor activity due to computerization in recent years.

#### Acknowledgements

The authors wish to thank Dr. Saman Mohazzab-Torabi (S.M.), Ms. Frida Jabbari-Azad (F.J.) and Ms. Mojgan Pakbin (M.P.) for their help with the literature review.

#### References

1. Pascolini D, Mariotti SP. Global estimates of visual impairment: 2010. *Br J Ophthalmol.* 2012;96(5):614–618.



2. Naidoo KS, Leasher J, Bourne RR, et al. Global vision impairment and blindness due to uncorrected refractive error, 1990–2010. *Optom Vis Sci*. 2016;93(3):227–234.
3. Fricke TR, Holden BA, Wilson DA, et al. Global cost of correcting vision impairment from uncorrected refractive error. *Bull World Health Organ*. 2012;90(10):728–738.
4. Smith TS, Frick KD, Holden BA, Fricke TR, Naidoo KS. Potential lost productivity resulting from the global burden of uncorrected refractive error. *Bull World Health Organ*. 2009;87(6):431–437.
5. Pan CW, Dirani M, Cheng CY, Wong TY, Saw SM. The age-specific prevalence of myopia in Asia: a meta-analysis. *Optom Vis Sci*. 2015;92(3):258–266.
6. Rudnicka AR, Kapetanakis VV, Wathern AK, et al. Global variations and time trends in the prevalence of childhood myopia, a systematic review and quantitative meta-analysis: implications for aetiology and early prevention. *Br J Ophthalmol*. 2016;100(7):882–890.
7. Dolgin E. The myopia boom. *Nature*. 2015;519(7543):276–278.
8. Foster PJ, Jiang Y. Epidemiology of myopia. *Eye*. 2014;28(2):202–208.
9. Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Ann Intern Med*. 2009;151(4):264–269. W264.
10. Hendler K, Mehravaran S, Lu X, Brown SI, Mondino BJ, Coleman AL. Refractive errors and amblyopia in the UCLA preschool vision program; first year results. *Am J Ophthalmol*. 2016;172:80–86.
11. Li H, Li S, Liu L, et al. Distribution of refraction and ocular biometric parameters in a population of junior middle school children in Anyang of Henan province. *Zhonghua Yi Xue Za Zhi*. 2014;94(17):1284–1288.
12. Kinge B, Midelfart A, Jacobsen G. Refractive errors among young adults and university students in Norway. *Acta Ophthalmol Scand*. 1998;76(6):692–695.
13. Zhu D, Wang Y, Yang X, et al. Pre- and postcycloplegic refractions in children and adolescents. *PLoS One*. 2016;11(12):e0167628.
14. Varma R, Torres M, McKean-Cowdin R, Rong F, Hsu C, Jiang X. Prevalence and risk factors for refractive error in adult Chinese Americans: the Chinese American Eye Study. *Am J Ophthalmol*. 2017;175:201–212.
15. Rim TH, Kim SH, Lim KH, Choi M, Kim HY, Baek SH. Refractive errors in Koreans: the Korea National Health and Nutrition Examination Survey 2008–2012. *Korean J Ophthalmol*. 2016;30(3):214–224.
16. Bin G, Liu H, Zhao C, et al. Refractive errors in northern China between the residents with drinking water containing excessive fluorine and normal drinking water. *Biol Trace Elem Res*. 2016;173(2):259–267.
17. Wu SY, Nemesure B, Leske MC. Refractive errors in a black adult population: the Barbados Eye Study. *Invest Ophthalmol Vis Sci*. 1999;40(10):2179–2184.
18. Rodriguez NM, Romero AF. The prevalence of refractive conditions in Puerto Rican adults attending an eye clinic system. *J Optom*. 2014;7(3):161–167.
19. Hendricks TJ, de Brabander J, Vankan-Hendricks MH, van der Horst FG, Hendrikse F, Knottnerus JA. Prevalence of habitual refractive errors and anisometropia among Dutch schoolchildren and hospital employees. *Acta Ophthalmol*. 2009;87(5):538–543.
20. Bourne RR, Dineen BP, Ali SM, Noorul Huq DM, Johnson GJ. Prevalence of refractive error in Bangladeshi adults: results of the National Blindness and Low Vision Survey of Bangladesh. *Ophthalmology*. 2004;111(6):1150–1160.
21. Rani PK, Raman R, Rachapalli SR, Kulothungan V, Kumaramanickavel G, Sharma T. Prevalence of refractive errors and associated risk factors in subjects with type 2 diabetes mellitus SN-DREAMS, report 18. *Ophthalmology*. 2010;117(6):1155–1162.
22. Durkin SR, Tan EW, Casson RJ, Selva D, Newland HS. Distance refractive error among aboriginal people attending eye clinics in remote South Australia. *Clin Exp Ophthalmol*. 2007;35(7):621–626.
23. Dandona R, Dandona L, Srinivas M, Giridhar P, McCarty CA, Rao GN. Population-based assessment of refractive error in India: the Andhra Pradesh eye disease study. *Clin Exp Ophthalmol*. 2002;30(2):84–93.
24. Prema R, George R, Sathyamangalam Ve R, et al. Comparison of refractive errors and factors associated with spectacle use in a rural and urban South Indian population. *Indian J Ophthalmol*. 2008;56(2):139–144.
25. Ma Y, Qu X, Zhu X, et al. Age-specific prevalence of visual impairment and refractive error in children aged 3–10 years in Shanghai, China. *Invest Ophthalmol Vis Sci*. 2016;57(14):6188–6196.
26. Wen G, Tarczy-Hornoch K, McKean-Cowdin R, et al. Prevalence of myopia, hyperopia, and astigmatism in non-Hispanic white and Asian children: multi-ethnic pediatric eye disease study. *Ophthalmology*. 2013;120(10):2109–2116.
27. Multi-Ethnic Pediatric Eye Disease Study Group. Prevalence of myopia and hyperopia in 6- to 72-month-old African American and Hispanic children: the multi-ethnic pediatric eye disease study. *Ophthalmology*. 2010;117(1):140–147 e143.
28. Ip JM, Robaei D, Kifley A, Wang JJ, Rose KA, Mitchell P. Prevalence of hyperopia and associations with eye findings in 6- and 12-year-olds. *Ophthalmology*. 2008;115(4):678–685 e671.
29. Castagno VD, Fassa AG, Vilela MA, Meucci RD, Resende DP. Moderate hyperopia prevalence and associated factors among elementary school students. *Cien Saude Colet*. 2015;20(5):1449–1458.
30. Dandona R, Dandona L, Srinivas M, et al. Refractive error in children in a rural population in India. *Invest Ophthalmol Vis Sci*. 2002;43(3):615–622.
31. Zhao J, Pan X, Sui R, Munoz SR, Sperduto RD, Ellwein LB. Refractive error study in children: results from Shunyi district, China. *Am J Ophthalmol*. 2000;129(4):427–435.
32. Goh PP, Abqariyah Y, Pokharell GP, Ellwein LB. Refractive error and visual impairment in school-age children in Gombak District, Malaysia. *Ophthalmology*. 2005;112(4):678–685.
33. Li SM, Liu LR, Li SY, et al. Design, methodology and baseline data of a school-based cohort study in Central China: the Anyang Childhood Eye Study. *Ophthalmic Epidemiol*. 2013;20(6):348–359.
34. Uzma N, Kumar BS, Khaja Mohinuddin Salar BM, Zafar MA, Reddy VD. A comparative clinical survey of the prevalence of refractive errors and eye diseases in urban and rural school children. *Can J Ophthalmol*. 2009;44(3):328–333.
35. Pai AS, Wang JJ, Samarawickrama C, et al. Prevalence and risk factors for visual impairment in preschool children the Sydney paediatric eye disease study. *Ophthalmology*. 2011;118(8):1495–1500.
36. Mashige KP, Jaggernath J, Ramson P, Martin C, Chinanayi FS, Naidoo KS. Prevalence of refractive errors in the INK area, Durban, South Africa. *Optom Vis Sci*. 2016;93(3):243–250.
37. Soler M, Anera RG, Castro JJ, Jiménez R, Jiménez JR. Prevalence of refractive errors in children in Equatorial Guinea. *Optom Vis Sci*. 2015;92(1):53–58.
38. Semanyenzi SE, Karimurio J, Nzayirambaho M. Prevalence and pattern of refractive errors in high schools of Nyarugenge district. *Rwanda Med J*. 2015;72(3):8–13.
39. Mehari ZA, Yimer AW. Prevalence of refractive errors among school-children in rural central Ethiopia. *Clin Exp Optom*. 2013;96(1):65–69.
40. Kumah BD, Ebri A, Abdul-Kabir M, et al. Refractive error and visual impairment in private school children in Ghana. *Optom Vis Sci*. 2013;90(12):1456–1461.
41. Bastawrous A, Mathenge W, Foster A, Kuper H. Prevalence and predictors of refractive error and spectacle coverage in Nakuru, Kenya: a cross-sectional, population-based study. *Int Ophthalmol*. 2013;33(5):541–548.
42. Ezelum C, Razavi H, Sivasubramaniam S, et al. Refractive error in Nigerian adults: prevalence, type, and spectacle coverage. *Invest Ophthalmol Vis Sci*. 2011;52(8):5449–5456.
43. Anera RG, Soler M, de la Cruz Cardona J, Salas C, Ortiz C. Prevalence of refractive errors in school-age children in Morocco. *Clin Exp Ophthalmol*. 2009;37(2):191–196.
44. Sounouvou I, Tchabi S, Doutetien C, Sonon F, Yehouessi L, Bassabi SK. A study of refractive errors in a primary school in Cotonou, Benin. *J Fr Ophthalmol*. 2008;31(8):771–775.
45. Naidoo KS, Raghunandan A, Mashige KP, et al. Refractive error and visual impairment in African children in South Africa. *Invest Ophthalmol Vis Sci*. 2003;44(9):3764–3770.



46. Kawuma M, Mayeku R. A survey of the prevalence of refractive errors among children in lower primary schools in Kampala district. *Afr Health Sci.* 2002;2(2):69–72.
47. Shiferaw Alemu D, Desalegn Gudeta A, Tsega Ferede A, Woretaw Alemu H. Prevalence and degrees of myopia and hyperopia at Gondar University Hospital Tertiary Eye Care and Training Center, Northwest Ethiopia. *Clin Optom.* 2016;8:85–91.
48. Sewunet SA, Aredo KK, Gedefew M. Uncorrected refractive error and associated factors among primary school children in Debre Markos District, Northwest Ethiopia. *BMC Ophthalmol.* 2014;14:95.
49. Yared AW, Belaynew WT, Destaye S, Ayanaw T, Zelalem E. Prevalence of refractive errors among school children in Gondar town, northwest Ethiopia. *Middle East Afr J Ophthalmol.* 2012;19(4):372–376.
50. Ferraz FH, Corrente JE, Opromolla P, Padovani CR, Schellini SA. Refractive errors in a Brazilian population: age and sex distribution. *Ophthalmic Physiol Opt.* 2015;35(1):19–27.
51. Schellini SA, Durkin SR, Hoyama E, et al. Prevalence of refractive errors in a Brazilian population: the Botucatu eye study. *Ophthalmic Epidemiol.* 2009;16(2):90–97.
52. Estacia P, Stramari LM, Schuch SB, Negrello D, Donato L. Prevalence of refractive errors in first grade school children of elementary schools of Northeast region of the Rio Grande do Sul State, Brazil. *Rev Bras Oftalmol.* 2007;66(5):297–303.
53. Garcia-Lievanos O, Sanchez-Gonzalez L, Espinosa-Cruz N, Hernandez-Flores LA, Salmeron-Leal L, Torres-Rodriguez HD. Myopia in school-children in a rural community in the State of Mexico, Mexico. *Clin Optom.* 2016;8:53–56.
54. De Amorim Garcia CA, Oréfice F, Dutra Nobre GF, De Brito Souza D, Ramalho Rocha ML, Garrido Vianna RN. Prevalence of refractive errors in students in Northeastern Brazil. *Arq Bras Oftalmol.* 2005;68(3):321–325.
55. Maul E, Barroso S, Munoz SR, Sperduto RD, Ellwein LB. Refractive error study in children: results from La Florida, Chile. *Am J Ophthalmol.* 2000;129(4):445–454.
56. Wang Q, Klein BE, Klein R, Moss SE. Refractive status in the Beaver Dam Eye Study. *Invest Ophthalmol Vis Sci.* 1994;35(13):4344–4347.
57. Haronian E, Wheeler NC, Lee DA. Prevalence of eye disorders among the elderly in Los Angeles. *Arch Gerontol Geriatr.* 1993;17(1):25–36.
58. Pi LH, Chen L, Liu Q, et al. Refractive status and prevalence of refractive errors in suburban school-age children. *Int J Med Sci.* 2010;7(6):342–353.
59. Hashim SE, Tan HK, Wan-Hazabbah WH, Ibrahim M. Prevalence of refractive error in Malay primary school children in suburban area of Kota Bharu, Kelantan, Malaysia. *Ann Acad Med Singap.* 2008;37(11):940–946.
60. Quek TP, Chua CG, Chong CS, et al. Prevalence of refractive errors in teenage high school students in Singapore. *Ophthalmic Physiol Opt.* 2004;24(1):47–55.
61. Congdon N, Wang Y, Song Y, et al. Visual disability, visual function, and myopia among rural Chinese secondary school children: the Xichang Pediatric Refractive Error Study (X-PRES) – report 1. *Invest Ophthalmol Vis Sci.* 2008;49(7):2888–2894.
62. Lan W, Zhao F, Lin L, et al. Refractive errors in 3–6 year-old Chinese children: a very low prevalence of myopia? *PLoS One.* 2013;8(10):e78003.
63. Pokharel A, Pokharel PK, Das H, Adhikari S. The patterns of refractive errors among the school children of rural and urban settings in Nepal. *Nepal J Ophthalmol Biannu Peer Rev Acad J Nepal Ophthalmic Soc NEPJOPH.* 2010;2(2):114–120.
64. Padhye AS, Khandekar R, Dharmadhikari S, Dole K, Gogate P, Deshpande M. Prevalence of uncorrected refractive error and other eye problems among urban and rural school children. *Middle East Afr J Ophthalmol.* 2009;16(2):69–74.
65. Gao Z, Meng N, Muecke J, et al. Refractive error in school children in an urban and rural setting in Cambodia. *Ophthalmic Epidemiol.* 2012;19(1):16–22.
66. Wong TY, Foster PJ, Hee J, et al. Prevalence and risk factors for refractive errors in adult Chinese in Singapore. *Invest Ophthalmol Vis Sci.* 2000;41(9):2486–2494.
67. Gupta A, Casson RJ, Newland HS, et al. Prevalence of refractive error in rural Myanmar: the Meiktila eye study. *Ophthalmology.* 2008;115(1):26–32.
68. Saw SM, Gazzard G, Koh D, et al. Prevalence rates of refractive errors in Sumatra, Indonesia. *Invest Ophthalmol Vis Sci.* 2002;43(10):3174–3180.
69. Sawada A, Tomidokoro A, Araie M, Iwase A, Yamamoto T. Refractive errors in an elderly Japanese population: the Tajimi study. *Ophthalmology.* 2008;115(2):363–370 e363.
70. Krishnaiah S, Srinivas M, Khanna RC, Rao GN. Prevalence and risk factors for refractive errors in the South Indian adult population: the Andhra Pradesh Eye disease study. *Clin Ophthalmol.* 2009;3:17–27.
71. Kim EC, Morgan IG, Kakizaki H, Kang S, Jee D. Prevalence and risk factors for refractive errors: Korean National Health and Nutrition Examination survey 2008–2011. *PLoS One.* 2013;8(11):e80361.
72. Jang JU, Park IJ. The status of refractive errors in elementary school children in South Jeolla Province, South Korea. *Clin Optom.* 2015;7:45–51.
73. Wang X, Liu D, Feng R, Zhao H, Wang Q. Refractive error among urban preschool children in Xuzhou, China. *Int J Clin Exp Pathol.* 2014;7(12):8922–8928.
74. Paudel P, Ramson P, Naduvilath T, et al. Prevalence of vision impairment and refractive error in school children in Ba Ria – Vung Tau province, Vietnam. *Clin Exp Ophthalmol.* 2014;42(3):217–226.
75. Li Z, Xu K, Wu S, et al. Population-based survey of refractive error among school-aged children in rural northern China: the Heilongjiang eye study. *Clin Exp Ophthalmol.* 2014;42(4):379–384.
76. Yoo YC, Kim JM, Park KH, Kim CY, Kim TW. Refractive errors in a rural Korean adult population: the Namil Study. *Eye (Lond).* 2013;27(12):1368–1375.
77. Adhikari S, Nepal B, Shrestha J, Khandekar R. Magnitude and determinants of refractive error among school children of two districts of Kathmandu, Nepal. *Oman J Ophthalmol.* 2013;6(3):175–178.
78. Nangia V, Jonas JB, Sinha A, Gupta R, Bhojwani K. Prevalence of undercorrection of refractive error in rural Central India: the Central India eye and medical study. *Acta Ophthalmol.* 2012;90(2):e166–e167.
79. Casson RJ, Kahawita S, Kong A, Muecke J, Sisaleumsak S, Visonnavong V. Exceptionally low prevalence of refractive error and visual impairment in schoolchildren from Lao People's Democratic Republic. *Ophthalmology.* 2012;119(10):2021–2027.
80. Wei RH, Li XR, Zhao SZ, et al. Prevalence of refractive error in a rural minority population in southwestern China. *Int J Ophthalmol.* 2011;11(6):1031–1034.
81. Tan CSH, Chan YH, Wong TY, et al. Prevalence and risk factors for refractive errors and ocular biometry parameters in an elderly Asian population: the Singapore Longitudinal Aging Study (SLAS). *Eye.* 2011;25(10):1294–1301.
82. Pan CW, Wong TY, Lavanya R, et al. Prevalence and risk factors for refractive errors in Indians: the Singapore Indian Eye Study (SINDI). *Invest Ophthalmol Vis Sci.* 2011;52(6):3166–3173.
83. Yingyong P. Refractive errors survey in primary school children (6–12 year old) in 2 provinces: Bangkok and Nakhonpathom (one year result). *J Med Assoc Thai.* 2010;93(10):1205–1210.
84. Li Z, Sun D, Cui H, et al. Refractive error among the elderly in rural Southern Harbin, China. *Ophthalmic Epidemiol.* 2009;16(6):388–394.
85. Saw SM, Chan YH, Wong WL, et al. Prevalence and risk factors for refractive errors in the Singapore Malay Eye Survey. *Ophthalmology.* 2008;115(10):1713–1719.
86. He M, Zeng J, Liu Y, Xu J, Pokharel GP, Ellwein LB. Refractive error and visual impairment in urban children in southern China. *Invest Ophthalmol Vis Sci.* 2004;45(3):793–799.
87. Sun Y, Cao H, Yan ZG. Prevalence of refractive errors in middle school students in Lanzhou city. *Int J Ophthalmol.* 2007;7(5):1240–1242.
88. Li JJ, Xu L, Li JJ, Cui TT, Yang H. Prevalence of refractive error in defined populations in rural and urban areas in Beijing. *Ophthalmol China.* 2007;16(3):206–211.

89. He M, Huang W, Zheng Y, Huang L, Ellwein LB. Refractive error and visual impairment in school children in rural southern China. *Ophthalmology*. 2007;114(2):374–382.e371.
90. Khan AA, Nasti AR, Ayoub Dar M, Lone SA. Prevalence of refractive errors in school children. *JK Pract*. 2005;12(3):156–159.
91. Raju P, Ramesh SV, Arvind H, et al. Prevalence of refractive errors in a rural South Indian population. *Invest Ophthalmol Vis Sci*. 2004;45(12):4268–4272.
92. Murthy GV, Gupta SK, Ellwein LB, et al. Refractive error in children in an urban population in New Delhi. *Invest Ophthalmol Vis Sci*. 2002;43(3):623–631.
93. Pokharel GP, Negrel AD, Munoz SR, Ellwein LB. Refractive error study in children: results from Mechi zone, Nepal. *Am J Ophthalmol*. 2000;129(4):436–444.
94. Czepita D, Mojsa A, Ustianowska M, Czepita M, Lachowicz E. Prevalence of refractive errors in schoolchildren ranging from 6 to 18 years of age. *Ann Acad Med Stetin*. 2007;53(1):53–56.
95. Czepita D, Zejmo M, Mojsa A. Prevalence of myopia and hyperopia in a population of Polish schoolchildren. *Ophthalmic Physiol Opt*. 2007;27(1):60–65.
96. Gronlund MA, Andersson S, Aring E, Hard AL, Hellstrom A. Ophthalmological findings in a sample of Swedish children aged 4–15 years. *Acta Ophthalmol Scand*. 2006;84(2):169–176.
97. Williams KM, Verhoeven VJM, Cumberland P, et al. Prevalence of refractive error in Europe: the European Eye Epidemiology (E3) Consortium. *Eur J Epidemiol*. 2015;30(4):305–315.
98. Sherwin JC, Khawaja AP, Broadway D, et al. Uncorrected refractive error in older British adults: the EPIC-Norfolk Eye Study. *Br J Ophthalmol*. 2012;96(7):991–996.
99. Wolfram C, Hohn R, Kottler U, et al. Prevalence of refractive errors in the European adult population: the Gutenberg Health Study (GHS). *Br J Ophthalmol*. 2014;98(7):857–861.
100. Anton A, Andrada MT, Mayo A, Portela J, Merayo J. Epidemiology of refractive errors in an adult European population: the Segovia study. *Ophthalmic Epidemiol*. 2009;16(4):231–237.
101. Pateras E. Prevalence of refractive errors amongst adults, located at the north suburbs of Athens-Greece. *Health Sci J*. 2012;6(1):102–114.
102. Villarreal MG, Ohlsson J, Abrahamsson M, Sjöström A, Sjöstrand J. Myopisation: the refractive tendency in teenagers. Prevalence of myopia among young teenagers in Sweden. *Acta Ophthalmol Scand*. 2000;78(2):177–181.
103. Caca I, Cingu AK, Sahin A, et al. Amblyopia and refractive errors among school-aged children with low socioeconomic status in southeastern Turkey. *J Pediatr Ophthalmol Strabismus*. 2013;50(1):37–43.
104. Gull A, Raza A. Visual screening and refractive errors among school aged children. *J Rawalpindi Med Coll (JRMC)*. 2014;18:97–100.
105. Gursoy H, Basmak H, Yaz Y, Colak E. Vision screening in children entering school: Eskisehir, Turkey. *Ophthalmic Epidemiol*. 2013;20(4):232–238.
106. Yekta AA, Fotouhi A, Khabazkhoob M, et al. The prevalence of refractive errors and its determinants in the elderly population of Mashhad, Iran. *Ophthalmic Epidemiol*. 2009;16(3):198–203.
107. Yekta A, Fotouhi A, Hashemi H, et al. Prevalence of refractive errors among schoolchildren in Shiraz, Iran. *Clin Exp Ophthalmol*. 2010;38(3):242–248.
108. Hashemi H, Rezvan F, Ostadimoghaddam H, Abdollahi M, Hashemi M, Khabazkhoob M. High prevalence of refractive errors in a rural population: 'Nooravaran Salamat' Mobile Eye Clinic experience. *Clin Exp Ophthalmol*. 2013;41(7):635–643.
109. Rezvan F, Khabazkhoob M, Fotouhi A, et al. Prevalence of refractive errors among school children in Northeastern Iran. *Ophthalmic Physiol Opt*. 2012;32(1):25–30.
110. Yekta A, Hashemi H, Ostadimoghaddam H, Shafae S, Norouzirad R, Khabazkhoob M. Prevalence of refractive errors among the elderly population of Sari, Iran. *Iran J Ophthalmol*. 2013;25(2):123–132.
111. Ziaei H, Katibeh M, Solaimanizad R, et al. Prevalence of refractive errors; the Yazd eye study. *J Ophthalmic Vis Res*. 2013;8(3):227–236.
112. Bataineh HA, Khatatbeh AE. Prevalence of refractive errors in school children of Tafila City. *Rawal Med J*. 2008;33(1):85–87.
113. Mullen EAH, Gammoh Y, Al-Bdour M, Sayegh FN. Refractive error and ocular biometry in Jordanian adults. *Ophthalmic Physiol Opt*. 2005;25(4):302–309.
114. Al-Rowaily MA. Prevalence of refractive errors among pre-school children at King Abdulaziz Medical City, Riyadh, Saudi Arabia. *Saudi J Ophthalmol*. 2010;24(2):45–48.
115. Rowaily MAA, Alanizi BM. Prevalence of uncorrected refractive errors among adolescents at King Abdul-Aziz medical city, Riyadh, Saudi Arabia. *J Clin Exp Ophthalmol*. 2010;1:114.
116. Abdullah AS, Jadoon MZ, Akram M, et al. Prevalence of uncorrected refractive errors in adults aged 30 years and above in a rural population in Pakistan. *J Ayub Med Coll Abbottabad*. 2015;27(1):8–12.
117. Hashemi H, Yekta A, Jafarzadehpur E, et al. High prevalence of refractive errors in 7 year old children in Iran. *Iran J Public Health*. 2016;45(2):194–202.
118. Hameed A. Screening for refractive errors and visual impairment among school children in Kohat, Pakistan. *Rawal Med J*. 2016;41(4):437–440.
119. Rajavi Z, Sabbaghi H, Baghini AS, et al. Prevalence of amblyopia and refractive errors among primary school children. *J Ophthalmic Vis Res*. 2015;10(4):408–416.
120. Norouzirad R, Hashemi H, Yekta A, et al. The prevalence of refractive errors in 6- to 15-year-old schoolchildren in Dezful, Iran. *J Curr Ophthalmol*. 2015;27(1-2):51–55.
121. Hashemi H, Asgari S, Yekta AA, et al. Prevalence of astigmatism in 4- to 6-year-old population of Mashhad, Iran. *J Compr Ped*. 2015;6(1):e25277.
122. Atta Z, Arif AS, Ahmed I, Farooq U. Prevalence of refractive errors in Madrasa students of Haripur district. *J Ayub Med Coll Abbottabad*. 2015;27(4):850–852.
123. Latif MA, Latif MZ, Hussain I, Nizami R, Jamal K. Prevalence of refractive errors in a public school children of Lahore. *Pak J Med Health Sci*. 2014;8(4):1016–1019.
124. Hashemi H, Rezvan F, Yekta AA, Hashemi M, Norouzirad R, Khabazkhoob M. The prevalence of astigmatism and its determinants in a rural population of Iran: the "Nooravaran Salamat" mobile eye clinic experience. *Middle East Afr J Ophthalmol*. 2014;21(2):175–181.
125. Hashemi H, Rezvan F, Beiranvand A, et al. Prevalence of refractive errors among high school students in western Iran. *J Ophthalmic Vis Res*. 2014;9(2):232–239.
126. Hashemi H, Khabazkhoob M, Yazdani N, et al. The prevalence of refractive errors among Iranian university students. *Iran J Ophthalmol*. 2014;26(3):129–135.
127. Hashemi H, Khabazkhoob M, Yekta A, et al. High prevalence of astigmatism in the 40- to 64-year-old population of Shahroud, Iran. *Clin Exp Ophthalmol*. 2012;40(3):247–254.
128. Ostadimoghaddam H, Fotouhi A, Hashemi H, et al. Prevalence of the refractive errors by age and gender: the Mashhad eye study of Iran. *Clin Exp Ophthalmol*. 2011;39(8):743–751.
129. Khalaj M, Gasemi M, Zeidi IM. Prevalence of refractive errors in primary school children [7–15 years] of Qazvin City. *Eur J Sci Res*. 2009;28(2):174–185.
130. He M, Huang W, Li Y, Zheng Y, Yin Q, Foster PJ. Refractive error and biometry in older Chinese adults: the Liwan eye study. *Invest Ophthalmol Vis Sci*. 2009;50(11):5130–5136.
131. Shah SP, Jadoon MZ, Dineen B, et al. Refractive errors in the adult Pakistani population: the national blindness and visual impairment survey. *Ophthalmic Epidemiol*. 2008;15(3):183–190.
132. Fotouhi A, Hashemi H, Khabazkhoob M, Mohammad K. The prevalence of refractive errors among schoolchildren in Dezful, Iran. *Br J Ophthalmol*. 2007;91(3):287–292.
133. Das A, Dutta H, Bhaduri G, De Ajay S, Sarkar K, Bannerjee M. A study on refractive errors among school children in Kolkata. *J Indian Med Assoc*. 2007;105(4):169–172.
134. Junghans BM, Crewther SG. Little evidence for an epidemic of myopia in Australian primary school children over the last 30 years. *BMC Ophthalmol*. 2005;5:1.

135. Junghans BM, Crewther SG. Prevalence of myopia among primary school children in eastern Sydney. *Clin Exp Optom*. 2003;86(5):339–345.
136. Attebo K, Ivers RQ, Mitchell P. Refractive errors in an older population: the Blue Mountains Eye Study. *Ophthalmology*. 1999;106(6):1066–1072.
137. Pan CW, Zheng YF, Anuar AR, et al. Prevalence of refractive errors in a multiethnic Asian population: the Singapore epidemiology of eye disease study. *Invest Ophthalmol Vis Sci*. 2013;54(4):2590–2598.
138. Hashemi H, Khabazkhoob M, Jafarzadehpur E, et al. High prevalence of myopia in an adult population, Shahroud, Iran. *Optom Vis Sci*. 2012;89(7):993–999.
139. Cortinez MF, Chiappe JP, Iribarren R. Prevalence of refractive errors in a population of office-workers in Buenos Aires, Argentina. *Ophthalmic Epidemiol*. 2008;15(1):10–16.
140. Liang YB, Wong TY, Sun LP, et al. Refractive errors in a rural Chinese adult population the Handan eye study. *Ophthalmology*. 2009;116(11):2119–2127.
141. Hashemi H, Fotouhi A, Mohammad K. The age- and gender-specific prevalences of refractive errors in Tehran: the Tehran Eye Study. *Ophthalmic Epidemiol*. 2004;11(3):213–225.
142. Xu L, Li J, Cui T, et al. Refractive error in urban and rural adult Chinese in Beijing. *Ophthalmology*. 2005;112(10):1676–1683.
143. Cheng CY, Hsu WM, Liu JH, Tsai SY, Chou P. Refractive errors in an elderly Chinese population in Taiwan: the Shihpai Eye Study. *Invest Ophthalmol Vis Sci*. 2003;44(11):4630–4638.
144. Wickremasinghe S, Foster PJ, Uranchimeg D, et al. Ocular biometry and refraction in Mongolian adults. *Invest Ophthalmol Vis Sci*. 2004;45(3):776–783.
145. Wensor M, McCarty CA, Taylor HR. Prevalence and risk factors of myopia in Victoria, Australia. *Arch Ophthalmol*. 1999;117(5):658–663.
146. Nangia V, Jonas JB, Sinha A, Matin A, Kulkarni M. Refractive error in central India: the Central India Eye and Medical Study. *Ophthalmology*. 2010;117(4):693–699.
147. Landers J, Henderson T, Craig J. Prevalence and associations of refractive error in indigenous Australians within central Australia: the Central Australian Ocular Health Study. *Clin Exp Ophthalmol*. 2010;38(4):381–386.
148. Sherwin JC, Kelly J, Hewitt AW, Kearns LS, Griffiths LR, Mackey DA. Prevalence and predictors of refractive error in a genetically isolated population: the Norfolk Island Eye Study. *Clin Exp Ophthalmol*. 2011;39(8):734–742.
149. Pan CW, Klein BEK, Cotch MF, et al. Racial variations in the prevalence of refractive errors in the United States: the multi-ethnic study of atherosclerosis. *Am J Ophthalmol*. 2013;155(6):1129–1138.e1121.
150. Jamali P, Fotouhi A, Hashemi H, Younesian M, Jafari A. Refractive errors and amblyopia in children entering school: Shahrood, Iran. *Optom Vis Sci*. 2009;86(4):364–369.
151. Morgan A, Young R, Narankhand B, Chen S, Cottrill C, Hosking S. Prevalence rate of myopia in schoolchildren in rural Mongolia. *Optom Vis Sci*. 2006;83(1):53–56.
152. Pavithra M, Maheshwaran R, Sujatha R. A study on the prevalence of refractive errors among school children of 7–15 years age group in the field practice areas of a medical college in Bangalore. *Int J Med Sci Public Health*. 2013;2(3):641–645.
153. Villarreal GM, Ohlsson J, Cavazos H, Abrahamsson M, Mohamed JH. Prevalence of myopia among 12- to 13-year-old schoolchildren in Northern Mexico. *Optom Vis Sci*. 2003;80(5):369–373.
154. Aldebasi YH. Prevalence of correctable visual impairment in primary school children in Qassim Province, Saudi Arabia. *J Optom*. 2014;7(3):168–176.
155. Fotouhi A, Khabazkhoob M, Hashemi H, Yekta AA, Mohammad K. Importance of including refractive error tests in school children's vision screening. *Arch Iran Med*. 2011;14(4):250–253.
156. Dirani M, Chan YH, Gazzard G, et al. Prevalence of refractive error in Singaporean Chinese children: the Strabismus, Amblyopia, and Refractive Error in young Singaporean children (STARS) study. *Invest Ophthalmol Vis Sci*. 2010;51(3):1348–1355.
157. Vitale S, Ellwein L, Cotch MF, Ferris Iii FL, Sperduto R. Prevalence of refractive error in the United States, 1999–2004. *Arch Ophthalmol*. 2008;126(8):1111–1119.
158. Jobke S, Kasten E, Vorwerk C. The prevalence rates of refractive errors among children, adolescents, and adults in Germany. *Clin Ophthalmol*. 2008;2(3):601–607.
159. He MG, Lin Z, Huang J, Lu Y, Wu CF, Xu JJ. Population-based survey of refractive error in school-aged children in Liwan District, Guangzhou. *Zhonghua Yan Ke Za Zhi*. 2008;44(6):491–496.
160. Huynh SC, Kifley A, Rose KA, Morgan I, Heller GZ, Mitchell P. Astigmatism and its components in 6-year-old children. *Invest Ophthalmol Vis Sci*. 2006;47(1):55–64.
161. Logan NS, Davies LN, Mullen EA, Gilmartin B. Ametropia and ocular biometry in a U.K. university student population. *Optom Vis Sci*. 2005;82(4):261–266.
162. Woo WW, Lim KA, Yang H, et al. Refractive errors in medical students in Singapore. *Singap Med J*. 2004;45(10):470–474.
163. Shih YF, Hsiao CK, Tung YL, Lin LL, Chen CJ, Hung PT. The prevalence of astigmatism in Taiwan schoolchildren. *Optom Vis Sci*. 2004;81(2):94–98.
164. Tong L, Saw SM, Carkeet A, Chan WY, Wu HM, Tan D. Prevalence rates and epidemiological risk factors for astigmatism in Singapore school children. *Optom Vis Sci*. 2002;79(9):606–613.
165. Midelfart A, Kinge B, Midelfart S, Lydersen S. Prevalence of refractive errors in young and middle-aged adults in Norway. *Acta Ophthalmol Scand*. 2002;80(5):501–505.
166. Pensyl CD, Harrison RA, Simpson P, Waterbor JW. Distribution of astigmatism among Sioux Indians in South Dakota. *J Am Optom Assoc*. 1997;68(7):425–431.
167. Kalikivayi V, Naduvilath TJ, Bansal AK, Dandona L. Visual impairment in school children in southern India. *Indian J Ophthalmol*. 1997;45(2):129–134.
168. Lewallen S, Lowdon R, Courtright P, Mehl GL. A population-based survey of the prevalence of refractive error in Malawi. *Ophthalmic Epidemiol*. 1995;2(3):145–149.
169. Lam CS, Goh WS, Tang YK, Tsui KK, Wong WC, Man TC. Changes in refractive trends and optical components of Hong Kong Chinese aged over 40 years. *Ophthalmic Physiol Opt*. 1994;14(4):383–388.
170. Rai S, Thapa HB, Sharma MK, Dhakhwa K, Karki R. The distribution of refractive errors among children attending Lumbini Eye Institute, Nepal. *Nepal J Ophthalmol*. 2012;4(1):90–95.
171. Young FA, Leary GA, Baldwin WR, et al. The transmission of refractive errors within Eskimo families. *Am J Optom Arch Am Acad Optom*. 1969;46(9):676–685.
172. Hashemi H, Khabazkhoob M, Iribarren R, Emamian MH, Fotouhi A. Five-year change in refraction and its ocular components in the 40- to 64-year-old population of the Shahrood eye cohort study. *Clin Exp Ophthalmol*. 2016;44(8):669–677.
173. Ostadimoghaddam H, Mirhajian H, Yekta A, et al. Eye problems in children with hearing impairment. *J Curr Ophthalmol*. 2015;27(1-2):56–59.
174. Hashemi H, Yekta A, Nabovati P, Khoshhal F, Riazi A, Khabazkhoob M. The prevalence of refractive errors in 5–15 year-old population of two underserved rural areas of Iran. *J Curr Ophthalmol*. 2017.
175. Ostadi-Moghaddam H, Fotouhi A, Khabazkhoob M, Heravian J, Yekta AA. Prevalence and risk factors of refractive errors among schoolchildren in Mashhad, 2006–2007. *Iran J Ophthalmol*. 2008;20(3):3–9.
176. Read SA, Collins MJ, Carney LG. A review of astigmatism and its possible genesis. *Clin Exp Optom*. 2007;90(1):5–19.
177. Ramamurthy D, Lin Chua SY, Saw SM. A review of environmental risk factors for myopia during early life, childhood and adolescence. *Clin Exp Optom*. 2015;98(6):497–506.
178. Pan CW, Ramamurthy D, Saw SM. Worldwide prevalence and risk factors for myopia. *Ophthalmic Physiol Opt*. 2012;32(1):3–16.
179. Buehren T, Collins MJ, Loughridge J, Carney LG, Iskander DR. Corneal topography and accommodation. *Cornea*. 2003;22(4):311–316.
180. Yasuda A, Yamaguchi T. Steepening of corneal curvature with contraction of the ciliary muscle. *J Cataract Refract Surg*. 2005;31(6):1177–1181.

181. Hashemi H, Khabazkhoob M, Peyman A, et al. The association between residual astigmatism and refractive errors in a population-based study. *J Refract Surg.* 2013;29(9):624–628.
182. Heidary G, Ying GS, Maguire MG, Young TL. The association of astigmatism and spherical refractive error in a high myopia cohort. *Optom Vis Sci.* 2005;82(4):244–247.
183. Fotouhi A, Hashemi H, Yekta AA, Mohammad K, Khoob MK. Characteristics of astigmatism in a population of schoolchildren, Dezful, Iran. *Optom Vis Sci.* 2011;88(9):1054–1059.
184. Hashemi H, Dadbin N, Yekta A, et al. Relation between near work and ocular biometric components. *Int J Occup Hyg.* 2015;7(3):153–158.
185. He M, Xiang F, Zeng Y, et al. Effect of time spent outdoors at school on the development of myopia among children in China: a randomized clinical trial. *JAMA.* 2015;314(11):1142–1148.
186. Jin JX, Hua WJ, Jiang X, et al. Effect of outdoor activity on myopia onset and progression in school-aged children in northeast China: the Sujiatun Eye Care Study. *BMC Ophthalmol.* 2015;15:73.
187. Ashby R, Ohlendorf A, Schaeffel F. The effect of ambient illuminance on the development of deprivation myopia in chicks. *Invest Ophthalmol Vis Sci.* 2009;50(11):5348–5354.
188. Ashby RS, Schaeffel F. The effect of bright light on lens compensation in chicks. *Invest Ophthalmol Vis Sci.* 2010;51(10):5247–5253.