

REVIEW

Open Access

A review on the epidemiology of myopia in school children worldwide



Andrzej Grzybowski^{1,2*} , Piotr Kanclerz³, Kazuo Tsubota^{4,5}, Carla Lanca⁶ and Seang-Mei Saw^{6,7,8}

Abstract

Background: Due to high prevalence myopia has gained importance in epidemiological studies. Children with early onset are at particular risk of complications associated with myopia, as progression over time might result in high myopia and myopic macular degeneration. Both genetic and environmental factors play a role in the increasing prevalence of myopia. The aim of this study is to review the current literature on epidemiology and risk factors for myopia in school children (aged 6–19 years) around the world.

Main body: PubMed and Medline were searched for the following keywords: prevalence, incidence, myopia, refractive error, risk factors, children and visual impairment. English language articles published between Jan 2013 and Mar 2019 were included in the study. Studies were critically reviewed for study methodology and robustness of data. Eighty studies were included in this literature review.

Myopia prevalence remains higher in Asia (60%) compared with Europe (40%) using cycloplegic refraction examinations. Studies reporting on non-cycloplegic measurements show exceptionally high myopia prevalence rates in school children in East Asia (73%), and high rates in North America (42%). Low prevalence under 10% was described in African and South American children. In recent studies, risk factors for myopia in schoolchildren included low outdoor time and near work, dim light exposure, the use of LED lamps for homework, low sleeping hours, reading distance less than 25 cm and living in an urban environment.

Conclusion: Low levels of outdoor activity and near work are well-established risk factors for myopia; this review provides evidence on additional environmental risk factors. New epidemiological studies should be carried out on implementation of public health strategies to tackle and avoid myopia. As the myopia prevalence rates in non-cycloplegic studies are overestimated, we recommend considering only cycloplegic measurements.

Keywords: Myopia, Epidemiology, Risk factors, Children

Background

The modern rise in myopia mirrors a trend with children in many countries spending considerable amounts of time engaged in reading, studying or — more recently — using computer and smartphones. The evidence suggests that not only genetic, but also environmental factors such as time spent outdoors [1–4], play a major role in this rise, and probably explain the epidemic of myopia that has appeared in East Asia. In other parts of the world, the prevalence of myopia also seems to be

increasing. Therefore, myopia has gained particular importance in epidemiological studies. It is estimated that 1.4 billion people were myopic in 2000, and it is predicted that by 2050 the number will reach 4.8 billion [5]. Socioeconomically, refractive errors, particularly if uncorrected, can affect school performance, limit employability and impair quality of life. Myopia is known to be associated with several ocular complications such as retinal detachment, glaucoma, cataract, optic disk changes and maculopathy [6]. High prevalence rates pose a major public health challenge due to visual impairment. The global potential productivity loss associated with the burden of visual impairment in 2015 was estimated at US\$244 billion from uncorrected myopia, and US\$6 billion from myopic macular degeneration [7]. Children with early onset myopia are the group at major risk, as

* Correspondence: ae.grzybowski@gmail.com

¹Department of Ophthalmology, University of Warmia and Mazury, Olsztyn, Poland

²Foundation for Ophthalmology Development, Institute for Research in Ophthalmology, Górczyczewskiego 2/3, 60-554 Poznań, Poland
Full list of author information is available at the end of the article



they will have higher duration of the disease, higher myopia progression and will be at risk of developing high myopia plus myopic macular degeneration. Age of myopia onset or duration of myopia progression is the most significant prognosticator of high myopia in later childhood [8].

The aim of this study is to present a review on the current epidemiology and risk factors for myopia in school children aged 6–19 years.

Main body

Methodology

Literature search

PubMed and Medline were searched to identify the prevalence of myopia among children, as reported in articles between January 2013 and March 2019. The following keywords were used in various combinations: *prevalence, incidence, myopia, refractive error, and visual impairment* (“prevalence”[All Fields] OR “incidence”[All Fields]) AND (“refractive error”[All Fields] OR “myopia”[All Fields] OR “visual impairment”[All Fields]). All publications in English and abstracts from non-English publications were reviewed. The reference lists of relevant publications were also considered as a potential source of information. If other studies (e.g., older than 5 years) were essential to draw conclusions, they were included in the discussion section. Studies were critically reviewed for study methodology and robustness of data, particularly the myopia definition and measurements under cycloplegia. No attempts to discover unpublished data were made.

Study selection

Full-text articles included in the prevalence analysis were required to meet the following criteria: 1) a cross-sectional or cohort design 2) refractive error measurements taken with a refractometer 3) clear definition of myopia and information on cycloplegic or non-cycloplegic measurements 4) prevalence assessed in children aged 6–19 years 5) studies with a minimum sample of 100 children. If more than one definition of myopia was used in a study, the prevalence for the more commonly used one was selected in order to enable comparison. Results for up to two age-groups were presented, and if data for more than two cohorts were reported, the average for the study or the most common age-group was selected. Studies were excluded from the prevalence analysis if they presented self-reported near-sightedness, reported the prevalence of visual impairment (but not myopia) or included animals.

Results

The search identified 1627 unique articles. Twenty-eight articles fulfilled the criteria for being included in the main analysis (myopia prevalence). One study was excluded, as it presented data from primary care optometry clinics [9].

Additionally, 55 articles were included in the analysis of risk factors.

Prevalence of myopia in school children

The prevalence of myopia was determined by the spherical equivalent refraction (SER) calculated as sphere plus the half of the cylindrical error. The reported prevalence of myopia is shown in Table 1 (list of studies in Additional file 1), with a geographical and age breakdown in Fig. 1 (cycloplegic measurements) and Fig. 2 (non-cycloplegic measurements). The prevalence ranged from 0.7% in Saudi Arabia (children aged 3 to 10 years) [35], 1.4% in South America (children aged 5–15 years) [28] to 65.5% in a cohort of 3rd year junior high school students (age 14–15 years; mean 15.25 ± 0.46 years) in the Haidian district of Beijing. The highest prevalence of myopia in schoolchildren was reported in East Asia and Singapore, urban areas of China, Taiwan and South Korea [39, 40]. In Europe the prevalence rates reached 42.7% in a 10–19-year French cohort [24].

Compared with cycloplegic measurements the majority of the studies reporting on myopia prevalence with non-cycloplegic measurements reported much higher prevalence rates. For example, a prevalence of 73% was found in South Korean children aged 12 to 18 years old [33]. However, there are some countries where the prevalence rates remain low, such as Brazil (3.14 and 9.6%) [26, 27] and Ghana (3.4%) [25]. In these countries even when considering an overestimated non-cycloplegic measurements the prevalence of myopia in school children remains low e.g., in the Republic of South Africa (7%) [38] or in Colombia (11.2%) [25, 37]. Carter et al. also found a very low prevalence of myopia (with relatively common hyperopia) in indigenous schoolchildren from Paraguay (1.4%) [28].

A critical parameter for epidemiological analysis of myopia is age, as prevalence rates of myopia are known to increase significantly with age (Table 1). For example, in the Shandong Children Eye Study, only $1.76 \pm 1.2\%$ of four-year-old children had myopia, while at the age of 17 years the prevalence was $84.6 \pm 3.2\%$ [14]. In another study, the one-year incidence of myopia among grade 1 (age 6–7 years) Chinese students was 33.6% (95% CI: 31.7–35%), with a progression rate of -0.97 D (95% CI: -1.22 to -0.71 D) [41]. Moreover, myopia beginning at school continues to progress up to adulthood in almost half of the patients [42].

Change over time

In several countries the prevalence of myopia has increased in the last years. In a study from the Haidian District in Beijing, China, the prevalence of myopia in a cohort of 15-year-old schoolchildren increased from 55.95% in 2005 to 65.48% in 2015 [10]. In Fenghua city, eastern China, the prevalence of myopia in high school students increased from 79.5% in 2001 to


Table 1 Cross-sectional studies reporting the prevalence of myopia in school children

Study	Region	Country	Cycloplegia	Definition	Participants	Age range/cohort	Mean Age	Prevalence (95% CI)
Li et al. 2017 [10]	Asia (East)	China	Yes	SE < -0.5 D	37424	3rd year junior high school	15.25 ± 0.46	65.48% (N/A)
Pan et al. 2017 [11]	Asia (East)	China	Yes	SE < -0.5 D	2346	13–14 years (grade 7)	13.8 ± 0.8	29.5% (27.7–31.4%)
Guo et al. 2016 [12]	Asia (East)	China	Yes	SE ≤ -0.5 D	3055	primary and middle school (grades 1–9)	13.6 ± 1.6	47.4% (45.6–49.2%)
Zhou et al. 2016 [13]	Asia (East)	China	Yes	SE ≤ -0.5 D	3469	follow-up from 6 to 15 years of age	N/A	54.9% (45.2–63.5%) ^E
He et al. 2015 [1]	Asia (East)	China	Yes	SE ≤ -0.5 D	951	6 years	6.6 ± 0.34	39.5% (N/A)
Wu et al. 2013 [14]	Asia (East)	China (East)	Yes	SE ≤ -0.5 D	6026	4–18 years	9.7 ± 3.3	36.9% (36.0–38.0%)
Guo et al. 2015 [15]	Asia (East)	China (West)	Yes	SE ≤ -0.5 D	1565	6–21 years	11.9 ± 3.5	60.0% (N/A)
Lin et al. 2014 [16]	Asia (East)	China (rural)	Yes	SE < -0.5 D	585	6–17 years	10.6 ± 2.5	23.3% (N/A)
Saxena et al. 2015 [17]	Asia (South)	India	Yes	SE ≤ -0.5 D	9884	urban school-children	11.6 + 2.2	13.1% (12.5–13.8%)
Aldebasi et al. 2013 [20]	Asia (Middle East)	Saudi Arabia	Yes	SE ≤ -0.5 D	846	primary school children	9.5 ± 1.8	6.5% (N/A)
Al Wadaani et al. 2013 [21]	Asia (Middle East)	Saudi Arabia	Yes	SE ≤ -0.75 D	2002	6–14 years (primary school children)	9.4 ± 2.3	9.0% (7.7–10.2%)
Lundberg et al. [22]	Europe	Denmark	Yes	SE ≤ -0.5 D	307	average age: 9.7, 11.0, 12.9 and 15.4 years (screened at 1–2.5 year intervals)	15.4 ± 0.7	17.9% (N/A)
Tideman et al. 2017 [23]	Europe	Netherlands	Yes	SE ≤ -0.5 D	5711	6 years	6.37 ± 0.7 (myopia) 6.16 ± 0.5 (non-myopia)	2.4% (N/A)
Matamoros et a. 2015 [24]	Europe	France	Yes	SE ≤ -0.5 D ^c	1781	0–9 years	N/A	19.6% (N/A)
Matamoros et a. 2015 [24]	Europe	France	Yes	SE ≤ -0.5 D ^c	8289	10–19 years	N/A	42.7% (N/A)
Kumah et al. 2013 [25]	Africa	Ghana	Yes	SE ≤ -0.5 D ^d	2435	12–15 years	N/A	3.4% (2.7–4.2%)
Lira et al. 2016 [26]	South America	Brazil	Yes	SE < -0.5 D	778	6–17 years	N/A	9.6% (N/A)
Moraes Ibrahim et al. 2013 [27]	South America	Brazil	Yes	SE ≤ -0.5 D	1590	10–15 years	12.2 ± 1.6 (boys) 12.4 ± 1.6 (girls)	3.14% (2.28–4.0%)
Carter et al. 2013 [28]	South America	Paraguay	Yes	SE ≤ -0.5 D	168	5–16 years	N/A	1.4% (N/A)
French et al. 2013 [29]	Australia	Australia	Yes	SE ≤ -0.5 D	863	6 years (younger cohort)	N/A	14.8% (N/A)
French et al. 2013 [29]	Australia	Australia	Yes	SE ≤ -0.5 D	1196	12 years (older cohort)	N/A	17.3% (N/A)
Guo et al. 2017 [30]	Asia (East)	China	No	SE ≤ -0.5 D	35,745	6–18 years (school-based)	12.6 ± 3.4	70.9% (70.5–71.4%)
You et al. 2014 [31]	Asia (East)	China	No ^a	SE ≤ -0.5 D	15,066	7–18 years	13.2 ± 3.4	64.9% (64.2–65.7%)

Table 1 Cross-sectional studies reporting the prevalence of myopia in school children (Continued)


Study	Region	Country	Cycloplegia	Definition	Participants	Age range/cohort (first grade of each school level)	Mean Age	Prevalence (95% CI)
Gong et al. 2014 [32]	Asia (East)	China (Beijing)	No	SE \leq -0.75 D	15,316	7–18 years	12.1 \pm 3.3	53.4% (52.6–54.19%)
Rim et al. 2016 [33]	Asia (East)	South Korea	No	SE < -0.5	7486	7–11 years	N/A	48.2% (45.9–50.6%)
Rim et al. 2016 [33]	Asia (East)	South Korea	No	SE < -0.5	7486	12–18 years	N/A	73.0% (71.0–74.8%)
Mahayana et al. 2017 [34]	Asia (South East)	Indonesia	No	SE \leq -0.5 D	410	N/A (school-based)	10.01 \pm 1.84	32.68% (N/A)
Alrahili et al. 2017 [35]	Asia (Middle East)	Saudi Arabia	No	SE \leq -1.0 D in aged 6 years and more SE \leq -3.0 D in aged 3–6 years	1893	3–10 years	6.2 \pm 1.9	0.7% (N/A)
Hynchak et al. 2013 [36]	North America	United States of America	No	SE < -0.5D ^b	370	10–15 years	N/A	42.2% (N/A)
Galvis et al. 2017 [37]	South America	Colombia	No	SE \leq -0.5 D	1228	8–17 years	11.4 \pm 2.1	11.2% (9.5–13.0%)
Wajuhian et al. 2017 [38]	Africa	South Africa (rural)	No	SE \leq -0.5 D	1586	13–18 years (high-school)	15.81 \pm 1.56	7% (6–9%)
Total					166,934			

Abbreviations CI confidence interval, N/A not assessed, SE spherical equivalent; ^ameasurement under cycloplegia in 1082 of 15,066 children; ^bcycloplegic measurements in about 3% of patients; ^ccycloplegic autorefraction in children, non-cycloplegic refraction in adults, age range not presented; ^dmeasurements under cycloplegia if unaided visual acuity was 20/40 or worse; ^epresented as a population-based cross-sectional survey, however, the results report that the prevalence rate among initial emmetropes and hyperopes, after 5 years



87.7% in 2015, and high myopia (SER greater than -6.0 D) was a major contributor to this increase [43]. In Western China not only myopia prevalence increased, but also a higher rate of annual myopia progression was recently noted [13]. The Waterloo Eye Study showed a long-term increase in myopia

prevalence also in the United States [36]. The prevalence rate reached 42.4% in 10 to 15-year-old children, and 53.9% in 15 to 20-year-old; this was significantly higher than the 21% peak value (in those aged 20–30 years) reported in a comparable study done in 1892 [43].



Risk factors

Risk factors influencing the prevalence of myopia are presented in Table 2. There are some risks factors that can contribute for the prevalence increase such as parental myopia, ethnic differences, less time outdoors, increased near work, population density and socioeconomic status.

Parental myopia

In a study conducted by Lim et al. children (aged 6–18 years) with two myopic parents had a mean refractive error of -2.33 D and the odds ratio of having myopia in childhood with two myopic parents was 2.83, compared with no parental myopia [45]. Although genetic factors

have some impact on eye growth, the development of myopia appears to be mainly influenced by environmental factors such as education [49]. Data from the Handan Offspring Myopia Study the children's refraction was similar to that of their parents at the age of 14 [16]. The inter-generational myopic shift was estimated to be only 1 D at 18 years of age. Thus, it might be concluded that environmental factors such as education influence emmetropization [49]. Looking at between-sibling refractive error in 700 families from the United States, Jones-Jordan et al. found that environmental factors reduced the estimated refractive error correlation between siblings by only 0.5% [50]. This was confirmed by the Collaborative Longitudinal Evaluation of Ethnicity and Refractive Error Study [51]. Less hyperopic and more myopic refractive error at the ages of 7 to 13 years was consistently associated with myopia onset, while having myopic parents, near work and time outdoors were not. The SAVES study revealed that parental myopia was a risk factor for myopia only for the 6-year-old children, but not in 12-year-old cohort [29].

A recent study from Netherlands found seven independently parameters associated with faster axial elongation (AL) in children with 6 to 9 years of age: parental myopia, 1 or more books read per week, time spent reading, no participation in sports, non-European ethnicity, less time spent outdoors, and baseline AL-to-Corneal Radius ratio [52]. Based on the aforementioned results, the authors suggested that behavioural changes are of the highest importance in these children, and employing preventive measures should be considered.

Outdoor time

Outdoor time has been proven to be the strongest environmental factor that can delay myopia onset. The Sydney Adolescent Vascular and Eye Study (SAVES) evaluated the risk factors for the incidence of myopia in Australian schoolchildren during a 5–6 year follow-up period in two cohorts: younger ($n = 892$; aged 6 years at baseline) and older ($n = 1211$; aged 12 years at baseline) [29]. The children that became myopic spent less time outdoors compared to those who remained nonmyopic (16.3 vs 21.0 h in the younger cohort, $p < 0.0001$; and 17.2 vs 19.6 h in the older cohort, $p = 0.001$). The Avon Longitudinal Study of Parents and Children confirmed the negative association between the time outdoors and myopia; additionally the time outdoors in 3 to 9 years-old age range was associated with reduced incidence of myopia at the age of 10 to 15 years [53]. Another study showed that patterns of daily outdoor light exposure differed substantially between Australian (105 ± 42 min/d) and Singaporean children (61 ± 40 min/d; $p = 0.005$) [54].

Myopia progression was not strongly associated either with near work or outdoor/sports activity in siblings with common environmental exposures [50]. In a

Table 2 Risk factors for the prevalence of myopia in the analyzed studies

Risk factor	Country where the study was conducted
Female gender	China [10, 12, 14, 15] Colombia [37] India [17] Saudi Arabia [21, 35]
Low outdoor activity	Australia [27] China [1, 15] Netherlands [23]
Parental myopia	Australia (6-year-old cohort) [29] China [12, 32, 45] Finland [42] India [17] Japan [46]
Increasing age	Brazil [26] China [12, 14, 15] India [17] Poland [18, 19] Saudi Arabia [21, 35]
Time spent on near work/studying	Australia [29] China [12, 32] India (over 5 h daily) [17] Taiwan [47]
Higher socio-economic status	India [17]
Low family income	Netherlands [23]
Higher body mass index	Japan [46] Netherlands [23]
Use of LED lamps for homework (compared to incandescent or fluorescent lamps)	China [11]
Urban environment, high population density and small home size	China [14] Hong Kong [48] Indonesia [34]
Rural environment	Saudi Arabia (only in girls) [21]
Private schooling and watching TV over 2 h daily and playing mobile/video games	India [17]
Low sleeping hours	China [32]
Lower vitamin D levels, less participation in sports and foreign descent	Netherlands [23]
Westernized dietary habits	Japan [46]

randomized clinical trial by He et al., a 40-min class of outdoor activities on each school day for 3 years resulted in a reduced incidence of myopia from 39.5 to 30.4% [1]. A recent RCT showed that outdoor activities can inhibit progression in myopic children aged 6 to 7 years old by 30% in 1 year [2]. These results might indicate that high-risk patients require a sum of treatments for the control of the condition, including changes in lifestyle (increase in outdoor time) and treatment with atropine eye drops, progressive contact lenses or orthokeratology.

Interesting results were presented in a recent study on 16–19-year-old Norwegian Caucasians ($n = 393$) living in 60° latitude North, where autumn-winter is 50 days longer than summer [55]. In their investigation the total time spent doing outdoor was not associated with myopia (3.65 ± 1.5 h in myopes, and 3.81 ± 1.9 in non-myopes, $p = 0.64$). Moreover, the prevalence of myopia was quite low (13% for SER lower than ≤ -0.5 D), despite the few daylight hours in the autumn-winter period (10 h 36 min–11 h 5 min) and high levels of indoor activity and near work [55]. The commonly agreed underlying mechanism of time spent outdoors proposed by researchers is based on the release of retinal dopamine that controls scleral growing and remodeling. Genetic observations add credence to the current notion that myopia is caused by a retina-to-sclera signaling cascade that induces scleral remodeling in response to light stimuli [56]. However, it is possible that other variables may influence emmetropization, including ultraviolet light [57] or blue-light [58]. Moreover, a recent systematic review found that lower blood vitamin D concentrations are associated with increased risk of myopia; on the other hand serum vitamin D levels may be just a proxy for time outdoors [59]. Viewing distances are also much greater outdoors, with the accommodative requirements being smaller and giving a more uniform dioptric space [60]. Animal studies provided evidence that sustained hyperopic defocus, which is generated indoors, promotes local eye growth and myopia [61].

The Childhood Health, Activity, and Motor Performance Eye Study determined the association between physical activity and myopia; in a group of 307 Danish children accelerometer measurements were conducted at mean ages of 9.7, 11.0, 12.9 and 15.4 years [22]. The prevalence of myopia at the final time-point was 17.9% and was not associated with physical activity. In an American cohort, exercise was associated with a lower rate of myopia [62]. Tideman et al. found that myopic children ($n = 5711$, six-year-old children) in Europe spent less time outdoors, have lower vitamin D3 and higher body mass index than non-myopic children [23]. Similarly, Terasaki et al. analysed lifestyle factors related to myopia progression in the third-year elementary school students in Japan [46]. High body weight, parental

myopia and Westernized dietary habits were associated with increased myopia prevalence. In Finland higher myopia during maturity was related to parental myopia, less time spent on sports and outdoor activities during school-years [42]. However, less time spent on sports might also be a proxy of low outdoor activity. Another recent investigation reported a relationship between myopia and BMI, with obese children having higher risk of developing myopia [63]. These results should be interpreted carefully as obese children may engage less in exercise and outdoor activities, as this may be a confound factor that needs further research. Those represent an important potentially modifiable risk factor that may be a target for future public health efforts, involving the protection of children not only from myopia, but also from other unhealthy behaviors that can impact health.

Translation of research findings, regarding outdoor time, to clinical practice is also growing rapidly. A recent questionnaire applied to paediatric ophthalmologists all over the world show that 86 % of the respondents advised children with myopia to spend more time outdoors [64].

Near work

The SAVES study revealed that near work was a risk factor for myopia but only for the 6-year-old children, and not in 12-year-old cohort [29]. This result might indicate that near work can be a factor for inducing the earlier onset of myopia in smaller children. There might be a difference in the mechanism of setting myopia development between early onset and later onset myopes. Children who became myopic performed significantly more near work (19.4 vs 17.6 h), which was statistically significant ($p = 0.02$), however, the association was lower than for outdoor activity. A combination of both factors may be implicated in the myopia development. Shorter reading distance and higher myopia at the baseline exam (1-year prior to the final assessment) were risk factors for myopia progression in a cohort of second-grade primary school (age 7–8 years) children in Taipei [47]. In this study, fast myopia progression in children was associated with more myopia at baseline and shorter reading distance. Similarly, in a study of primary and middle school-aged pupils in Guangzhou ($n = 3055$, mean age of 13.6 ± 1.6 years), children whose reading distance was less than 25 cm were more likely to have myopia than those reading from a distance of 25–29 cm or over 29 cm ($p < 0.001$) [12]. In the same study, reading for more than 2 hours daily was positively associated with myopia in boys, while spending time watching television per week was associated with myopia in girls. Taiwanese children attending private classes outside the regular school system in the evening or on the weekends for ≥ 2 h/d had increased risk of myopia

occurrence [65]. The authors hypothesize that this effect may be due to increase near visual activity or reduced time outdoors. Because the effect of near distance activities on myopia onset and progression was shown to be higher in younger children, it seems to be reasonable to limit unnecessary time spent on near distance activities (including electronic devices) by pre-school children, and these activities should be under strict parental control.

LED lamps and hours of sleep

In a study by Pan et al., conducted on 2346 Chinese children aged 13 to 14-years-old using LED lamps for homework after school had a higher prevalence of myopia (SER less than -0.75 D) and longer axial length than those using incandescent ($p = 0.04$ and $p = 0.007$, respectively) or fluorescent lamps ($p = 0.02$ and $p = 0.003$, respectively) [11]. Gong et al. found low hours of sleep to be an independent risk factor for myopia in 15,316 students of mean age of 12.1 ± 3.3 years from 18 districts of Beijing. Children with 7 hours or less of sleep (odds ratio 3.37, 95% confidence interval (CI): 3.07–3.70, $p < 0.001$) or about 8 h of sleep (odds ratio 2.12, 95% CI 1.94–2.31, $p < 0.001$) had a higher risk compared to those who slept 9 hours or more daily [32]. A disadvantage of the study was that it analysed non-cycloplegic SER. The mechanism underlying the sleeping time-myopia relationship is not well understood yet and future research is needed; presumably inactivity of the ciliary muscle during the sleep could prevent or alleviate the myopic progression. However, other new variables may be involved such as the effect of dim light. New findings suggest that in addition to bright light exposure, rod pathways stimulated by dim light exposure could be important to human myopia development [66]. One study with Australian children aged 10 to 15 years old demonstrated that myopic children spent less time in both scotopic and outdoor photopic light conditions compared with non-myopic children. Myopes may also have reduced sensitivity to low spatial frequency S-cone stimuli with consequences in their failure to emmetropize normally [67].

Population density

Higher population density seems to be associated with myopia risk, independent of time spent outdoors and other environmental factors [68]. High population density and small home size was also associated with longer axial length and refractive error in children in Hong Kong and Beijing [48, 69]. The Beijing study reported other risks factors that are associated with myopia, such as urban region of habitation [69]. In a cohort of 12-year-old children from the urban region of greater Beijing the prevalence of myopia was 70.9% [30]. The mean refractive error in 18-year-olds was -3.74 ± 2.56 D.

The prevalence of myopia was highest in provincial capitals in Hubei province, followed by non-provincial cities, and the lowest in rural areas, with a statistically significant difference ($p < 0.05$) [70]. On the other hand, the incidence of myopia in a rural area in southwestern Japan was very low, from 0.3 to 4.9% over a five-year study in the late 1990s [71]. In general, high population density might be a surrogate of outdoor and near work; children in urban areas might spend less time outdoors, as they might may not have available places where to play.

Socioeconomic status (SES)

In the North India Myopia Study, the prevalence of myopia was 13.1% [17]. Myopia was more common among children with higher SES and among private school students, compared to governmental school pupils. Presumably children in private schools spend more hours at school compared to children in public schools; they spend more time reading and writing at home, with significantly more pressure and a greater likelihood of extra classes. Studying and reading for over 5 hours daily, watching television for over 2 hours daily, and playing video/mobile games were also significantly associated with myopia. In this study there was no obvious mechanism linking higher SES and attending private schools to myopia, except through the education that the children received. A plausible hypothesis would be that children from higher SES families and private schools would be getting more intensive education, as within the study children from private schools spent more time reading at home than those from government schools ($p < 0.001$).

Contradictory findings were reported by a Dutch study of a multi-ethnic cohort of 6-year-old children, revealing a significant influence of socioeconomic factors on the prevalence of myopia [23]. In particular, children of non-European descent, with children from low maternal education, low family income, were more likely to be myopic. These findings are in contrast to the results cited earlier in North Indian children [17]. However, children from families with a non-European ethnic background, similarly to those in private schools in North India, spend lower time outdoors [23]. The children in the Rotterdam study are still very young, and the effects of education are unlikely to be clear, and although their parents have low incomes, they might have a greater commitment to education as a pathway to success.

Discussion

Cycloplegic refraction is established as the gold standard for epidemiological studies on refractive errors. Nevertheless, within our review nine studies used non-cycloplegic measures, while 19 studies presented cycloplegic refraction. Studies reporting non-cycloplegic measurements of

prevalence cannot be considered as reliable; application of non-cycloplegic measurements leads to substantial errors, both in prevalence rates and associations with risk factors [77, 72]. For example, Lundberg et al. reported the prevalence of myopia in children reaching 33.6% using non-cycloplegic measurements and 17.9% under cycloplegia [22]. In the Shandong Children Eye Study the difference between cycloplegic and non-cycloplegic SER was 0.78 ± 0.79 D; this difference decreased with age and increased with greater hyperopic refractive error [73]. In the study by Fotedar the difference between cycloplegic and non-cycloplegic SER was 1.18 D (95% CI: 1.05–1.30 D) for 6 year-old-children, and 0.84 D (0.81–0.87 D) for 12-year-old-children [74]. Thus, the refractive error was misclassified in 9.5% 6-year-old children, and 17.8% 12-year-old children [74]. Interestingly, the Beijing Myopia Progression Study (which enrolled children aged 6 to 17 years) found that a major difference between non-cycloplegic and cycloplegic was associated with the progression of myopia in children, but not with the onset of myopia [75].

The different studies included in this review used different definitions of myopia. Most studies define myopia as a SER less than or equal to -0.5 D. Some studies use a criterion of SER less than -0.5 D, or less than or equal to -0.75 D. Myopia was also defined as a SER less than or equal to -1.0 D in children aged 6 years. A SER greater than or equal to -3.0 D in children aged 3–6 years was reported in one study [35]. The definition of myopia is of extreme importance, and even small changes in the threshold definition (± 0.25 D) have been shown to affect significantly the conclusions of epidemiological studies [76, 77, 78]. Recently, the International Myopia Institute suggested employing a ≤ -0.5 D threshold as an evidence based consensus [79].

Another issue is the choice of eye; usually measurements of the right eye are included in the analysis. In one study conducted in Seoul, South Korea, the age standardized prevalence was reported to be as high as 80% in children aged 12–18 years [33]. However, myopia was classified according to the Korea Centers for Disease Control and Prevention; it employs definition of myopia as a SER greater than or equal to -0.75 D in either (worse) eye. When applying a definition of < -0.5 D in the right eye, the prevalence rates dropped down to 73%. Moreover, one should consider that these results were significantly biased as the measurements were done without cycloplegia.

Interestingly, in some migrant groups, primarily of East Asian origin, the children were significantly more myopic than those of European origin, presumably because of the intensive education that the children are receiving [80, 81]. Children of East Asian ethnicity spend less time outdoors and more time in near work activities compared to European Caucasian children at all school

ages [80]. Rudnicka et al. found that the increase in myopia prevalence over the last decade is related with ethnic differences, with only a small change seen in whites but a significant increase observed in East Asians and a weaker increase among South Asians [82]. Myopia was also common in a diverse Southern Californian pediatric cohort and children of Asian ancestry had the highest prevalence. Particular lifestyle habits in different populations may partially explain dissimilarities in myopia prevalence [62]. It has been suggested that a probable causative role in the development of myopia is the competitive and stressful education systems in some East Asian countries [83].

New risk factors, apart from outdoor time, such as the use of LED lamps for homework, dim light, low sleeping hours, reading distance less than 25 cm and living in an urban environment were described in recent studies. Additional epidemiological studies should be carried out to further expand the knowledge of outdoors on myopia progression. Interventional studies might be also needed to better understand the effectiveness of preventive methods in different settings and age groups. Although light intensity patterns in humans have been implicated in myopia protection, research needs to be further expanded to understand how bright needs to be the exposure to avoid myopia. Longitudinal patterns of light exposure in different refractive errors (e.g. myopes, hyperopes and emmetropes) are needed to understand which of light parameters is the most important (e.g. light intensity, duration or regularity). This study did not focus on the prevalence of high myopia, which is an important indicator and should be further developed.

Conclusion

It can be concluded that prevalence rates were shown to increase in Asia, but also in Europe and North America. Particular lifestyle habits in different populations may partially explain dissimilarities in myopia prevalence between geographical regions. Preventive measures such as outdoor programs and changes on near distance activities in preschool children should be implemented.

Supplementary information

Supplementary information accompanies this paper at <https://doi.org/10.1186/s12886-019-1220-0>.

Additional file 1. The list of cross-sectional studies reporting the prevalence of myopia in school children.

Acknowledgements

Nothing to acknowledge.

Authors' contributions

AG concept and design, data acquisition and analysis, critical revision of the manuscript. PK: data acquisition and analysis, interpretation of data for the work, drafting the manuscript. KT, SSW, CL: interpretation of data for the

work, critical revision of the manuscript. All authors read and approved the final manuscript.

Funding

None.

Availability of data and materials

N/A (review article).

Ethics approval and consent to participate

N/A

Consent for publication

N/A (no identifying data).

Competing interests

Dr. Grzybowski reports grants, personal fees and non-financial support from Bayer, non-financial support from Novartis, non-financial support from Alcon, non-financial support from Thea, personal fees and non-financial support from Valeant, non-financial support from Santen, outside the submitted work. Dr. Kanclerz reports non-financial support from Visim. Dr. Tsubota reports grant support from Jins, Inc., Toshiba Materials Co., Ltd., Asahi Glass, Rohto, Toray and Plus One Marketing; in addition, patents are pending with Jins, Inc., Asahi Glass, Toshiba Materials Co., Ltd. for myopia prevention-related devices. Dr. Carla Lanca and Dr. Seang-Mei Saw have no conflicts of interest to declare.

Author details

¹Department of Ophthalmology, University of Warmia and Mazury, Olsztyn, Poland. ²Foundation for Ophthalmology Development, Institute for Research in Ophthalmology, Gorzyczewskiego 2/3, 60-554 Poznan, Poland. ³Private Practice, Gdańsk, Poland. ⁴Department of Ophthalmology, Keio University School of Medicine, Tokyo, Japan. ⁵Tsubota Laboratory, Inc., Tokyo, Japan. ⁶Singapore Eye Research Institute, Singapore, Singapore. ⁷Saw Swee Hock School of Public Health, National University of Singapore, Singapore, Singapore. ⁸Duke-NUS Medical School, Singapore, Singapore.

Received: 29 March 2019 Accepted: 14 October 2019

Published online: 14 January 2020

References

- He M, Xiang F, Zeng Y, et al. Effect of time spent outdoors at school on the development of myopia among children in China. *JAMA*. 2015;314:1142.
- Wu P-C, Chen C-T, Lin K-K, et al. Myopia prevention and outdoor light intensity in a school-based cluster randomized trial. *Ophthalmology*. 2018; 125:1239–50.
- Wu P-C, Tsai C-L, Wu H-L, et al. Outdoor activity during class recess reduces myopia onset and progression in school children. *Ophthalmology*. 2013;120: 1080–5.
- Jin J-X, Hua W-J, Jiang X, et al. Effect of outdoor activity on myopia onset and progression in school-aged children in Northeast China: the Sujiatun eye care study. *BMC Ophthalmol*. 2015;15:73.
- Holden BA, Fricke TR, Wilson DA, et al. Global prevalence of myopia and high myopia and temporal trends from 2000 through 2050. *Ophthalmology*. 2016;123:1036–42.
- Ikuno Y. Overview of the complications of high myopia. *Retina*. 2017;37: 2347–51.
- Naidoo KS, Fricke TR, Frick KD, et al. Potential lost productivity resulting from the global burden of myopia: systematic review, meta-analysis, and modeling. *Ophthalmology*. 2018;126:338–46.
- Chua SYL, Sabanayagam C, Cheung Y-B, et al. Age of onset of myopia predicts risk of high myopia in later childhood in myopic Singapore children. *Ophthalmic Physiol Opt*. 2016;36:388–94.
- Gomez-Salazar F, Campos-Romero A, Gomez-Campaña H, et al. Refractive errors among children, adolescents and adults attending eye clinics in Mexico. *Int J Ophthalmol*. 2017;10:796–802.
- Li Y, Liu J, Qi P. The increasing prevalence of myopia in junior high school students in the Haidian District of Beijing, China: a 10-year population-based survey. *BMC Ophthalmol*. 2017;17:88.
- Pan C-W, Wu R-K, Liu H, et al. Types of lamp for homework and myopia among Chinese school-aged children. *Ophthalmic Epidemiol*. 2017;25:1–7.
- Guo L, Yang J, Mai J, et al. Prevalence and associated factors of myopia among primary and middle school-aged students: a school-based study in Guangzhou. *Eye*. 2016;30:796–804.
- Zhou W-J, Zhang Y-Y, Li H, et al. Five-year progression of refractive errors and incidence of myopia in school-aged children in Western China. *J Epidemiol*. 2016;26:386–95.
- Wu JF, Bi HS, Wang SM, et al. Refractive error, visual acuity and causes of vision loss in children in Shandong, China. *The Shandong Children Eye Study*. *PLoS One*. 2013;8:e82763.
- Guo K, Yang DY, Wang Y, et al. Prevalence of myopia in schoolchildren in Ejina: the Gobi Desert children eye study. *Invest Ophthalmol Vis Sci*. 2015;56: 1769–74.
- Lin Z, Gao TY, Vasudevan B, et al. Generational difference of refractive error and risk factors in the Handan offspring myopia study. *Invest Ophthalmol Vis Sci*. 2014;55:5711–7.
- Saxena R, Vashist P, Tandon R, et al. Prevalence of myopia and its risk factors in urban school children in Delhi: the North India myopia study (NIM study). *PLoS One*. 2015;10:e0117349.
- Grzybowski A, Nowak MS. The review of refractive studies of nonadults in Poland. *Acta Ophthalmol*. 2018;96:e755–6.
- Kanclerz P, Przewlocka K. Results of refractive error screening of Polish schoolchildren aged 6 to 14 years. *Przegl Lek*. 2019;76:468–71.
- Aldebari YH. Prevalence of correctable visual impairment in primary school children in Qassim Province, Saudi Arabia. *J Optom*. 2014;7:168–76.
- Al Wadaani FA, Amin TT, Ali A, et al. Prevalence and pattern of refractive errors among primary school children in Al Hassa, Saudi Arabia. *Glob J Health Sci*. 2013;5:125–34.
- Lundberg K, Suhr Thykjaer A, Søgaard Hansen R, et al. Physical activity and myopia in Danish children-The CHAMPS Eye Study. *Acta Ophthalmol*. 2017; 96:134–41.
- Tideman JW, Polling JR, Hofman A, et al. Environmental factors explain socioeconomic prevalence differences in myopia in 6-year-old children. *Br J Ophthalmol*. 2017;102:243–7.
- Matamoros E, Ingrand P, Pelen F, et al. Prevalence of myopia in France: a cross-sectional analysis. *Medicine*. 2015;94:e1976.
- Kumah BD, Ebri A, Abdul-Kabir M, et al. Refractive error and visual impairment in private school children in Ghana. *Optom Vis Sci*. 2013;90:1456–61.
- Lira RPC, Arieta CEL, Passos THM, et al. Distribution of ocular component measures and refraction in Brazilian school children. *Ophthalmic Epidemiol*. 2017;24:29–35.
- Moraes Ibrahim F, Moraes Ibrahim M, Pompo de Camargo JR, et al. Visual impairment and myopia in Brazilian children: a population-based study. *Optom Vis Sci*. 2013;90:223–7.
- Carter MJ, Lansingh VC, Schacht G, et al. Visual acuity and refraction by age for children of three different ethnic groups in Paraguay. *Arq Bras Oftalmol*. 2013;76:94–7.
- French AN, Morgan IG, Mitchell P, et al. Risk factors for incident myopia in Australian schoolchildren: the Sydney adolescent vascular and eye study. *Ophthalmology*. 2013;120:2100–8.
- Guo Y, Duan JL, Liu LJ, et al. High myopia in greater Beijing school children in 2016. *PLoS One*. 2017;12:e0187396.
- You QS, Wu LJ, Duan JL, et al. Prevalence of myopia in school children in greater Beijing: the Beijing childhood eye study. *Acta Ophthalmol*. 2014;92:e398–406.
- Gong Y, Zhang X, Tian D, et al. Parental myopia, near work, hours of sleep and myopia in Chinese children. *Health*. 2014;06:64–70.
- Rim TH, Kim S-H, Lim KH, et al. Refractive errors in Koreans: the Korea National Health and nutrition examination survey 2008–2012. *Korean J Ophthalmol*. 2016;30:214–24.
- Mahayana IT, Indrawati SG, Pawiroranu S. The prevalence of uncorrected refractive error in urban, suburban, exurban and rural primary school children in Indonesian population. *Int J Ophthalmol*. 2017;10:1771–6.
- Alrahili NHR, Jadidy ES, Alahmadi BSH, et al. Prevalence of uncorrected refractive errors among children aged 3–10 years in western Saudi Arabia. *Saudi Med J*. 2017;38:804–10.
- Hrynchak PK, Mittelstaedt A, Machan CM, et al. Increase in myopia prevalence in clinic-based populations across a century. *Optom Vis Sci*. 2013;90:1331–41.
- Galvis V, Tello A, Otero J, et al. Refractive errors in children and adolescents in Bucaramanga (Colombia). *Arq Bras Oftalmol*. 2017;80:359–63.
- Wajuhian SO, Hansraj R. Refractive error in a sample of black high school children in South Africa. *Optom Vis Sci*. 2017;94:1145–52.

39. Xiang F, He M, Zeng Y, et al. Increases in the prevalence of reduced visual acuity and myopia in Chinese children in Guangzhou over the past 20 years. *Eye*. 2013;27:1353–8.
40. Ding B-Y, Shih Y-F, Lin LLK, et al. Myopia among schoolchildren in East Asia and Singapore. *Surv Ophthalmol*. 2017;62:677–97.
41. Li L, Zhong H, Li J, et al. Incidence of myopia and biometric characteristics of premyopic eyes among Chinese children and adolescents. *BMC Ophthalmol*. 2018;18:178.
42. Pärssinen O, Kauppinen M, Viljanen A. The progression of myopia from its onset at age 8–12 to adulthood and the influence of heredity and external factors on myopic progression. A 23-year follow-up study. *Acta Ophthalmol*. 2014;92:730–9.
43. Chen M, Wu A, Zhang L, et al. The increasing prevalence of myopia and high myopia among high school students in Fenghua city, eastern China: a 15-year population-based survey. *BMC Ophthalmol*. 2018;18:159.
44. Herrnhaiser J. Die Refraktionsentwicklung des menschlichen Auges. *Zeitschrift fuer Heilkunde*. 1892;13:342–77.
45. Lim LT, Gong Y, Ah-Kee EY, et al. Impact of parental history of myopia on the development of myopia in mainland China school-aged children. *Ophthalmol Eye Dis*. 2014;6:31–5.
46. Terasaki H, Yamashita T, Yoshihara N, et al. Association of lifestyle and body structure to ocular axial length in Japanese elementary school children. *BMC Ophthalmol*. 2017;17:123.
47. Hsu C-C, Huang N, Lin P-Y, et al. Risk factors for myopia progression in second-grade primary school children in Taipei: a population-based cohort study. *Br J Ophthalmol*. 2017;101:1611–7.
48. Choi KY, Yu WY, Lam CHI, et al. Childhood exposure to constricted living space: a possible environmental threat for myopia development. *Ophthalmic Physiol Opt*. 2017;37:568–75.
49. Goldschmidt E, Jacobsen N. Genetic and environmental effects on myopia development and progression. *Eye*. 2014;28:126–33.
50. Jones-Jordan LA, Sinnott LT, Graham ND, et al. The contributions of near work and outdoor activity to the correlation between siblings in the collaborative longitudinal evaluation of ethnicity and refractive error (CLEERE) study. *Invest Ophthalmol Visual Sci*. 2014;55:6333.
51. Zadnik K, Sinnott LT, Cotter SA, et al. Prediction of juvenile-onset myopia. *JAMA Ophthalmol*. 2015;133:683–9.
52. Tideman JWL, Polling JR, Jaddoe VVW, et al. Environmental risk factors can reduce axial length elongation and myopia incidence in 6- to 9-year-old children. *Ophthalmology*. 2018;126:127–36.
53. Shah RL, Huang Y, Guggenheim JA, et al. Time outdoors at specific ages during early childhood and the risk of incident myopia. *Invest Ophthalmol Vis Sci*. 2017;58:1158–66.
54. Read SA, Vincent SJ, Tan C-S, et al. Patterns of daily outdoor light exposure in Australian and Singaporean children. *Transl Vis Sci Technol*. 2018;7:8.
55. Hagen LA, Gjelle JVB, Arnegard S, et al. Prevalence and possible factors of myopia in Norwegian adolescents. *Sci Rep*. 2018;8:13479.
56. Tedja MS, Wojciechowski R, Hysi PG, et al. Genome-wide association meta-analysis highlights light-induced signaling as a driver for refractive error. *Nat Genet*. 2018;50:834–48.
57. Jiang X, Kurihara T, Torii H, et al. Progress and control of myopia by light environments. *Eye Contact Lens*. 2018;44:273–8.
58. Rucker F, Henriksen M, Yanase T, et al. The role of temporal contrast and blue light in emmetropization. *Vis Res*. 2018;151:78–87.
59. Tang SM, Lau T, Rong SS, et al. Vitamin D and its pathway genes in myopia: systematic review and meta-analysis. *Br J Ophthalmol*. 2019;103:8–17.
60. French AN, Ashby RS, Morgan IG, et al. Time outdoors and the prevention of myopia. *Exp Eye Res*. 2013;114:58–68.
61. Flitcroft DL. The complex interactions of retinal, optical and environmental factors in myopia aetiology. *Prog Retin Eye Res*. 2012;31:622–60.
62. Theophanous C, Modjtahedi BS, Batech M, et al. Myopia prevalence and risk factors in children. *Clin Ophthalmol*. 2018;12:1581–7.
63. Lim DH, Han J, Chung T-Y, et al. The high prevalence of myopia in Korean children with influence of parental refractive errors: the 2008–2012 Korean National Health and nutrition examination survey. *PLoS One*. 2018;13:e0207690.
64. Zloto O, Wygnanski-Jaffe T, Farzavandi SK, et al. Current trends among pediatric ophthalmologists to decrease myopia progression—an international perspective. *Graefes Arch Clin Exp Ophthalmol*. 2018;256:2457–66.
65. Ku P-W, Steptoe A, Lai Y-J, et al. The associations between near visual activity and incident myopia in children: a Nationwide 4-year follow-up study. *Ophthalmology*. 2018;126:214–20.
66. Landis EG, Yang V, Brown DM, et al. Dim light exposure and myopia in children. *Invest Ophthalmol Vis Sci*. 2018;59:4804–11.
67. Taylor CP, Shepard TG, Rucker FJ, et al. Sensitivity to S-cone stimuli and the development of myopia. *Invest Ophthalmol Vis Sci*. 2018;59:4622–30.
68. Zhang M, Li L, Chen L, et al. Population density and refractive error among Chinese children. *Invest Ophthalmol Vis Sci*. 2010;51:4969–76.
69. Guo Y, Liu LJ, Xu L, et al. Outdoor activity and myopia among primary students in rural and urban regions of Beijing. *Ophthalmology*. 2013;120:277–83.
70. Zeng CQ, Zhou LH, Zhang P, et al. The epidemiology of myopia in primary school students of grade 1 to 3 in Hubei province. *Zhonghua Yan Ke Za Zhi*. 2018;54:756–61.
71. Yamashita T, Watanabe S, Ohba N. A longitudinal study of cycloplegic refraction in a cohort of 350 Japanese schoolchildren. *Anisometropia. Ophthalmic Physiol Opt*. 1999;19:30–3.
72. Fotouhi A, Morgan IG, Iribarren R, et al. Validity of noncycloplegic refraction in the assessment of refractive errors: the Tehran eye study. *Acta Ophthalmol*. 2012;90:380–6.
73. Hu YY, Wu JF, Lu TL, et al. Effect of cycloplegia on the refractive status of children: the Shandong children eye study. *PLoS One*. 2015;10:e0117482.
74. Fotedar R, Rochtchina E, Morgan I, et al. Necessity of Cycloplegia for assessing refractive error in 12-year-old children: a population-based study. *Am J Ophthalmol*. 2007;144:307–9.
75. Lin Z, Vasudevan B, Ciuffreda KJ, et al. The difference between cycloplegic and non-cycloplegic autorefraction and its association with progression of refractive error in Beijing urban children. *Ophthalmic Physiol Opt*. 2017;37:489–97.
76. Cumberland PM, Bountziouka V, Rahi JS. Impact of varying the definition of myopia on estimates of prevalence and associations with risk factors: time for an approach that serves research, practice and policy. *Br J Ophthalmol*. 2018;102:1407–12.
77. Morgan IG, Iribarren R, Fotouhi A, et al. Cycloplegic refraction is the gold standard for epidemiological studies. *Acta Ophthalmol*. 2015;93:581–5.
78. Grzybowski A, Kanclerz P. The standardized definition of high myopia. *Graefes Arch Clin Exp Ophthalmol*. 2019;257:1805.
79. Flitcroft DL, He M, Jonas JB, et al. IML - defining and classifying myopia: a proposed set of standards for clinical and epidemiologic studies. *Invest Ophthalmol Vis Sci*. 2019;60:M20–30.
80. French AN, Morgan IG, Mitchell P, et al. Patterns of myopigenic activities with age, gender and ethnicity in Sydney schoolchildren. *Ophthalmic Physiol Opt*. 2013;33:318–28.
81. Ip JM, Huynh SC, Robaei D, et al. Ethnic differences in refraction and ocular biometry in a population-based sample of 11–15-year-old Australian children. *Eye*. 2008;22:649–56.
82. Rudnicka AR, Kapetanakis VV, Wathern AK, et al. Global variations and time trends in the prevalence of childhood myopia, a systematic review and quantitative meta-analysis: implications for aetiology and early prevention. *Br J Ophthalmol*. 2016;100:882–90.
83. Jiang Y, Tian B. Understanding Modifiable Risk Factors for the Development of Myopia. *Ophthalmology*. 2018;126:221–2.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

