

REVIEW

A review of environmental risk factors for myopia during early life, childhood and adolescence

Clin Exp Optom 2015; 98: 497–506

DOI:10.1111/cxo.12346

Dharani Ramamurthy* PhD BS**Sharon Yu Lin Chua†** BS**Seang-Mei Saw*†** PhD MPH MBBS FAMS FARVO†

*Saw Swee Hock School of Public Health, National University of Singapore, Singapore

†Myopia Unit, Singapore Eye Research Institute, Singapore

E-mail: seang_mei_saw@nuhs.edu.sg

Submitted: 19 December 2015

Revised: 16 August 2015

Accepted for publication 20 August 2015

Myopia is a significant public health problem worldwide, particularly in East Asian countries. The increasing prevalence of myopia poses a huge socio-economic burden and progressive high myopia can lead to sight-threatening ocular complications. Hence, the prevention of early-onset myopia progressing to pathological high myopia is important. Recent epidemiological studies suggest that increased outdoor time is an important modifiable environmental factor that protects young children from myopia. This protective effect may be due to high light intensity outdoors, the chromaticity of daylight or increased vitamin D levels. This review summarises the possible underlying biological mechanisms for the protective association between time outdoors and myopia, including the potential role of nicotinic acetylcholine receptors in refractive error development. Recent evidence for the role of other environmental risk factors such as near work, birth seasons, parental smoking and birth order are also summarised.

Key words: children's vision, myopia, risk factors

Myopia is a significant public health problem across the globe, especially in East Asian countries like Singapore and China. The overall prevalence of myopia in adults aged above 40 years is 38.9 per cent in Singapore,¹ whereas it is much lower in Western countries like the United States² (25.1 per cent), Barbados³ (21.9 per cent) and Australia⁴ (15 per cent). Likewise, the prevalence of myopia is much higher in East Asian children. In 12-year-old children, the prevalence of myopia is 62.0 per cent in Singapore^{5,6} and 49.7 per cent in Guangzhou, China⁷ compared with 20.0 per cent in the United States,⁸ 11.9 per cent in Australia,⁹ 9.7 per cent in urban India^{10,11} and 16.5 per cent in Nepal.¹² As a result of this high prevalence, the mean annual cost of myopia in Singapore teenagers is estimated to be US\$148 per child due to eye examinations and optical purchases.¹³ The mean annual cost of myopia is about US \$709 per adult in Singapore due to eye examinations, optical purchases and laser in situ keratomileusis (LASIK) surgery.¹⁴ Among US adults, the economic cost of myopia is estimated to be \$4.6 billion per year.¹⁵

High myopia may lead to potentially blinding complications, such as retinal tears, myopic macular degeneration and choroidal neovascularisation in both adults¹⁶ and

adolescents¹⁷ that may require surgery, medications, lifelong medical care as well as imposed high social costs.¹⁸

Both genetic and environmental factors play a role in the aetiology of myopia. Near work is an important environmental factor associated with myopia but recent evidence suggests that time spent outdoors is another modifiable risk factor.^{19–22}

This paper provides a review of the major environmental factors that are associated with myopia during early life, childhood and adolescence, in contrast to a number of recent reviews that have mainly focused on the role of genetics in myopia.^{23–25} This review not only considers time spent outdoors and near work but also other early life factors, such as parental smoking, birth season and post-natal light levels, birth order and maternal age that are associated with myopia.

ASSOCIATION BETWEEN TIME SPENT OUTDOORS AND MYOPIA

Evidence from epidemiological studies

Previous cross-sectional and cohort studies have shown a significant association between myopia and outdoor activity among

Australian,^{26,27} Singapore Chinese,²⁸ Taiwanese²⁹ and Caucasian children.^{30–32}

In the Collaborative Longitudinal Evaluation of Ethnicity and Refractive Error study of six- to 14-year-old children of diverse ethnicities, time outdoors/sports activity was one to two hours lower in children who developed myopia compared to stable emmetropes ($p < 0.01$).³¹ Similarly, in a recent population-based cohort study of 3,241 British children aged seven to 15 years,²⁰ children who spent more time outdoors had a lower risk of developing myopia compared to those who spent less time outdoors (hazards ratio [HR] = 0.75, 95 per cent CI: 0.60, 0.96, $p = 0.023$). Although physical activity also had an independent association with incident myopia, time outdoors was a stronger predictor for incident myopia than physical activity²⁰ (Table 1).

There are some potential limitations of this study.²⁰ Refractive error was measured using non-cycloplegic refraction, which tends to overestimate myopia in children.²⁰ This may result in the inadvertent misclassification of children into refractive error groups (incident myopia stable emmetropia), introducing some bias. There was also a significant loss to follow-up in older age groups (up to 85 per cent) and missing data (up

Author	Study design (location)	Sample size	Age	Definition of outdoor activity	Main findings
French et al. ²¹	Population-based cohort study (Australia)	2103	6-12 years	Hours per week on outdoors	Increased OR for incident myopia in children with low and moderate levels of outdoor activity compared to those with high levels of outdoor activity.
Guggenheim et al. ²⁰	Population-based cohort study (ALSPAC [†]) (United Kingdom)	3241	7-15 years	Hours per day on outdoors/physical activity	Time spent outdoors and physical activity was significantly associated with incident myopia
Jones-Jordan et al. ³¹	Longitudinal cohort (CLEERE [‡]) (United States)	731	6-14 years	Hours per week on sports/outdoor activities	Reduced time spent outdoors in children who became myopic compared to emmetropes
Jones-Jordan et al. ³²	Population-based cohort study (CLEERE) (United States)	835	6-14 years	Hours per week on outdoor/ sports activity	No significant association between time spent outdoors and myopia progression

[†]Avon Longitudinal Study of Parents and Children
[‡]Collaborative Longitudinal Evaluation of Ethnicity and Refractive Error study

Table 1. Association between outdoor activity and myopia – evidence from epidemiological studies

to 13 per cent). Consequently, the association between myopia and time spent outdoors may have differed between children who remained in the study compared to those who discontinued.

In another population-based cohort study²¹ of six- and 12-year-old Australians, including Caucasian and East Asian children, those with low and moderate levels of outdoor activity were more likely to develop myopia compared to children who performed more outdoor activity in both the younger (adjusted odds ratios [ORs] = 2.84, 95 per cent CI: 1.56, 5.17 and 1.14, 95 per cent CI: 0.59, 2.21, respectively for low and moderate groups, $p_{\text{trend}} < 0.0001$) and older cohort (adjusted ORs = 2.15, 95 per cent CI: 1.35, 3.42 and 2.00, 95 per cent CI: 1.28, 3.14, respectively for low and moderate groups, $p_{\text{trend}} < 0.001$) (Table 1).

Time spent outdoors/sports activities was not significantly associated with myopia progression ($\beta = 0.03$, 99 per cent CI: -0.03, 0.08, $p > 0.01$) in six- to 14-year-old children of various ethnicities.³² While increased outdoor time seems to have a protective effect on incident myopia, this effect may not be pertinent for myopic progression following its initial onset. A recent intervention study has also shown a significant reduction in the rate of incident myopia but not for myopic progression.⁴⁸

A significant limitation of these studies is that outdoor activities were assessed subjectively using questionnaires, rather than an objective measure of light exposure.^{20,21,26-32} Precise quantification of time spent outdoors is a major challenge in epidemiological

studies. Questionnaires have been widely used but they only provide a subjective measure of time outdoors, are susceptible to recall bias and need to be validated against objective measures. Few studies have used objective measures, such as light meters,^{33,34} conjunctival ultraviolet autofluorescence^{35,36} (UVAF) and UV dosimeter.³⁷ A diary recording child activities over a one-week period in conjunction with objective measures of light intensity using a HOBO[®] pendant light meter may provide more accurate data as the corresponding light intensities could provide a guide to the precision of diary data.³³ Conjunctival UVAF has been used as a bio-marker for outdoor light exposure in Australian young adults^{35,36} and is associated with lower myopic prevalence.³⁷

Evidence from other studies

In a recent cross-sectional study of 681 Chinese children in Beijing, time spent outdoors was significantly associated with myopia (adjusted OR = 0.32, 95 per cent CI: 0.21, 0.48, $p < 0.001$).³⁸ Lin and colleagues³⁹ also observed that Chinese children ($n = 370$) with low levels of outdoor activity were significantly more myopic (-1.34 ± 2.45 D) than those with moderate (-0.29 ± 2.11 D) and higher levels of outdoor activity (-0.25 ± -2.06 D; $p_{\text{trend}} = 0.003$). In a school-based longitudinal study of five- to 13-year-old Chinese children,⁴⁰ outdoor time was negatively correlated with axial elongation ($\beta = -0.12$, 95 per cent CI: -0.15, -0.02, $p = 0.01$) and positively correlated with myopic shift ($\beta = 0.13$, $p = 0.005$) (Table 1); however, the Xichang Pediatric Refractive

Error study, which examined 13- to 17-year-old school children in China⁴¹ did not report any association between myopia and outdoor activities (adjusted OR = 1.14, 95 per cent CI: 0.69, 1.89, $p = 0.61$).

There are some potential limitations in this study. Time spent outdoors was self-reported by participants and this may be affected by recall bias. Additionally, 19 per cent of the participants did not complete the survey and the time spent outdoors may have differed between respondents and non-respondents. Thus, these limitations may have resulted in an underestimation of the true association between myopia and time spent outdoors.

There is also substantial evidence that the rate of myopic progression varies across different seasons (Table 2). Fujiwara and colleagues⁴² observed that axial length elongation was slower in summer (0.13 ± 0.008 mm) than in winter for Japanese children (0.15 ± 0.01 mm, $p = 0.04$). Similar results were reported for six- to 12-year-old Chinese children⁴³ with a slower myopic shift (-0.31 ± 0.25 D versus -0.53 ± 0.29 D; $p < 0.001$) and reduced axial elongation in summer than in winter (0.17 ± 0.10 mm versus 0.24 ± 0.09 mm; $p < 0.001$). Danish children with $2,782 \pm 19$ hours of cumulative available daylight showed less myopia progression and axial elongation (0.26 ± 0.27 D and 0.12 ± 0.09 mm), compared to children with 1681 ± 24 hours of daylight (0.32 ± 0.27 D and 0.19 ± 0.10 mm; $p < 0.01$).⁴⁴ This seasonal pattern was also observed in six- to 12-year-old ethnically diverse children from the COMET study,⁴⁵ with a slower myopic shift

Author	Study design (location)	Sample size	Age range	Definition of season/day light hours	Main findings
Fujiwara et al. ⁴²	Longitudinal (Japan)	92	6 to 12 years	Summer: June-Sep Winter: Dec-Mar	Significantly reduced axial elongation in summer compared to winter:
Donovan et al. ⁴³	Longitudinal (China)	85	6 to 12	Summer: June-Aug Autumn: Sep-Nov Winter: Dec-Feb Spring: Mar-May	Significantly lower myopia progression and axial elongation in summer compared to other seasons
Cui et al. ⁴⁴	Cross-sectional (Denmark)	235	8 to 14 years	Cumulative available daylight hours derived from astronomical tables	Significantly lower myopia progression and axial elongation in children with more available daylight hours compared to those with fewer daylight hours
Gwiazda et al. ⁴⁵	Longitudinal (United States)	469	6 to 12 years	Winter: Oct-Mar Summer: Apr-Sep	Significantly lower myopia progression and axial elongation in summer compared to winter

Table 2. Evidence for seasonal variations in myopic progression

in summer than in winter (-0.14 ± 0.32 D versus -0.35 ± 0.34 D; $p < 0.0001$). The slower rate of myopic progression in summer could either be due to increased outdoor activity and reduced near work during the school break in summer⁴⁶ or may be due to more exposure to daylight or blue light.⁴⁷ Both myopic and non-myopic Caucasian children have been shown to spend more time outdoors (21.76 ± 13.80 versus 10.34 ± 6.10 hours per week; $p < 0.001$) and less time studying (1.69 ± 3.71 versus 9.51 ± 6.96 hours per week; $p < 0.001$) during summer break compared to the school year.⁴⁶ The light exposure levels among young adults in the United Kingdom were higher in summer than in winter (586.8 ± 89.4 lux versus 209.9 ± 43.0 lux, $p = 0.0002$) with a greater proportion of blue light exposure during summer than in winter⁴⁷ (41.3 versus 37.4 per cent; $p < 0.0001$).

Evidence from intervention studies

Few randomised controlled trials have been conducted to assess the effect of structured outdoor programs, as an intervention to increase the time spent outdoors by children and retard myopic incidence and progression^{19,48} (Table 3). In the Guangzhou Outdoor Activity Longitudinal study (GOAL),¹⁹ a randomised controlled trial of 1,789 Chinese children aged six to seven years, the two-year incident rate of myopia was significantly lower among children enrolled in an after-school structured outdoor activity (intervention) group compared to a control group (30.4 per cent versus 39.5 per cent, $p < 0.001$). Myopic progression and axial elongation were significantly higher in the control group compared to the intervention group (0.86 ± 0.77 D versus

0.75 ± 0.69 D, $p < 0.01$ and 0.61 ± 0.33 mm versus 0.59 ± 0.33 mm, $p < 0.01$, respectively). Another school-based intervention study involving 571 Chinese Taiwanese children aged seven to 11 years⁴⁸ reported a significantly lower incident rate of myopia in children who were in the Recess Outdoor the Classroom (ROC) Program compared to a control group (8.4 versus 17.65 per cent, $p = 0.001$). Myopic progression was significantly lower in the intervention group compared to the control group among non-myopic subjects (-0.25 ± 0.68 D versus -0.39 ± 0.69 D, $p = 0.02$). A proportion of myopic children was also under atropine treatment in both the intervention (~ 29 per cent) and control (~ 20 per cent) groups and the outdoor intervention program did not show a combined effect with atropine on myopic progression ($p = 0.85$). The Family Incentive Trial (FIT), a randomised controlled trial of 285 Singapore children estimated time spent outdoors using a questionnaire and a one-week child activities diary, and demonstrated that structured weekend outdoor intervention programs significantly increased outdoor time; 14.75 ± 7.52 hours per week for the intervention group compared to 12.40 ± 6.94 hours per week for the control group, ($p = 0.004$) at six months.⁴⁹

MECHANISMS UNDERLYING THE PROTECTIVE EFFECT

High light levels – evidence from animal studies

Animal studies have shown that light levels as high as 15,000 to 30,000 lux either in the form of artificial laboratory light or natural

daylight retard experimental myopia in chicks,^{50–53} guinea pigs⁵⁴ and monkeys.^{55,56} Chicks reared under either light–dark cycles or continuous illumination (50, 500 or 10,000 lux) display increased dopamine concentration for higher luminance levels in both groups.⁵¹ Under both experimental conditions, high light intensity and dopamine concentration were significantly associated with less myopic development (light–dark cycle group: $r = 0.91$, $p < 0.0001$ and continuous light group: $r = 0.74$, $p < 0.0001$).⁵¹ Backhouse, Collins and Phillips⁵² showed that chick eyes exposed to continuous lighting of 2,000 lux developed significantly less form deprivation myopia (-4.94 ± 1.21 D) compared to those exposed to continuous lighting of 300 lux (-9.73 ± 0.96 D, $p = 0.022$) or brief periods of 10,000 lux (-9.98 ± 0.85 D; $p = 0.017$). In addition, the intravitreal injection of the dopamine agonist spiperone retards the development of form deprivation myopia.⁵³ A recent study has shown that drugs that activate D1-like receptors (SKF 38390) inhibit the progression of naturally occurring progressive myopia in albino guinea pigs, whereas activation of D2-like receptors by quinpirole promoted progressive myopia.⁵⁴ In monkeys, light intensity as high as 25,000 lux is protective against form deprivation myopia⁵⁵ but not against lens-induced myopia.⁵⁶

Based on the results from animal models and epidemiological studies, it is hypothesised that high light levels outdoors or rapid luminance changes⁶⁰ trigger the release of dopamine, which is an ocular growth inhibitor,^{22,26,27,57–59} which inhibits myopic development.⁶⁰

Author	Study design (location)	Sample size	Age	Type of intervention	Main findings
Morgan et al. ¹⁹	School-based cluster randomised trial (GOAL study [†]) (China)	1789	6-7 years	Structured outdoor activity outside school hours	Lower incident myopia rate in the intervention group compared to the control group
Wu et al. ⁴⁸	Prospective school-based intervention study (Taiwan)	571	7-11 years	Recess outside the classroom (ROC) program to increase outdoor time	Lower incident myopia rate in the intervention group compared to the control group
Ngo et al. ⁴⁹	Randomised controlled trial (FIT [‡]) (Singapore)	285	6-12 years	Structured weekend outdoor activities, incentives and myopia education	Increased outdoor time in the intervention group compared to the control group

[†] Guangzhou Outdoor Activity Longitudinal study;
[‡] Family Incentive Trial

Table 3. Association between outdoor activity and myopia – evidence from intervention studies

Light chromaticity and spectral composition – evidence from animal studies

There is some evidence that the chromaticity and spectral composition of ambient light may influence myopic inhibition. The longitudinal chromatic aberration of the eye means that not all wavelengths are equally focused on the retina, thus causing reduced contrast of the wavelengths that are focused away from the retina. Hyperopic defocus reduces the contrast of long wavelength components to a greater extent, resulting in a chromatic blur of the retinal image, which could be a guide for the axial elongation of the eye.⁶¹

The spectral composition of light has a significant impact on ocular growth in guinea pigs.^{62–64} Guinea pigs reared in long wavelength light display a significantly more myopic refraction ($+1.78 \pm 1.22$ D) relative to those reared in mixed wavelength light ($+3.60 \pm 1.65$ D) and white light⁶² ($+5.20 \pm 1.67$ D, $p < 0.05$). Guinea pigs reared under short wavelength light developed significantly more hyperopia ($+6.08 \pm 0.80$ D) compared to those reared under medium wavelength light ($+2.96 \pm 0.68$ D, $p < 0.01$) and broadband light⁶³ ($+1.36 \pm 0.65$ D, $p < 0.001$). Vitreous chamber depth was also shorter in guinea pigs raised under blue light (3.23 ± 0.09 mm) compared to those reared under medium wavelength light (3.36 ± 0.10 mm, $p < 0.01$) and broadband light⁶³ (3.51 ± 0.11 mm, $p = 0.0001$). Guinea pigs reared in red light developed nearly 2.50 D more myopia ($p < 0.01$) and substantially longer eyes (0.20 mm longer, $p = 0.02$) compared to those reared under blue light or white light.⁶⁴ Rearing guinea pigs under blue light suppresses lens-induced myopia, whereas white light leads to the development of significant myopia.⁶⁴

A similar phenomenon has been observed in chicks with red-green flicker producing increased axial elongation compared to luminance flicker⁶¹ (50 per cent more ocular growth, $p < 0.05$). Altering the chromaticity of ambient light appears to induce and reverse myopia and hyperopia in chicks.⁶⁵ In chicks, an excess of red light caused myopia (-2.83 ± 0.25 D), while an excess of blue light caused hyperopia⁶⁵ ($+4.55 \pm 0.21$ D). Additionally, red light-induced myopia (-2.21 ± 0.21 D) in chicks could be reversed to hyperopia ($+2.50 \pm 0.29$ D) by changing the excess red light to blue light. Blue light-induced hyperopia ($+4.21 \pm 0.19$ D) could also be reversed to myopia (-1.23 ± 0.12 D) by changing blue light to red light.⁶⁵ Together these results suggest that exposure to shorter wavelength blue light is protective against myopia. Since daylight is predominantly composed of blue light, the observed association between time spent outdoors and myopic inhibition in humans may be related to the spectral composition of light.

Increased blood vitamin D levels – evidence from epidemiological and other studies

Exposure to solar ultra violet B radiation (UV-B) outdoors may trigger vitamin D synthesis.⁶⁶ In a cross-sectional study of 13- to 25-year-old subjects⁶⁷ blood vitamin D levels were inversely related to myopia ($\beta = -3.4$, $p = 0.005$) with myopes having a lower blood level of vitamin D by 3.4 ng/ml compared to non-myopes. Similar results were found among 2,038 adolescents aged 13 to 18 years⁶⁸ with a positive association between serum vitamin D levels and refractive error ($\beta = 0.03$, 95 per cent CI: 0.00, 0.06, $p < 0.05$). Those with higher levels of serum vitamin D were less likely to have high myopia of 6.00 D or more

(adjusted OR=0.55; 95 per cent CI: 0.34, 0.90; $p = 0.02$) compared to those with lower serum vitamin D levels. Another cross-sectional study of 946 young Australian adults showed significantly lower serum levels of vitamin D in myopes compared to non-myopes (67.6 nmol versus 72.5 nmol, $p = 0.003$). Subjects with vitamin D deficiency were more likely to be myopic (adjusted OR=2.07, 95 per cent CI: 1.29 – 3.32, $p = 0.002$) compared to those with sufficient levels of vitamin D.⁶⁹

Similarly, in seven- to 15-year-old British children,⁷⁰ total vitamin D level was significantly higher in children who spent more time outdoors; however, blood vitamin D levels were not associated with incident myopia after controlling for time outdoors,⁷⁰ which suggests that blood vitamin D levels may be a bio-marker for time outdoors. Children who spend more time outdoors may have increased exposure to UV-B radiation and higher blood vitamin D levels.

Vitamin D receptor (VDR) polymorphisms are associated with low to moderate myopia in Whites⁷¹; however, these polymorphisms were insignificant, when myopia of greater than -4.00 D and other ethnicities were taken into account. Annamaneni and colleagues⁷² also found a decreased allele frequency of the *FokI* VDR gene in high myopes compared to controls, but the *FokI* VDR polymorphism was not significantly associated with myopia.

Results from these studies should be interpreted with caution, as these are case-controlled studies with smaller samples than epidemiological studies. Replication studies including larger samples are needed to further confirm the association between these polymorphisms and myopia. Although vitamin D receptor polymorphisms are related to myopia, the functional role of these polymorphisms remains unclear. These

associated polymorphisms may be non-functional, resulting in a failure to obtain any consistent associations with refractive error.

It is speculated that increased vitamin D levels secondary to daylight exposure may inhibit myopia by regulating scleral growth through its anti-proliferative effect or it may be important for the functioning of the smooth ciliary muscle involved in accommodation to achieve a clear retinal image both at distance and near.^{67,71,72} Increased levels of vitamin D and retinoic acid, an ocular growth regulator, may also be involved in signalling and regulation of the cell cycle;^{67,68,71} however, it is difficult to separate the direct effect of vitamin D and vitamin D as a surrogate for outdoor sunlight exposure in human and experimental studies.

Other possible mechanisms underlying the protective effect

The dioptric pattern of the outdoor visual environment may be protective against myopia.⁷³ In an outdoor visual environment, objects are typically further away with less dioptric variations across the visual scene. Thus, an outdoor visual environment is composed of a more uniform dioptric pattern and subsequently the retinal image has a more uniform pattern of retinal defocus in the periphery compared to indoor viewing conditions, which may influence ocular growth and prevent myopia.⁷³

In contrast, objects are much nearer in an indoor visual environment with a higher dioptric

value at the point of fixation, which decreases towards the periphery. Thus, an indoor visual environment consists of greater dioptric variations across the visual scene and the retinal image has higher levels of retinal defocus in the periphery that may accelerate ocular growth.⁷³

Other mechanisms for the protective effect of time outdoors include an increased depth of focus and retinal image clarity (a reduction in higher-order aberrations) due to pupillary constriction under high light intensity outdoors and a reduced accommodative demand for distance viewing, while in outdoor environments.^{26,27,42,46}

ASSOCIATION BETWEEN NEAR WORK AND MYOPIA

Evidence from epidemiological and other studies

Epidemiological studies such as the Orinda Longitudinal Study of Myopia⁷⁴ (OLSM), the Singapore Cohort of Risk factors for Myopia^{5,6} (SCORM) and the Sydney Myopia Study⁷⁵ (SMS) examined the relationship between myopia and near work and have shown equivocal results. Near work appears to be associated with myopia among Caucasian⁷⁴ and Australian children⁷⁵ but was not significantly associated with incident myopia in Singaporean children.^{5,6}

Despite a number of more recent studies investigating the association between myopia

and near work, a clear understanding of the nature of this relationship remains elusive, with some studies showing positive findings^{21,38,31,40,76–78} and others reporting no relationship^{32,39} (Table 4).

In the Beijing Eye Study,⁷⁶ a population-based cross-sectional study of 15,066 children aged seven to 18 years, the odds of having myopia was significantly higher in children, who performed prolonged near work (adjusted OR: 1.15; 95 per cent CI: 1.11 to 1.20; $p < 0.001$) and took less rest during studying (adjusted OR: 1.17; 95 per cent CI: 1.13 to 1.21; $p < 0.001$). A school-based study of Chinese school children aged five to 13 years showed that increased reading time was associated with myopia³⁸ (adjusted OR: 1.38; 95 per cent CI: 1.09 to 1.75; $p = 0.009$) and more axial elongation⁴⁰ ($\beta = 0.13$, 95 per cent CI: 0.02 to 0.12, $p = 0.005$). A cross-sectional study of 5,048 Taiwan military conscripts aged 18 to 24 years, showed that increased near work was associated with a more myopic refractive error⁷⁷ ($\beta = -0.18$; 95 per cent CI: -0.22 to -0.15; $p < 0.001$) and longer axial length ($\beta = 0.10$, 95 per cent CI: 0.07 to 0.13, $p < 0.01$). In a cross-sectional study of 15,316 Chinese children aged six to 18 years,⁷⁸ myopia was significantly associated with increasing levels of near work (adjusted ORs: = 1.14, 95 per cent CI: 1.04 to 1.26; 1.39, 95 per cent CI: 1.24 to 1.56; = 1.43, 95 per cent CI: 1.25 to 1.64, for low, moderate and high near work activity, respectively; $p_{\text{trend}} < 0.001$) and closer reading

Author	Study design (location)	Sample size	Age range	Definition of near work activity	Main findings
French et al. ²¹	Population-based cohort study (Australia)	2103	6 and 12 years	Diopter hours of near work per week	Higher ORs for incident myopia in children with moderate and high levels of near work compared to those with lower levels of near work
Jones-Jordan et al. ²⁹	Longitudinal cohort (‡CLEERE) (United States)	731	6-14 years	Hours per week on reading, studying and computer work	Significantly higher near work in myopic children compared to emmetropes at myopia onset and 3 years after myopia onset.
Jones-Jordan et al. ³¹	Population-based cohort study (United States)	835	6-14 years	Dioptre hours of near work per day	No significant association between dioptré hours of near work and reading
You et al. ⁷⁶	Population-based cross-sectional (China)	15,066	7-18 years	Hours of studying per day	Significant association between myopia and longer hours of reading with less frequent breaks
Lee et al. ⁷⁷	Cross-sectional (Taiwan)	5048	18-24 years	Hours of near work activities per day	Significant association between myopia and time spent on reading
Gong et al. ⁷⁸	Cross-sectional (China)	15,316	6-18 years	Hours of near work activities per day and reading distance	Increasing OR for myopia with increasing near work time and closer reading distances

‡ Collaborative Longitudinal Evaluation of Ethnicity and Refractive Error study

Table 4. Association between near work and myopia – evidence from epidemiological and other studies

distances (adjusted ORs: = 1.39, 95 per cent CI: 1.21 to 1.60; 1.95, 95 per cent CI: 1.24 to 1.69, for 33 cms and more than 33 cms, respectively; $p_{\text{trend}} < 0.001$). In a population-based cohort study of incident myopia among 2,103 Australian children aged six and 12 years, six-year-olds with moderate and higher dioptric hours of near work were more likely to become myopic (adjusted ORs: 1.68; 95 per cent CI: 0.89 to 3.16 and 2.35, 95 per cent CI: 1.30 to 4.27, respectively, $p_{\text{trend}} < 0.001$) compared to those with lower dioptric hours.²¹ Among six- to 14-year-old children from diverse ethnicities, hours spent per week for reading/studying was 0.7 to 1.5 hours more and 0.8 to 1.9 hours more for computer work/video games in children who became myopic compared to emmetropes at the time of onset of myopia and three years after onset of myopia ($p < 0.01$) but not before onset of myopia.³¹

Conversely, several studies have not reported an association between near work and myopia. Near work was not significantly associated with myopia in Chinese children³⁹ or with myopic progression in children from a diverse ethnic group.³² The mean myopic refractions were not significantly different between Chinese children with varying levels of near work activity ($p_{\text{trend}} = 0.94$ and 0.63 , respectively, for primary and secondary school children).³⁹ Myopia progression did not increase with additional 10 hours of near work in children from varying ethnicities ($\beta = -0.007$, 99 per cent CI: -0.02 to 0.004 , $p > 0.01$).³² Recent evidence suggests that the intensity of near work, that is, sustained reading at closer distance (less than 30 cms) with fewer breaks, may be more important than the total hours of near work^{75,76,78}; however, precise quantification of near work is difficult and all studies have used a questionnaire-based approach, which is subjective and may not be reliable.

Possible mechanism - evidence from animal studies

Animal models suggest that retinal image defocus may play a major role in ocular growth and refractive error development in chicks^{79–82} and primates.^{83–85} Negative lens-induced hyperopic retinal defocus results in axial elongation and myopia, whereas positive lens-induced myopic retinal defocus produces hyperopia.⁸⁶ Alterations in the refractive state are accompanied by transient changes in choroidal thickness followed by longer-term changes in scleral ocular growth

in both chicks^{87,88} and primates.^{89,90} Hyperopic defocus causes a rapid thinning of the choroid, thus shifting the retina posteriorly, whereas myopic defocus leads to choroidal thickening, shifting the retina anteriorly to achieve a clear retinal image. In humans, a deficit in the accommodative response (a lag of accommodation), which places the image behind the retina during near work is analogous to the negative lens-induced defocus in animal models. This hyperopic retinal defocus might trigger the growth of the posterior segment to move the retina toward the point of clear focus, leading to axial elongation and myopia. Thus, an individual with a greater lag of accommodation and excessive near work may develop myopia due to hyperopic retinal defocus, which could provide a stimulus for axial elongation.^{91–93} Changes in the size of the retinal defocus area or the blur circle size may regulate the release of retinal neuro-modulators that control ocular growth.⁹⁴

ASSOCIATION BETWEEN SEASON OF BIRTH, POST-NATAL LIGHT LEVELS AND MYOPIA

Evidence from epidemiological and other studies

Birth season and post-natal light levels have also been linked to myopia.^{95–99} A cross-sectional study including 276,911 Israeli defense subjects aged 16 to 22 years, showed that those born during months with longer daylight hours (12.27 to 13.57 hours and 13.58 to 14.23 hours) were more likely to have moderate myopia (adjusted ORs = 1.06, 95 per cent CI: 1.02, 1.10, $p = 0.002$ and 1.08, 95 per cent CI: 1.04, 1.13; $p = 0.002$, respectively) and high myopia (adjusted ORs = 1.11, 95 per cent CI: 1.03, 1.19; $p = 0.004$ and 1.24, 95 per cent CI: 1.16, 1.33, $p < 0.001$, respectively) compared to those born during months with shorter daylight hours.⁹⁵ A cross-sectional study of older British adults reported that subjects born during summer and autumn were more likely to be highly myopic compared to those born in winter⁹⁶ (adjusted ORs = 1.17; 95 per cent CI: 1.05, 1.30, $p = 0.006$ and 1.16; 95 per cent CI: 1.04, 1.30, $p = 0.002$, respectively). Similarly, a cross-sectional study of 722 Caucasian infants aged one to three months⁹⁷ revealed that 28.5 per cent of infants born during longest daylight hours were myopic, whereas only 17.5 per cent of infants born during shortest daylight hours were myopic ($p = 0.02$). Results

from two studies in Finland and China were contradictory^{98,99} and did not show any significant trend of increasing myopic prevalence across birth seasons, quartiles of global irradiance or daily hours of darkness during the birth season.⁹⁸ These results should be considered with caution, as there may be inaccuracies in the assessment or measurement of refractive error. Vannas and colleagues⁹⁸ ascertained myopic status through a questionnaire, whereas other studies^{96–99} used non-cycloplegic refraction to measure refractive error, which tends to overestimate myopia due to uncontrolled accommodation in children and young adults. Thus, there may be a random misclassification of myopia that may have caused a spurious association between birth season and myopia.

Possible mechanism - evidence from animal and human studies

Longer photoperiods may influence ocular growth through abnormal diurnal growth rhythms.^{100–102} Eyes of young chicks reared under a 12 hour/12 hour light/dark cycle with normal visual experience display an increase in axial length by about 0.13 mm during the day and a decrease of 0.04 mm during the night, whereas in form-deprived eyes axial length increases during both day and night, resulting in myopia primarily due to the inhibition of normal ocular shrinkage observed during the night.¹⁰⁰ Form-deprived chick eyes develop myopia and longer axial lengths, when reared under photoperiods of eight to 18 hours, whereas eyes remain hyperopic with shorter axial length under a photoperiod of 23 hours.¹⁰¹ These results suggest that chick eyes exhibit a photo period-dependant diurnal growth pattern. Similar diurnal rhythms have also been observed in marmosets.¹⁰² The diurnal growth pattern is age-dependant in marmosets, in which the eyes elongate by 25 μm during the day and decrease in length by 22 μm during the night in juveniles. The reverse is true for adolescents with axial length decreasing by 20 μm during the day and increasing by 38 μm during the night. Choroidal thickness also shows a diurnal rhythm with a decrease during the day and an increase during the night in both juvenile ($-12 \mu\text{m}$ versus $+18 \mu\text{m}$) and adolescent marmosets ($-22 \mu\text{m}$ versus $+21 \mu\text{m}$).¹⁰²

Diurnal variations in ocular growth have also been observed in young adults, with the longest axial length and vitreous chamber depth during the day and the shortest during the night. Choroidal thickness also displays

diurnal variations opposite in phase to that of axial length, with reduced thickness during the day and increased thickness during the night¹⁰³; however, season of birth is not a good measure of the overall lifetime effect of seasons on refractive error shifts.

ASSOCIATION BETWEEN PARENTAL SMOKING AND MYOPIA

Evidence from epidemiological and other studies

Several studies have shown that parental smoking, in particular maternal smoking during pregnancy, may influence refractive error development in children^{104–109} (Table 5).

Parental smoking during a child's lifetime is associated with a lower myopic prevalence,¹⁰⁴ less myopic refraction and shorter axial length compared to those without a history of maternal smoking.^{104,105} A recent population-based study of 3,009 Singapore Chinese children aged six to 72 months showed that the ORs for myopia were lower in children with a maternal and paternal history of smoking (adjusted ORs=0.50, 95 per cent CI: 0.30, 0.84; $p=0.01$; adjusted OR=0.72; 95 per cent CI: 0.54, 0.96, $p=0.02$, respectively) compared to those without.¹⁰⁶ A similar study of children from diverse ethnicities showed that a maternal history of smoking during pregnancy was significantly associated hyperopia greater than 2.00 D (adjusted OR=1.41; 95 per cent CI: 1.18, 1.69; $p<0.05$) compared to those who did not have a history of maternal smoking.¹⁰⁷ Another cross-sectional study of 300 Egyptian children aged five to 12 years¹⁰⁸ showed that urine cotinine levels (a biomarker for exposure to tobacco smoke) were significantly

higher in hyperopes ($64.46 \pm 15.82 \mu\text{g/l}$) compared to myopes ($51.96 \pm 18.66 \mu\text{g/l}$, $p<0.0001$) and emmetropes ($40.28 \pm 15.76 \mu\text{g/l}$) ($p<0.0001$). Conversely, a cross-sectional analysis of a British cohort of 2,487 subjects aged 44 years showed that a maternal history of smoking in early pregnancy was associated with increased risk of high myopia (adjusted relative risks [RRs] = 2.40, 95 per cent CI: 1.10, 5.40, $p<0.05$) compared to those without a maternal history of smoking.¹⁰⁹ The exact role of parental smoking in myopia development is still unknown. Studies that report a protective association between parental smoking and myopia either did not adjust for¹⁰¹ or inadequately adjusted for socio-economic status.^{104–107} Borchert and colleagues¹⁰⁷ did not adjust for socio-economic status, whereas other studies^{104–106} adjusted only for parental education and income. It should be noted that Rahi, Cumberland and Peckham¹⁰⁹ did not control for parental myopia, which is an important risk factor associated with myopia. In addition, there were few high myopes who reported a maternal history of smoking during early pregnancy.

Possible mechanism - evidence from animal studies

Exposure to tobacco smoke may influence ocular growth through nicotine, an important component of cigarette smoke that may influence myopia development through nicotinic acetylcholine receptors.^{110,111} The non-selective nicotinic antagonist drugs, namely chlorisondamine and mecamylamine, show the greatest efficacy in inhibiting axial growth and reducing myopic shift in refraction. Two of the nicotinic antagonists (methylcaconitine

and dihydro- β -erythroidine) inhibited myopic growth at high doses but the same effect has not been replicated in low doses, revealing multiphasic dose-response curves.¹¹⁰ Due to its complex nature of signalling mechanisms, the multiple nicotinic receptor subtypes, differing drug affinities with different receptor subtypes, potential biological effects of other constituents of tobacco smoke, the specific mechanism is yet to be defined.

Although human studies do not provide clear evidence for an association between passive smoking and myopia, animal studies have shown several antagonistic drugs to the neural types of nicotinic acetylcholine receptors which inhibit form-deprivation myopia in chicks.^{110,111}

ASSOCIATION BETWEEN BIRTH ORDER AND MYOPIA

Evidence from epidemiological studies

Evidence from epidemiological studies suggests that first-born individuals are more likely to be myopic.^{112,113} A meta-analysis of three British cohorts¹¹² showed that an increase in the number of older siblings was significantly associated with a reduced OR for a visual acuity of 6/12 or worse in both 10- and 11-year-olds (adjusted OR=0.89; 95 per cent CI: 0.83, 0.94; $p<0.01$) and 15- and 16-year-old subjects (adjusted OR=0.84; 95 per cent CI: 0.80, 0.89; $P<0.01$). A meta-analysis of four population-based cohort studies¹¹³ suggested that first-born subjects were more likely to be myopic compared to non-first-born subjects in the ALSPAC (adjusted OR=1.31; 95 per cent CI: 1.05, 1.64; $p=0.016$) and IDFC cohorts (adjusted OR=1.04, 95 per cent CI: 1.03, 1.06; $p<0.001$).

Author	Study design (location)	Sample size	Age range	Main findings
Iyer et al. ¹⁰⁶	Population based study (Singapore)	3009	6-72 months	Significantly lower ORs for myopia in children with history of parental smoking during pregnancy compared to those without
El Shazly ¹⁰⁸	Cross-sectional (Egypt)	300	5-12 years	Significantly higher urine cotinine levels in hyperopes compared to myopes
Borchert et al. ¹⁰⁷	Population-based cross-sectional (United States) (MEPDS and BPEDS [†])	9970	6-72 months	Maternal smoking during pregnancy was significantly associated with hyperopia $\geq +2.00\text{D}$
Rahi et al. ¹⁰⁹	Cohort study (United Kingdom)	2487	44 years	Maternal smoking during early pregnancy was significantly associated with high myopia

[†] Multi-Ethnic Pediatric Eye Disease and Baltimore Pediatric Eye Disease Studies

Table 5. Evidence for the association between parental smoking and myopia

Despite larger samples, these studies have some limitations. Rudnicka and colleagues¹¹² classified myopia based on unaided visual acuity, which is not a precise method. The use of non-cycloplegic auto-refraction and subjective refraction to determine refractive error may have overestimated the prevalence or magnitude of myopic refractive error in children and young adults.¹¹³ Thus, there may be a random misclassification of myopia in these studies, which could result in a spurious association between myopia and birth order.

Possible mechanisms

Possible mechanisms underlying the association between birth order and myopia include low birth weight, post-natal catch-up growth and insulin resistance. Most first-born babies tend to have low birth weight for their gestational age compared to second and third-born babies^{114,115} and are more likely to have intrauterine growth restraint, which leads to post-natal catch-up growth during the first two years of life.^{116–118} Such children are likely to have increased levels of plasma insulin and insulin resistance.^{118–120}

High levels of insulin may trigger myopia in a mechanism similar to that observed in chick eyes. Experiments in chicks have shown that intravitreal administration of insulin inhibits positive lens-induced hyperopia and accelerates negative lens-induced axial myopia through the inhibition of choroid thickening, elongation of the anterior chamber and crystalline lens thickening.^{121,122}

A large body of evidence shows a trend of increasing educational attainment with decreasing birth order, with first-born children attaining the highest educational level compared to non-first borns.^{123–129} Given the link between higher educational level and myopia^{74,130–136} and a significant interaction between myopic genetic loci and higher educational level,^{137,138} it is likely that myopia is more prevalent in firstborn individuals and education level may be a contributing factor.

ASSOCIATION BETWEEN MATERNAL AGE AND MYOPIA - EVIDENCE FROM EPIDEMIOLOGICAL STUDIES

Maternal age is another important early-life factor associated with myopia. Maternal age greater than 35 years increases the likelihood of varying degrees of myopia (adjusted OR: 1.5, 95 per cent CI: 1.1, 2.0, $p < 0.05$)¹⁰⁹ and reduced unaided distance vision (adjusted OR: 1.10, 95 per cent CI 1.04, 1.17; $p < 0.01$).¹¹² Maternal age may be linked to myopia

as older mothers are more likely to give birth to low birth weight infants¹³⁹ and myopia is associated with low birth weight.¹⁰⁹

CONCLUSION

Population-based data show a consistent protective association between time outdoors and myopia. Results from clinical trials for outdoor intervention programs to reduce incident myopia are promising. Increasing time outdoors during childhood and adolescence could be accomplished through conducting lessons outside the classroom, incorporating outdoor activity in the school curriculum or through structured weekend outdoor programs for families. Evidence for the association of near work with myopia is not as robust as time outdoors and may be difficult to quantify. Experimental models and epidemiological associations suggest a role for neural nicotinic acetylcholine receptors on ocular growth but more studies are warranted.

REFERENCES

- Pan CW, Zhen YF, Anuar AR, Chew M, Gazzard G, Aung T et al. Prevalence of refractive errors in a multi-ethnic Asian population: The Singapore Epidemiology of Eye Disease Study. *Invest Ophthalmol Vis Sci* 2013; 54: 2590–2598.
- Pan CW, Klein BE, Cotch MF, Shrager S, Klein R, Folsom A et al. Racial variations in the prevalence of refractive errors in the United States: The Multi-Ethnic study of Atherosclerosis. *Am J Ophthalmol* 2013; 155: 1129–1138.
- Wu SY, Nemesure B, Leske MC. Refractive errors in a black adult population: The Barbados Eye Study. *Invest Ophthalmol Vis Sci* 1999; 40: 2179–2184.
- Attebo K, Ivers RQ, Mitchell P. Refractive errors in an older population: the Blue Mountains Eye Study. *Ophthalmology* 1999; 106: 1066–1072.
- Saw SM, Shankar A, Tan SB, Taylor H, Tan DT, Stone RA et al. A cohort study of incident myopia in Singaporean children. *Invest Ophthalmol Vis Sci* 2006; 47: 1839–1844.
- Saw SM, Tong L, Chua WH, Chia KS, Koh D, Tan DT et al. Incidence and progression of myopia in Singaporean school children. *Invest Ophthalmol Vis Sci* 2005; 46: 51–57.
- He M, Huang W, Zheng Y, Huang L, Ellwein LB. Refractive error and visual impairment in school children in rural southern China. *Am J Ophthalmol* 2000; 129: 427–435.
- Zadnik K, The Glenn A. Fry award lecture. Myopia development in childhood. *Optom Vis Sci* 1997; 74: 603–608.
- Ip JM, Huynh SC, Robaei D, Kifley A, Rose KA, Morgan IG et al. Ethnic differences in refraction and ocular biometry in population-based sample of 11–15 year old Australian children. *Eye* 2008; 22: 649–656.
- Dandona R, Dandona L, Srinivas M, Sahare P, Narsaiah S, Muñoz SR et al. Refractive error in children in a rural population in India. *Invest Ophthalmol Vis Sci* 2002; 43: 615–622.
- Murthy GV, Gupta SK, Ellwein LB, Muñoz SR, Pokharel GP, Sanga L et al. Refractive error in children in an urban population in New Delhi. *Invest Ophthalmol Vis Sci* 2002; 43: 623–631.
- Pokharel GP, Negrel AD, Munoz SR, Ellwein LB. Refractive Error Study in Children: results from Mechi Zone, Nepal. *Am J Ophthalmol* 2000; 129: 436–444.
- Lim MC, Gazzard G, Sim EL, Tong L, Saw SM. Direct costs of myopia in Singapore. *Eye* 2009; 23: 1086–1089.
- Zheng YF, Pan CW, Chay J, Wong TY, Finkelstein E, Saw SM. The economic cost of myopia in adults aged aged over 40 years in Singapore. *Invest Ophthalmol Vis Sci* 2013; 54: 7532–7537.
- Javitt JC, Chiang YP. The socioeconomic aspects of laser refractive surgery. *Arch Ophthalmol* 1994; 112: 1526–1530.
- Chang L, Pan CW, Ohno-Matsui L, Lin X, Cheung G, Gazzard G et al. Myopia related fundus changes in Singapore adults with high myopia. *Am J Ophthalmol* 2013; 155: 991–999.
- Samarawickrama C, Mitchell P, Tong L, Gazzard G, Lim L, Wong TY et al. Myopia related optic disc and retinal changes in adolescent children from Singapore. *Ophthalmology* 2011; 118: 2050–2057.
- Saw SM, Gazzard G, Shih-Yen EC, Chua WH. Myopia and associated pathological complications. *Ophthalmic Physiol Opt* 2005; 25: 381–391.
- Morgan IG, Xiang F, Zeng Y, Mai J, Chen Q, Zhang J et al. Increased outdoor time reduces the risk of myopia - the Guangzhou Outdoor Activity Longitudinal study. *Invest Ophthalmol Vis Sci* 2014; 55: E-Abstract 1272.
- Guggenheim JA, Northstone K, McMahon G, Ness AR, Deere K, Mattocks C et al. Time outdoors and physical activity as predictors of incident myopia in childhood: a prospective cohort study. *Invest Ophthalmol Vis Sci* 2012; 53: 2856–2865.
- French AN, Morgan IG, Mitchell P, Rose KA. Risk factors for incident myopia in Australian schoolchildren: the Sydney adolescent vascular and eye study. *Ophthalmology* 2013; 120: 2100–2108.
- French AN, Ashby RS, Morgan IG, Rose KA. Time outdoors and the prevention of myopia. *Exp Eye Res* 2013; 114: 58–68.
- Wojciechowski R. Nature and nurture: the complex genetics of myopia and refractive error. *Clin Genet* 2011; 79: 301–320.
- Stambolian D. Genetic susceptibility and mechanisms for refractive error. *Clin Genet* 2013; 84: 102–108.
- Baird PN, Schache M, Dirani M. The Genes in Myopia (GEM) study in understanding the aetiology of refractive errors. *Prog Retin Eye Res* 2010; 29: 520–542.
- Rose KA, Morgan IG, Smith W, Burlutsky G, Mitchell P, Saw SM. Myopia, lifestyle, and schooling in students of Chinese ethnicity in Singapore and Sydney. *Arch Ophthalmol* 2008; 126: 527–530.
- Rose KA, Morgan IG, Ip J, Kifley A, Huynh S, Smith W et al. Outdoor activity reduces the prevalence of myopia in children. *Ophthalmology* 2008; 115: 1279–1285.
- Dirani M, Tong L, Gazzard G, Zhang X, Chia A, Young TL et al. Outdoor activity and myopia in Singapore teenage children. *Br J Ophthalmol* 2009; 93: 997–1000.
- Wu PC, Tsai CL, Hu CH, Yang YH. Effects of outdoor activities on myopia among rural school children in Taiwan. *Ophthalmic Epidemiol* 2010; 17: 338–342.

30. Jones LA, Sinnott LT, Mutti DO, Mitchell GL, Moeschberger ML, Zadnik K. Parental history of myopia, sports and outdoor activities, and future myopia. *Invest Ophthalmol Vis Sci* 2007; 48: 3524–3532.
31. Jones-Jordan LA, Mitchell GL, Cotter SA, Kleinstein RN, Manny RE, Mutti DO et al. Visual activity before and after the onset of juvenile myopia. *Invest Ophthalmol Vis Sci* 2011; 52: 1841–1850.
32. Jones-Jordan LA, Sinnott LT, Cotter SA, Kleinstein RN, Manny RE, Mutti DO et al. Time outdoors, visual activity, and myopia progression in juvenile-onset myopes. *Invest Ophthalmol Vis Sci* 2012; 53: 7169–7175.
33. R Dharani, Lee C-F, Theng ZX, Drury VB, Ngo C, Sandar M et al. Comparison of measurements of time outdoors and light levels as risk factors for myopia in young Singapore children. *Eye* 2012; 26: 911–918.
34. Read SA, Collins MJ, Vincent SJ. Light exposure and physical activity in myopic and emmetropic children. *Optom Vis Sci* 2014; 91: 330–341.
35. McKnight CM, Sherwin JC, Yazar S, Forward H, Tan AX, Hewitt AW et al. Myopia in young adults is inversely related to an objective marker of ocular sun exposure: The Western Australian Raine Cohort Study. *Am J Ophthalmol* 2014; 158: 1079–1085.
36. Sherwin JC, Hewitt AW, Coroneo MT, Kearns LS, Griffiths LR, Mackey DA. The association between time spent outdoors and myopia using a novel bio marker of outdoor exposure. *Invest Ophthalmol Vis Sci* 2012; 53: 4363–4370.
37. Schmid KL, Leyden K, Chiu YH, Lind SR, Vos DJ, Kimlin M et al. Assessment of daily light and ultra violet exposure in young adults. *Optom Vis Sci* 2013; 90: 148–155.
38. Guo Y, Liu LJ, Xu L, Lv YY, Tang P, Feng Y et al. Outdoor activity and myopia among primary students in rural and urban regions of Beijing. *Ophthalmology* 2013; 120: 277–283.
39. Lin Z, Vasudevan B, Jhanji V, Mao GY, Gao TY, Wang FH et al. Near work, outdoor activity, and their association with refractive error. *Optom Vis Sci* 2014; 91: 376–382.
40. Guo Y, Liu LJ, Xu L, Tang P, Lv YY, Feng Y et al. Myopic shift and outdoor activity among primary school children: one-year follow-up study in Beijing. *PLoS One* 2013; 8: e75260.
41. Lu B, Congdon N, Liu X, Choi K, Lam DSC, Zhang M et al. Associations between near work, outdoor activity and myopia among adolescent students in rural China. *Arch Ophthalmol* 2009; 127: 769–775.
42. Fujiwara M, Hasebe S, Nakanishi R, Tanigawa K, Ohtsuki H. Seasonal variation in myopia progression and axial elongation: an evaluation of Japanese children participating in a myopia control trial. *Jpn J Ophthalmol* 2012; 56: 401–406.
43. Donovan L, Sankaridurg P, Ho A, Chen X, Lin Z, Thomas V et al. Myopia progression in Chinese children is slower in summer than in winter. *Optom Vis Sci* 2012; 89: 1196–1202.
44. Cui D, Trier K, Munk Ribel-Madsen S. Effect of day length on eye growth, myopia progression, and change of corneal power in myopic children. *Ophthalmology* 2013; 120: 1074–1079.
45. Gwiazda J, Deng L, Manny R, Norton TT, COMET Study Group. Seasonal variations in the progression of myopia in children enrolled in the correction of myopia evaluation trial. *Invest Ophthalmol Vis Sci* 2014; 55: 752–758.
46. Deng L, Gwiazda J, Thorn F. Children's refractions and visual activities in the school year and summer. *Optom Vis Sci* 2010; 87: 406–413.
47. Thorne HC, Jones KH, Peters SP, Archer SN, Dijk DJ. Daily and seasonal variations in the spectral composition of light exposure humans. *Chronobiol Int* 2009; 26: 854–866.
48. Wu PC, Tsai CL, Wu HL, Yang YH, Kuo HK. Outdoor activity during class recess reduces myopia onset and progression in school children. *Ophthalmology* 2013; 120: 1080–1085.
49. Ngo CS, Pan CW, Finkelstein EA, Lee CF, Wong IB, Ong J et al. A cluster randomized controlled trial evaluating an incentive-based outdoor physical activity programme to increase outdoor activity and prevent myopia in children. *Ophthalmic Physiol Opt* 2014; 34: 362–368.
50. Ashby R, Ohlendorf A, Schaeffel F. The effect of ambient illuminance on the development of deprivation myopia in chicks. *Invest Ophthalmol Vis Sci* 2009; 50: 5348–5354.
51. Cohen Y, Peleg E, Belkin M, Polat U, Solomon AS. Ambient illuminance, retinal dopamine release and refractive development in chicks. *Exp Eye Res* 2012; 103: 33–40.
52. Backhouse S, Collins AV, Phillips JR. Influence of periodic vs. continuous daily bright light exposure on development of experimental myopia in the chick. *Ophthalmic Physiol Opt* 2013; 33: 563–572.
53. Ashby RS, Schaeffel F. The effect of bright light on lens compensation in chicks. *Invest Ophthalmol Vis Sci* 2010; 51: 5247–5253.
54. Jiang L, Long K, Schaeffel F, Zhou X, Zheng Y, Ying H et al. Effects of dopaminergic agents on progression of naturally occurring myopia in albino guinea pigs. *Invest Ophthalmol Vis Sci* 2014; 55: 7508–7519.
55. Smith EL 3rd, Hung LF, Huang J. Protective effects of high ambient lighting on the development of form deprivation myopia in rhesus monkeys. *Invest Ophthalmol Vis Sci* 2012; 53: 421–428.
56. Smith EL 3rd, Hung LF, Arumugam B, Huang J. Negative lens induced myopia in infant monkeys: effects of high ambient lighting. *Invest Ophthalmol Vis Sci* 2013; 54: 2959–2969.
57. Sherwin JC, Reacher MH, Keogh RH, Khawaja AP, Mackey DA, Foster PJ. The association between time spent outdoors and myopia in children and adolescents: a systematic review and meta-analysis. *Ophthalmology* 2012; 119: 2141–2151.
58. Norton TT, Siegwart JT Jr. Light levels, refractive development and myopia. A speculative review. *Exp Eye Res* 2013; 114: 48–57.
59. Feldkaemper M, Schaeffel F. An updated view on the role of dopamine in myopia. *Exp Eye Res* 2013; 114: 106–119.
60. Schwahn HN, Schaeffel F. Flicker parameters are different for suppression of myopia and hyperopia. *Vision Res* 1997; 37: 2661–2673.
61. Rucker FJ, Wallman J. Chicks use changes in luminance and chromatic contrast as indicators of the sign of defocus. *J Vis* 2012; 12: 1–13.
62. Long Q, Chen D, Chu R. Illumination with monochromatic long-wavelength light promotes myopic shift and axial elongation in new born pigmented guinea pigs. *Cutan Ocul Toxicol* 2009; 28: 176–180.
63. Liu R, Qian YF, He JC, Hu M, Zhou XT, Dai JH et al. Effects of different monochromatic lights on refractive development and eye growth in guinea pigs. *Exp Eye Res* 2011; 92: 447–453.
64. Jiang L, Zhang S, Schaeffel F, Xiong S, Zheng Y, Zhou X et al. Interactions of chromatic and lens induced-defocus during visual control of eye growth in guinea pigs. *Vision Res* 2014; 94: 24–32.
65. Foulds WS, Barathi VA, Luu CD. Progressive myopia or hyperopia can be induced in chicks and reversed by manipulation of the chromaticity of the ambient light. *Invest Ophthalmol Vis Sci* 2013; 54: 8004–8012.
66. Holick MF. Sunlight and vitamin D for bone health and prevention of autoimmune diseases, cancers, and cardiovascular disease. *Am J Clin Nutr* 2004; 80 (6 suppl): 1678S–1688S.
67. Mutti DO, Marks AR. Blood levels of vitamin D in teens and young adults with myopia. *Optom Vis Sci* 2011; 88: 377–382.
68. Choi JA, Han K, Park YM, La TY. Low serum 25-hydroxyvitamin D is associated with myopia in Korean adolescents. *Invest Ophthalmol Vis Sci* 2014; 55: 2041–2047.
69. Yazar S, Hewitt AW, Black LJ, McKnight CM, Mountain JA, Sherwin JC et al. Myopia is associated with lower vitamin D status in young adults. *Invest Ophthalmol Vis Sci* 2014; 55: 4552–4559.
70. Guggenheim JA, Williams C, Northstone K, Howe LD, Tilling K, St Pourcain B et al. Does vitamin D mediate the protective effects of time outdoors on myopia? Findings from a prospective birth cohort. *Invest Ophthalmol Vis Sci* 2014; 55: 8550–8558.
71. Mutti DO, Cooper ME, Dragan E, Jones-Jordan LA, Bailey MD, Marazita ML et al. Vitamin D receptor (VDR) and group-specific component (GC, vitamin D-binding protein) polymorphisms in myopia. *Invest Ophthalmol Vis Sci* 2011; 52: 3818–3824.
72. Annamalai S, Bindu CH, Reddy KP, Vishnupriya S. Association of vitamin D receptor gene start codon (FokI) polymorphism with high myopia. *Oman J Ophthalmol* 2011; 4: 57–62.
73. Flitcroft DI. The complex interactions of retinal, optical and environmental factors in myopia aetiology. *Prog Retin Eye Res* 2012; 31: 622–660.
74. Mutti DO, Mitchell GL, Moeschberger ML, Jones LA, Zadnik K. Parental myopia, near work, school achievement and children's refractive error. *Invest Ophthalmol Vis Sci* 2002; 43: 3633–3640.
75. Ip JM, Rose KA, Morgan IG, Burlutsky G, Mitchell P. Myopia and the urban environment: findings in a sample of 12-year-old Australian schoolchildren. *Invest Ophthalmol Vis Sci* 2008; 49: 3858–3863.
76. You QS, Wu LJ, Duan JL, Luo YX, Liu LJ, Li X et al. Factors associated with myopia in children in China: the Beijing childhood Eye Study. *PLoS One* 2012; 7: e52668.
77. Lee YY, Lo CT, Sheu SJ, Lin JL. What factors are associated with myopia in young adults? A survey study in Taiwan military conscripts. *Invest Ophthalmol Vis Sci* 2013; 54: 1026–1033.
78. Gong Y, Zhang X, Tian D, Wang D, Xiao G. Parental myopia, near work, hours of sleep and myopia in Chinese children. *Health* 2014; 6: 64–70.
79. Wallman J, Gottlieb MD, Rajaram NV, Fugate-Wentzek LA. Local retinal regions control local eye growth and myopia. *Science* 1987; 237: 73–77.
80. Troilo D, Gottlieb MD, Wallman J. Visual deprivation causes myopia in chicks with optic nerve section. *Curr Eye Res* 1987; 6: 993–999.
81. Choh V, Lew MY, Nadel MW, Wildsoet CF. Effects of interchanging hyperopic defocus and form deprivation stimuli in normal and optic nerve sectioned chicks. *Vision Res* 2006; 46: 1070–1079.
82. Tse DY, Lam CS, Guggenheim JA, Lam C, Li KK, Liu Q et al. Simultaneous defocus integration during refractive development. *Invest Ophthalmol Vis Sci* 2007; 48: 5352–5359.

83. Smith EL, Bradley DV, Fernandes A, Booth RG. Form deprivation myopia in adolescent monkeys. *Optom Vis Sci* 1999; 76: 428–432.
84. Smith EL, Hung LF. Form-deprivation myopia in monkeys is a graded phenomenon. *Vision Res* 2000; 40: 371–381.
85. Smith EL, Hung LF, Kee CS, Qiao Y. Effects of brief periods of unrestricted vision on the development of form-deprivation myopia in monkeys. *Invest Ophthalmol Vis Sci* 2002; 43: 291–299.
86. Winawer J, Wallman J. Temporal constraints in lens compensation in chicks. *Vision Res* 2002; 42: 2651–2668.
87. McBrien NA, Young TL, Pang CP, Hammond C, Baird P, Saw SM et al. Myopia: recent advances in molecular studies; prevalence, progression and risk factors; emmetropization; therapies; optical links; peripheral refraction; sclera and ocular growth; signalling cascades; and animal models. *Optom Vis Sci* 2009; 86: 45–66.
88. Wildsoet CF, Wallman J. Choroidal and scleral mechanisms of compensation for spectacle lenses in chicks. *Vision Res* 1995; 35: 1175–1194.
89. Hung LF, Wallman J, Smith EL. Vision-dependent changes in the choroidal thickness of macaque monkeys. *Invest Ophthalmol Vis Sci* 2000; 41: 1259–1269.
90. Troilo D, Nickla DL, Wildsoet CF. Choroidal thickness changes during altered eye growth and refractive state in a primate. *Invest Ophthalmol Vis Sci* 2000; 41: 1249–1258.
91. Goss DA, Wickham MG. Retinal image mediated ocular growth as mechanism for juvenile onset myopia and emmetropization. A review. *Doc Ophthalmol* 1995; 90: 341–375.
92. Zadnik K, Mutti DO. How applicable are animal myopia models to human juvenile onset myopia? *Vision Res* 1995; 35: 1283–1288.
93. Grosvenor T, Goss DA. *Clinical Management of Myopia*. Oxford: Butterworth-Heinemann Publishers, 1999.
94. Hung GK, Ciuffreda KJ. Incremental retinal defocus theory of myopia development - Schematic analysis and computer simulation. *Comput Biol Med* 2007; 37: 930–946.
95. Mandel Y, Grotto I, El-Yaniv R, Belkin M, Israeli E, Polat U et al. Season of birth, natural light, and myopia. *Ophthalmology* 2008; 115: 686–692.
96. McMahon G, Zayats T, Chen YP, Prashar A, Williams C, Guggenheim JA. Season of birth, daylight hours at birth, and high myopia. *Ophthalmology* 2009; 116: 468–473.
97. Deng L, Gwiazda J. Birth season, photoperiod, and infancy refraction. *Optom Vis Sci* 2011; 88: 383–387.
98. Vannas AE, Ying GS, Stone RA, Maguire MG, Jormanainen V, Tervo T. Myopia and natural lighting extremes: risk factors in Finnish army conscripts. *Acta Ophthalmol Scand* 2003; 81: 588–595.
99. Ma Q, Xu W, Zhou X, Cui C, Pan CW. The relationship of season of birth with refractive error in very young children in Eastern China. *PLoS One* 2014; 9: e100472.
100. Weiss S, Schaeffel F. Diurnal growth rhythms in the chicken eye: relation to myopia development and retinal dopamine level. *J Comp Physiol A* 1993; 172: 263–270.
101. Stone RA, Lin T, Desai D, Capehart C. Photoperiod, early post-natal eye growth, and visual deprivation. *Vision Res* 1995; 35: 1195–1202.
102. Nickla DL, Wildsoet CF, Troilo D. Diurnal rhythms in intraocular pressure, axial length, and choroidal thickness in a primate model of eye growth, the common marmoset. *Invest Ophthalmol Vis Sci* 2002; 43: 2519–2528.
103. Chakraborty R, Read SA, Collins MJ. Diurnal variations in axial length, choroidal thickness, intraocular pressure and ocular biometrics. *Invest Ophthalmol Vis Sci* 2011; 52: 5121–5129.
104. Saw SM, Chia KS, Lindstrom JM, Tan DT, Stone RA. Childhood myopia and parental smoking. *Br J Ophthalmol* 2004; 88: 934–937.
105. Stone RA, Wilson LB, Ying GS, Liu C, Criss JS, Orlov J et al. Associations between childhood refraction and parental smoking. *Invest Ophthalmol Vis Sci* 2006; 47: 4277–4287.
106. Iyer JV, Low WC, Dirani M, Saw SM. Parental smoking and childhood refractive error: the STARS study. *Eye* 2012; 26: 1324–1328.
107. Borchert MS, Varma R, Cotter SA, Tarczy-Hornoch K, McKean-Cowdin R, Lin JH et al. Risk factors for hyperopia and myopia in preschool children: the multi-ethnic pediatric eye disease and Baltimore pediatric eye disease studies. *Ophthalmology* 2011; 118: 1966–1973.
108. El-Shazly AA. Passive smoking exposure might be associated with hypermetropia. *Ophthalmic Physiol Opt* 2012; 32: 304–307.
109. Rahi JS, Cumberland PM, Peckham CS. Myopia over the life course: prevalence and early life influences in the 1958 British birth cohort. *Ophthalmology* 2011; 118: 797–804.
110. Stone RA, Sugimoto R, Gill AS, Liu J, Capehart C, Lindstrom JM. Effects of nicotinic antagonists on ocular growth and experimental myopia. *Invest Ophthalmol Vis Sci* 2001; 42: 557–565.
111. Stone RA, Pardue MT, Iuvone PM, Khurana TS. Pharmacology of myopia and potential role for intrinsic retinal circadian rhythms. *Exp Eye Res* 2013; 114: 35–47.
112. Rudnicka AR, Owen CG, Richards M, Wadsworth ME, Strachan DP. Effect of breastfeeding and socio-demographic factors on visual outcome in childhood and adolescence. *Am J Clin Nutr* 2008; 87: 1392–1399.
113. Guggenheim JA, McMahon G, Northstone K, Mandel Y, Kaiserman I, Stone RA et al. Birth order and myopia. *Ophthalmic Epidemiol* 2013; 20: 375–384.
114. Ghaemmaghami SJ, Nikniaz L, Mahdavi R, Nikniaz Z, Razmifard F, Afsharman F. Effects of infants' birth order, maternal age, and socio-economic status on birth weight. *Saudi Med J* 2013; 34: 949–953.
115. Ong KK, Preece MA, Emmett PM, Ahmed ML, Dunger DB, ALSPAC Study Team. Size at birth and early childhood growth in relation to maternal smoking, parity and infant breast-feeding: longitudinal birth cohort study and analysis. *Pediatr Res* 2002; 52: 863–867.
116. Ong KK, Ahmed ML, Emmett PM, Preece MA, Dunger DB. Association between postnatal catch-up growth and obesity in childhood: prospective cohort study. *Br Med J* 2000; 320: 967–971.
117. Ong KK, Dunger DB. Birth weight, infant growth and insulin resistance. *Eur J Endocrinol* 2004; 151: 131–139.
118. Ong KK, Petry CJ, Emmett PM, Sandhu MS, Kiess W, Hales CN et al. Insulin sensitivity and secretion in normal children related to size at birth, postnatal growth, and plasma insulin-like growth factor-I levels. *Diabetologia* 2004; 47: 1064–1070.
119. Bavdekar A, Yajnik CS, Fall CH, Bapat S, Pandit AN, Deshpande V et al. Insulin resistance syndrome in 8-year-old Indian children: small at birth, big at 8 years, or both? *Diabetes* 1999; 48: 2422–2429.
120. Soto N, Bazaes RA, Peña V, Salazar T, Avila A, Iniguez G et al. Insulin sensitivity and secretion are related to catch-up growth in small-for-gestational age infants at age 1-year: results from a prospective cohort. *J Clin Endocrinol Metab* 2003; 88: 3645–3650.
121. Feldkaemper MP, Neacsu I, Schaeffel F. Insulin acts as a powerful stimulator of axial myopia in chicks. *Invest Ophthalmol Vis Sci* 2009; 50: 13–23.
122. Zhu X, Wallman J. Opposite effects of glucagon and insulin on compensation for spectacle lenses in chicks. *Invest Ophthalmol Vis Sci* 2009; 50: 24–36.
123. Hauser RM, Sewell WH. Birth order and educational attainment in full sibships. *Am Educ Res J* 1985; 22: 1–23.
124. Booth AL, Kee HJ. Birth order matters: the effect of family size and birth order on educational attainment. *J Popul Econ* 2006; 22: 367–397.
125. Fergusson DM, Horwood LJ, Boden JM. Birth order and educational achievement in adolescence and young adulthood. *Au J Educ* 2006; 50: 122–139.
126. De Haan M. Birth order, family size and educational attainment. *Econ Educ Rev* 2010; 29: 576–588.
127. Harkonen J. Birth Order Effects on Educational Attainment and Educational Transitions in West Germany. *Eu Sociol Rev* 2014; 30: 166–179.
128. Black SE, Devereux PJ, Salvanes KG. The more the merrier? The effect of family size and birth order on children's education. *The Quart J Econ* 2005; 120: 669–700.
129. Clayton S, Apperley E, Hannon F, Karia A, Baxter V, Julious SA. A survey of birth order status of students studying for medical degree at the University of Sheffield. *JRSM Open* 2014; 5: 1–5.
130. Morgan IG, Cotch MF. Birth Order and Myopia: What are the Messages to Readers? *Ophthalmic Epidemiol* 2013; 20: 333–334.
131. Teasdale TW, Fuchs J, Goldschmidt E. Degree of myopia in relation to intelligence and educational level. *Lancet* 1988; 2: 1351–1354.
132. Tay MT, Au Eong KG, Ng CY, Lim MK. Myopia and educational attainment in 421,116 young Singaporean males. *Ann Acad Med Singapore* 1992; 21: 785–791.
133. Saw SM, Wu HM, Seet B, Wong TY, Yap E, Chia KS et al. Academic achievement, close up work parameters, and myopia in Singapore military conscripts. *Br J Ophthalmol* 2001; 85: 855–860.
134. Konstantopoulos A, Yadegarfar G, Elgohary M. Near work, education, family history and myopia in Greek conscripts. *Eye* 2008; 22: 542–546.
135. Gwiazda J, Deng L, Dias L, Marsh-Tootle W, COMET Study Group. Association of education and occupation with myopia in COMET parents. *Optom Vis Sci* 2011; 88: 1045–1053.
136. Jung SK, Lee JH, Kakizaki H, Jee D. Prevalence of myopia and its association with body stature and educational levels in 19-year-old male conscripts in Seoul, South Korea. *Invest Ophthalmol Vis Sci* 2012; 53: 5579–5583.
137. Verhoeven VJ, Buitendijk GH, Consortium for Refractive Error and Myopia (CREAM), Rivadeneira F, Uitterlinden AG, Vingerling JR et al. Education influences the role of genetics in myopia. *Eur J Epidemiol* 2013; 28: 973–980.
138. Fan Q, Wojciechowski R, Kamran Ikram M, Cheng CY, Chen P, Zhou X et al. Education influences the association between genetic variants and refractive error: a meta-analysis of five Singapore studies. *Hum Mol Genet* 2014; 23: 546–554.
139. Dennis JA, Mollborn S. Young maternal age and low birth weight risk: An exploration of racial/ethnic disparities in the birth outcomes of mothers in the United States. *Soc Sci J* 2013; 50: 625–634.