

RESEARCH

Open Access



Prevalence and pattern of refractive error and visual impairment among schoolchildren: the Lhasa childhood eye study

Jiantao Cui¹, Jing Fu^{1*}, Lei Li¹, Weiwei Chen^{1,2}, Zhaojun Meng¹, Han Su¹, Yao Yao¹ and Wei Dai¹

Abstract

Background: Early and effective ocular screening may help to eliminate treatable eye disorders. The Lhasa Childhood Eye Study (LCES) revealed the particular prevalence of refractive error and visual impairment in grade one schoolchildren (starting age of 6 years old) in Lhasa.

Methods: This is a cross-sectional part of school-based cohort study. One thousand nine hundred forty-three children were enrolled (median age, 6.78 years, range, 5.89 to 10.32). Each child underwent general and ocular examinations, including logarithm of the minimum angle of resolution (logMAR) visual acuity, cycloplegic autorefraction, and slit-lamp biomicroscopy evaluation. Multivariate and correlation analyses were performed to evaluate the association between refractive error with gender and ethnics.

Results: The prevalence of visual impairment (logMAR visual acuity ≥ 0.3 in the better-seeing eye) of uncorrected, presenting and best-corrected visual acuity (BCVA) was 12.2, 11.7 and 2.7%, respectively. Refractive error presented in 177 (78.0%) out of 227 children with bilateral visual impairment. Myopia (spherical equivalent refractor [SER] ≤ -0.50 diopter [D] in either eye) was present in 4.7% children when measured after cycloplegic autorefraction. Hyperopia (SER $\geq +2.00$ D) affected 12.1% children. Hyperopia was significantly associated with female gender ($P < 0.001$). Astigmatism (cylinder value ≤ -0.75 D) was present in 44.8% children. In multivariate regression and correlation analysis, SER had no significant difference between ethnic groups.

Conclusion: The Lhasa Childhood Eye Study is the first school-based cohort study to reveal the prevalence and pattern of refractive error and visual impairment in Lhasa. Effective strategies such as corrective spectacles should be considered to alleviate treatable visual impairment.

Keywords: Refractive error, Visual impairment, School children, Myopia, Lhasa

Background

Vision impairment caused by refractive error, strabismus, amblyopia and other visual disabilities in children is relatively increased in recent years [1]. Pediatric vision

screening is important in the early detection of children who have reduced visual acuity or risk factors that endanger eye health. Effective vision screening enables early treatment of preventable and curable ocular diseases. World Health Organization, American Academy of Ophthalmology and other ophthalmic associations recommend early and regular vision screening throughout childhood [2].

Among the numerous visual impairment disorders, uncorrected refractive error leads to a key problem of heavy burden on learning, employment, and quality of

*Correspondence: fu_jing@126.com

¹ Beijing Tongren Eye Center, Beijing Tongren Hospital, Capital Medical University, Beijing Key Laboratory of Ophthalmology & Visual Sciences, No.1, Dong Jiao Min Xiang Street, Dongcheng District, Beijing 100730, China

Full list of author information is available at the end of the article



© The Author(s) 2021. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

life [3]. Thus, more attention is being paid to prevent and alleviate uncorrected refractive error especially in school children. More important, other ocular conditions, such as amblyopia and strabismus, will obtain favorable therapeutic effect being treated at early age. Worldwide, the prevalence of visual impairment among school-age children differs due to country, region, ethnicity, social development level, medical resources, etc. [1, 4] Evidence shown that refractive error is a concerning issue in China. And different prevalence of refractive error was reported in different regions throughout China [5, 6]. Tibetan adolescents were reported have a lower prevalence of refractive errors compared to China inland without cycloplegia [7]. However, there is no school-based cohort study about school children refraction status follow up from first grade in Tibetan have been published. Reliable evidence is necessary for local public health strategy conducting.

Tibet is characterized by its high-altitude (an average of >3500m), low air pressure and oxygen concentration, intense solar infrared and ultra-violet radiation, which are obviously different from China inland. Other distinguishing features of Tibetan include a relative lower education level than China inland, lower population mobility and easy to follow up. The differences of regional environment and economics are reported to have influence on human body development and disease composition [8]. Whether the above unique differences affect pediatric eye diseases is not known until recently.

The ethnicity of elementary school children in Lhasa, the provincial capital of Tibet, mainly consists of Tibetan (90.7%), Han (8.2%) and other minority nationality (1.1%). Lhasa Childhood Eye Study is a school-based cohort study mainly designed to longitudinally observe the occurrence and development of different ocular diseases especially refractive error and visual impairment in school-aged children in Lhasa [9].

Methods

Informed consent and ethics approval

This research was reviewed by an independent ethical review board and conforms with the principles and applicable guidelines for the protection of human subjects in biomedical research. The protocols in this study were approved by the Institutional Review Board of Beijing Tongren Hospital, Capital Medical University (TRECKY2019–146), and adhere to the principles of the Declaration of Helsinki. This present study is part of Lhasa Childhood Eye Study, which has registered on Chinese Clinical Trial Registry (<http://www.chictr.org.cn>, ChiCTR1900026693). Lhasa Childhood Eye Study is mainly designed to investigate the prevalence and associated factors longitudinally of several ocular disorders in school-age children in Lhasa for continuous 5 years.

Written informed consent was obtained from a parent or legal guardian of each child prior to the examinations.

Study setting and population

Stratified random cluster sampling was employed in selecting schools and eligible children for participation. Elementary schools were stratified into three levels based on the evaluation of local government. A total of 1943 grade one students from 14 randomly selected classes represented 7 of the 27 elementary schools available joining the Lhasa Childhood Eye Study.

An experienced clinical team comprising 2 optometrists and 3 ophthalmologists from Beijing Tongren Hospital performed all examination procedures during the period from October to November 2019. All the examinations were performed in the health examination station of Lhasa Maternal and Child Health Care Center.

As Tibetan children were reported to have a lower prevalence of refractive errors than inland, we use a cumulative incidence of 40% to calculate the sample size [7, 9]. Assuming a design effect of 2.0, a tolerated error of 0.1 times the myopia incidence, and a loss of follow-up of 20%, a total of 1382 grade one students would be needed. Further detail of the method is available elsewhere [9].

Ocular examinations

Demographic data (including ethnicity, age and gender) was collected before examination. All participants underwent standardized ocular examinations included distant visual acuity and identification of visual impairment. Quality controls were implemented throughout the entire study. The detailed standard operation procedures was reported elsewhere [9].

Distant visual acuity

The ophthalmic examinations included measures of visual acuity and refraction. Distant visual acuity of both eyes was measured at 3m using a Lea Symbols ETDRS 3-m Set charts (250,300, Goodlite, IL, USA). Children were examined monocularly, right eye followed by left eye. A letter-by-letter logMAR visual acuity score was documented and calculated accordingly. For children wearing glasses, both presenting visual acuity (with glasses) and uncorrected visual acuity (without glasses) were measured. For students with unaided visual acuity 0.3 or worse in either eye, subjective refraction was performed to obtain best-corrected visual acuity by trained optometrists. The cycloplegic procedure referred to the Anyang Childhood Eye Study [10]. Cycloplegia was achieved with two drops of 1% cyclopentolate and 1 drop of Mydrin P at a 5-min interval. Refractive status was measured using an autorefractor (KR-800, Topcon, Tokyo, Japan) before and after cycloplegia.

Assessment of visual impairment

The prevalence of visual impairment and blindness were calculated using uncorrected (unaided), presenting, and best-corrected visual acuity [11]. Visual acuities evaluated via logMAR categories were defined as normal/near normal (≤ 0.2 in both eyes), unilateral visual impairment (0.2 in one eye only), mild visual impairment (0.3–0.5 in the better eye), moderate visual impairment (0.6–0.9 in the better eye), and blindness (≥ 1.0 in the better eyes). The causes of visual impairment of 0.3 or worse were assessed by the ophthalmologist.

Definitions

Diagnosis was based on the following definition: (1) Refractive error was assigned routinely if visual acuity improved to ≤ 0.2 with refractive correction. (2) Myopia was defined as spherical equivalent refraction ≤ -0.50 D, and (3) hyperopia as spherical equivalent refraction $\geq +2.00$ D. Children were considered myopic with at least one eye was diagnosed, and hyperopic with hyperopia at least one eye, so long as neither eye was myopic. Children were considered emmetropic if neither eye was myopic or hyperopic. (4) Unilateral amblyopia was considered as a 2-line interocular difference in eyes with BCVA 0.2 or worse. And at least one of the following risk factors must be presence: strabismus or history of strabismus surgery, past or present obstruction of visual axis, anisometropia consistent with the worse eye (≥ 1.00 D SER anisohyperopia, ≥ 3.00 D SER anisomyopia, or ≥ 1.50 D anisoastigmatism). Bilateral amblyopia was defined as bilateral BCVA >0.3 . And there must be presence of bilateral visual axis obstruction or bilateral ametropia (≥ 4.00 D SE hyperopia, ≥ 6.00 D SE myopia, or ≥ 2.50 D astigmatism) [12]. Estimates of refractive error prevalence were based on children with successful cycloplegia in both eyes [11].

Presenting visual acuity was defined by the detected visual acuity when tested wearing currently available refractive correction, if any. Best-corrected visual acuity was the visual acuity achieved by subjects tested with pinhole or refraction. Corneal opacity was considered as the cause of vision loss in an eye if there was an easily visible opacity overlying the pupil to the extent that at least part of the pupil margin is blurred. Retinal disease was assigned the cause of vision loss in an eye found with chorioretinitis, optic disk atrophy, macular hole, and inherited retinal diseases. Cataract was assigned if a pupil appeared grey or white when examined with oblique light in a shaded or darkened area [13]. Children requiring medical or surgical treatment were recommended for eye clinic referral.

Data management and analysis

Clinical examination forms were verified at least twice to evaluate integrity and precision before entered into the database using Epidata software 3.1 (The Epidata Association, Odense, Denmark) by two separated individuals.

Statistical analysis was performed using SPSS software (version 20.0). Means \pm standard deviation, frequencies, and percentages were used to summarize the characteristics of the research subjects. The association between refractive status and the gender and ethnicity was explored using logistic regression. Confidence intervals and *P* values (significant at the $P < 0.05$ level) for prevalence estimates and regression models were calculated and adjusted for clustering effects associated with the sampling design [9].

Results

Demographics of study population

A final of 1856 participants completed all the examinations, with an overall response rate of 95.52%. The mean age of students was 6.82 ± 0.47 years (median age, 6.78 years; range, 5.89 to 10.32), 984 (53.0%) were males, and 1762 (94.9%) were Tibetan minority nationalities.

Distribution of Visual Acuity and Prevalence of Visual Impairment.

Distribution of uncorrected visual acuity and best-corrected visual acuity is presented in Table 1. Measurements of uncorrected visual acuity were not possible in 2 children. By uncorrected visual acuity, normal/near normal visual acuity in at least one eye was found in 1627 children (87.8%). Two hundred twenty-seven children (12.2%) were visually impaired in both eyes, and 2 (0.1%) of them were blind in both eyes.

For children wearing glasses in daily life, presenting visual acuity was recorded with their current refractive correction. Twelve out of 16 children achieved normal/near normal visual acuity in at least one eye. A 4.4% reduction of bilateral visually impairment was noticed over uncorrected vision without glasses.

After subjective refraction was performed to achieve best-corrected visual acuity, bilateral visual impairment was reduced to 2.7% ($n = 50$). And only 1 child remained bilateral blind. Nearly 80% (78.0%) children with bilateral visual impairment could reach normal/near-normal vision in at least one eye with refractive correction. These findings illustrate the potential benefit of spectacles in 77.0% of the children who had bilateral vision impairment.

Refractive data

A total of 1837 (99.0%) children had cycloplegic autorefraction performed successfully on both eyes.

Table 1 Distribution of Uncorrected and Best-Corrected Visual Acuity

Visual Acuity Category	Uncorrected Visual Acuity			Wearing glasses		Presenting Visual Acuity			Best corrected Visual Acuity		
	n	%	95%CI	n	%	n	%	95%CI	n	%	95%CI
Normal/near normal	1450	78.2	76.33–80.09	6	0.3	1454	78.4	76.55–80.30	1735	93.6	92.47–94.70
Unilateral impairment	177	9.5	8.21–10.88	6	0.3	183	9.9	8.51–11.23	69	3.7	2.86–4.58
Mild impairment in better eye	205	11.1	9.63–12.48	3	0.2	198	10.7	9.27–12.90	46	2.5	1.77–3.19
Moderate impairment in better eye	20	1.1	0.61–1.55	1	0.1	17	0.9	0.48–1.35	3	0.2	0.00–0.34
Blind in both eyes	2	0.1	0.00–0.26	0	0	2	0.1	0.00–0.26	1	0.0	0.00–0.16
All	1854	100		16	0.9	1854	100		1854	100	

CI confidence interval

Normal/near normal means log MAR visual acuity ≤ 0.2 in both eyes

Unilateral visual impairment means log MAR visual acuity ≤ 0.2 in one eye only

Mild visual impairment means log MAR visual acuity 0.3–0.5 in the better eye

Moderate visual impairment means log MAR visual acuity 0.6–0.9 in the better eye

Blindness means log MAR visual acuity ≥ 1.0 in the better eye

Spherical equivalent in right eyes in boys (1.03 ± 0.90 D) was less positive compared to girls (1.12 ± 0.94 D), as shown in Fig. 1 ($P=0.025$). The prevalence of myopia and hyperopia was 4.8 and 12.3% separately as shown in Table 2. Girls were more often hyperopic than boys.

The prevalence of astigmatism was 44.8% in total as shown in Table 3. There was no statistically significant differences between each gender and ethnicity groups, either in low (cylinder value -1.75 D to -0.75 D) or high astigmatism (cylinder value ≤ -2.00 D).

Risk factor analyses

Hyperopia was significantly associated with female gender ($P<0.001$), while there were no differences between

gender and myopia or astigmatism. No significant association of ethnicity and myopia, hyperopia and astigmatism was observed.

Ocular abnormalities

Tropia was present in 67 (3.6%) children. In all, 42 (62.7%) of them were exotropia. Exterior and anterior segment abnormalities were observed in 35 (1.9%) of the 1856 children: ptosis was observed in 3 eyes of 3 (0.2%) children; corneal abnormalities were present in 13 eyes of 11 (0.6%) children; pupillary abnormalities were noted in 3 eyes of 2 (0.1%) children; lenticular abnormalities were present in 15 eyes of 9 (0.5%) children; and other anterior segment abnormalities were observed in 12 eyes of 10 (0.5%) children.


**Fig. 1** Distribution of spherical equivalent refractive error (D) in right eyes by gender

Table 2 Prevalence of Refractive Status by Gender and Ethnicity

	Myopia			Hyperopia		
	n	%	95%CI	n	%	95%CI
Gender						
Boys	44	4.5	3.19–5.78	93	9.5	7–65–11.31
Girls	44	5.1	3.59–6.50	132	15.1*	12.76–17.52
Ethnicity						
Tibetan	82	4.7	3.68–5.65	211	12.0	10.48–13.51
Other	6	6.4	1.44–11.32	14	14.9	7.70–22.09
All	88	4.8		225	12.3	

CI confidence interval

*P<0.001

Table 3 Prevalence of Astigmatism* by Gender and Ethnicity

Cylinder Value (D)	−1.75 to −0.75		≤ −2.00		All	
	n	%	n	%	n	%
Gender						
Boys	359	36.5	86	8.7	445	45.2
Girls	300	34.4	87	10.0	387	44.4
Ethnicity						
Tibetan	632	35.9	167	9.5	799	45.3
Other	27	28.7	6	6.4	33	35.1
All	659	35.5	173	9.3	832	44.8

Data are presented as number (%) of children in each gender or ethnicity groups

*Categorized using the cylinder value in the eye with greater astigmatism by cycloplegic refraction

Table 4 Causes of Uncorrected Visual Acuity (UCVA) of 0.3 or Worse

Cause	Eyes with UCVA 0.3 or Worse		Percent Prevalence in the Population
	n	%	
Refractive error	285	70.5	15.4
Amblyopia	33	8.2	1.8
Corneal opacity	2	0.5	0.1
Cataract/posterior capsular opacification	1	0.2	0.1
Other	3	0.7	0.2
Unexplained cause	54	13.4	2.9

Causes of visual impairment

Visual impairment was mainly associated with refractive error, as shown in Table 4. A total of 285 children with uncorrected visual acuity 0.3 or worse in at least one eye attained visual acuity 0.2 or better in both eyes

with refractive correction. A total of 83 (2.9%) children presented with best-corrected visual acuity 0.3 or worse in right eye. Among them, 33 were unilateral amblyopia ($n=10$) or bilateral amblyopia ($n=23$), 7 with a tropia; 2 with anisometropia 2.00 spherical equivalent diopters or more. Other causes of visual impairment were uncommon.

Discussion

Various visual impairment disorders, such as refractive error, are related to several impact factors, and there are significant differences between regions, living environments, medical care and nationalities [14–16]. Because of the geographical remoteness from inland and backward economy, little information is available on the prevalence of ocular diseases in Tibetan children. The Lhasa Childhood Eye Study provides evaluations of the prevalence of refractive error and visual impairment situations in grade one children in Lhasa.

The current Lhasa Childhood Eye Study reported a 21.6% of the research population presenting with visual

acuity of 0.3 or worse in at least one eye in daily living condition, which decreased to 6.4% with best corrected vision. With presenting visual acuity, 11.7% remained bilateral visual impaired (worse than 0.3 in better-seeing eye), which is significantly higher than the Ireland Eye Study (6–7 years 3.43%) [17], and the Northern Ireland Childhood Errors of Refraction study (6–7 years 1.5%) [18]. The higher percentage of poor presenting visual acuity may result from the poor economy and shortage of necessary glasses. A total of 177 (78.0%) out of the 227 children with bilateral visual impairment could achieve normal/near-normal vision in at least one eye with refractive correction. Only 10 of them achieved normal/near normal visual acuity in at least one eye with their own glasses. A total of 33 out of 83 children with best-corrected visual acuity 0.3 or worse in right eye had asthenopia.

The major cause of visual impairment, such as refractive error and amblyopia, are largely preventable or treatable in early age [19, 20]. More attention should be paid to such a high prevalence of visual impairment of children in a remote low-income area. School-based screening programs provide the platform for inspecting a scalable number of children. School-age children comprise about a quarter of the population in developing countries and are easy to intervene. Besides, some conditions that can become permanent could be eliminated in early stage with a lower cost, such as refractive error and amblyopia. After proper examinations and identifying at risk children, more appropriate eye care services could be provided. To our knowledge, there is no cohort study following up the refractive status and visually impairment being implemented in Lhasa. The Lhasa Childhood Eye Study will provide further information in the next 5 year follow-up and is believed to contribute to relieve the visual abnormality of Tibetan school children.

The myopia prevalence in Lhasa Childhood Eye Study (6.82 ± 0.47 years 4.8%) was broadly in line with that reported in the Poland (7 years 4.0%) and Aston Eye Study (6–7 years 5.7%) [21, 22], higher than that reported in the UK Northern Ireland Childhood Errors of Refraction study (6–7 years 2.8%) [18], Australia (6 years 1.6%) [23], South Africa (7 years 2.5%) [24], but lower than urban China inland (7 years, 7.7% by autorefractometry) [25]. This difference may result from outdoor activity duration and study burden between different countries and regions.

The Lhasa Childhood Eye Study hyperopia prevalence (6.82 ± 0.47 years 12.3%) was comparable with Australia (6 years 13.2%) [23], and non-white ethnic groups of the Ireland Eye Study (6–7 years 11.1%) [17], and considerably lower than that reported in the other ethnic groups of the Ireland Eye Study (6–7 years 25.8–35.4%) [17], Northern Ireland (6–7 years 26%) [18], and higher than urban

China inland (7 years, 4.0% by autorefractometry) [25]. Same as myopia prevalence, outdoor activity duration and study burden may be the reason of the difference.

The astigmatism prevalence in Lhasa Childhood Eye Study (6.82 ± 0.47 years 44.8%) was similar to that found in Native American children (5–16 years 42%) [26], and significantly higher than Ireland Eye Study (6–7 years 19.2%) [17], Northern Ireland (6–7 years 24%) and Australia (6 years 7.6%) [18, 23].

Evidence from Refractive Error Study in Children demonstrated that differences in the prevalence of myopia were associated with ethnicity [5, 11, 15, 27–29]. In consist with Qian et al. (carried out in primary and secondary schools in the Tibet Naidong district) [7], a lower prevalence of myopia in Tibet than China inland was witnessed. However, there were no significant associations of ethnic or gender with myopia or astigmatism in Lhasa Childhood Eye Study. The prevalence of hyperopia in females was significantly higher than that in males in our survey. The difference might result from the different lifestyle, such as outdoor activity [14, 30, 31]. Not inconsistent with the finding in Naidong, no difference of the prevalence of hyperopia between ethnicity was recorded in Lhasa Childhood Eye Study. It is possible that environmental and lifestyle differences underlie the difference among studies, as well as the difference of study design [4, 32].

There are a few limitations in this study. The prevalence of blindness and low vision may be underestimated for children with low vision in both eyes may dropped out school and went to schools for the blind, which might have biased the analysis of the distribution of eye diseases. The reason for children wear glasses is not recorded in the present article. In addition, we did not record sociodemographic characteristics that might have been associated with refractive error, such as outdoor activity and family income.

Conclusions

Our data revealed a lower prevalence of myopia compared to China inland, but a significantly higher prevalence of visual impairment compared to developed county [1, 33]. It is clear that vision screening in schools accompanied by health education is an important approach to alleviate the influence of visual impairment. Our survey demonstrated the unmet need for the necessity of corrective spectacles. Effective strategies are needed to eliminate this easily treated cause of significant visual impairment.

Acknowledgements

The authors would like to acknowledge the support and participation of the schools, the children, and their parents and guardians in the Lhasa Childhood Eye Study.

Authors' contributions

JC conducted data analysis and interpretation. JF designed and supervised this study and revised the final manuscript. WC, ZM, LL, HS, YY and WD assisted with the acquisition of the data. WC, ZM and LL helped develop the statistical method. All authors read and approved the final manuscript.

Funding

This work was supported by High Level Health Technical Talent Training Program of Beijing Municipal Health Bureau [2015–3-023]; Beijing Municipal Science & Technology Commission [Z171100001017066], and the Capital Health Development Special Fund of China -Major Project [SF-2018-1-2051], Youth Project [2018–4-1083]. The funders had no role in the study design, data collection, analysis or preparation of the manuscript.

Availability of data and materials

The datasets used and analysed during the current study available from the corresponding author on reasonable request.

Declarations

Ethics approval and consent to participate

The protocols in this study were approved by the Institutional Review Board of Beijing Tongren Hospital, Capital Medical University (TRECXY2019–146), and adhere to the principles of the Declaration of Helsinki. Written informed consent was obtained from a parent or legal guardian of each child prior to the examinations.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Beijing Tongren Eye Center, Beijing Tongren Hospital, Capital Medical University, Beijing Key Laboratory of Ophthalmology & Visual Sciences, No.1, Dong Jiao Min Xiang Street, Dongcheng District, Beijing 100730, China. ²Beijing Institute of Ophthalmology, Beijing, China.

Received: 26 January 2021 Accepted: 23 September 2021

Published online: 12 October 2021

References

- Naidoo KS, Leasher J, Bourne RR, et al. Global vision impairment and blindness due to uncorrected refractive error, 1990–2010. *Optom Vis Sci*. 2016;93:227–34.
- Simons K. Preschool vision screening: Rationale, methodology and outcome. 1996;41:3–30.
- Schneider J, Leeder SR, Gopinath B, et al. Frequency, course, and impact of correctable visual impairment (uncorrected refractive error). *Surv Ophthalmol* 2010; 55: 539–560. 2010/09/21. DOI: <https://doi.org/10.1016/j.survophthal.2010.02.004>.
- Elsman EBM, Al Baaj M, van Rens G, et al. Interventions to improve functioning, participation, and quality of life in children with visual impairment: a systematic review. *Surv Ophthalmol* 2019; 64: 512–557. 2019/02/01. DOI: <https://doi.org/10.1016/j.survophthal.2019.01.010>.
- Wu JF, Bi HS, Wang SM, et al. Refractive error, visual acuity and causes of vision loss in children in Shandong, China. The Shandong Children Eye Study. *PLoS One* 2013; 8: e82763. 2014/01/01. DOI: <https://doi.org/10.1371/journal.pone.0082763>.
- You QS, Wu LJ, Duan JL, et al. Prevalence of myopia in school children in greater Beijing: the Beijing Childhood Eye Study. *Acta Ophthalmol* 2014; 92: e398–406. 2014/08/29. DOI: <https://doi.org/10.1111/aos.12299>.
- Qian X, Liu B, Wang J, et al. Prevalence of refractive errors in Tibetan adolescents. *BMC Ophthalmol* 2018; 18: 118. 2018/05/12. DOI: <https://doi.org/10.1186/s12886-018-0780-8>.
- Bali J, Chaudhary KP and Thakur R. High altitude and the eye: a case controlled study in clinical ocular anthropometry of changes in the eye. *High Alt Med Biol* 2005; 6: 327–338. 2005/12/15. DOI: <https://doi.org/10.1089/ham.2005.6.327>.
- Chen W, Fu J, Meng Z, et al. Lhasa childhood eye study: the rationale, methodology, and baseline data of a 5 year follow-up of school-based cohort study in the Tibetan plateau region of Southwest China. *BMC Ophthalmol* 2020; 20: 250. 2020/06/24. DOI: <https://doi.org/10.1186/s12886-020-01522-w>.
- Li SM, Rafael I, He L, et al. Intraocular pressure and myopia progression in Chinese children: the Anyang Childhood Eye Study. *British Journal of Ophthalmology: bjophthalmol-2017-311831*.
- He M, Huang W, Zheng Y, et al. Refractive error and visual impairment in school children in rural southern China. *Ophthalmology* 2007; 114: 374–382. 2006/11/25. DOI: <https://doi.org/10.1016/j.ophtha.2006.08.020>.
- Multi-ethnic Pediatric Eye Disease Study Group. Prevalence of amblyopia and strabismus in African American and Hispanic children ages 6 to 72 months the multi-ethnic pediatric eye disease study. *Ophthalmology* 2008; 115: 1229–1236.e1221. 2007/10/24. DOI: <https://doi.org/10.1016/j.ophtha.2007.08.001>.
- WHO Programme for the Prevention of Blindness. Coding instructions for the WHO/PBL eye examination record (version III): World Health Organization; 1988.
- Wang GQ, Bai ZX, Shi J, et al. Prevalence and risk factors for eye diseases, blindness, and low vision in Lhasa, Tibet. *Int J Ophthalmol* 2013; 6: 237–241. 2013/05/03. DOI: <https://doi.org/10.3980/j.issn.2222-3959.2013.02.24>.
- Pokharel GP, Negrel AD, Munoz SR, et al. Refractive Error Study in Children: results from Mechi Zone, Nepal. *Am J Ophthalmol* 2000; 129: 436–444. 2000/04/15. DOI: [https://doi.org/10.1016/s0002-9394\(99\)00453-5](https://doi.org/10.1016/s0002-9394(99)00453-5).
- Pi LH, Chen L, Liu Q, et al. Prevalence of eye diseases and causes of visual impairment in school-aged children in Western China. *J Epidemiol* 2012; 22: 37–44. 2011/11/30. DOI: <https://doi.org/10.2188/jea.20110063>.
- Harrington SC, Stack J, Saunders K, et al. Refractive error and visual impairment in Ireland schoolchildren. *Br J Ophthalmol* 2019; 103: 1112–1118. 2018/10/14. DOI: <https://doi.org/10.1136/bjophthalmol-2018-312573>.
- O'Donoghue L, McClelland JF, Logan NS, et al. Refractive error and visual impairment in school children in Northern Ireland. *Br J Ophthalmol* 2010; 94: 1155–1159. 2010/05/25. DOI: <https://doi.org/10.1136/bjo.2009.176040>.
- McCann S, Flitcroft I, Lalor K, et al. Parental attitudes to myopia: a key agent of change for myopia control? *Ophthalmic Physiol Opt* 2018; 38: 298–308. 2018/04/25. DOI: <https://doi.org/10.1111/opo.12455>.
- Guo Y, Liu LJ, Xu L, et al. Outdoor activity and myopia among primary students in rural and urban regions of Beijing. *Ophthalmology* 2013; 120: 277–283. 2012/10/27. DOI: <https://doi.org/10.1016/j.ophtha.2012.07.086>.
- Logan NS, Shah P, Rudnicka AR, et al. Childhood ethnic differences in ametropia and ocular biometry: the Aston Eye Study. *Ophthalmic Physiol Opt* 2011; 31: 550–558. 2011/07/19. DOI: <https://doi.org/10.1111/j.1475-1313.2011.00862.x>.
- Czepiet D, Zejmo M and Mojsa A. Prevalence of myopia and hyperopia in a population of Polish schoolchildren. *Ophthalmic Physiol Opt* 2007; 27: 60–65. 2007/01/24. DOI: <https://doi.org/10.1111/j.1475-1313.2006.00419.x>.
- Robaei D, Rose K, Kifley A, et al. Patterns of spectacle use in young Australian school children: findings from a population-based study. *Journal of American Association for Pediatric Ophthalmology and Strabismus*. 2005;9:579–83. <https://doi.org/10.1016/j.jaapos.2005.07.005>.
- Naidoo KS, Raghunandan A, Mashige KP, et al. Refractive error and visual impairment in African children in South Africa. *Invest Ophthalmol Vis Sci* 2003; 44: 3764–3770. 2003/08/27. DOI: <https://doi.org/10.1167/iov.03-0283>.
- He M, Zeng J, Liu Y, et al. Refractive error and visual impairment in urban children in southern China. *Investigative Ophthalmology & Visual Science*. 2004;45. <https://doi.org/10.1167/iov.03-1051>.
- DOBSON V, MILLER JM and HARVEY EM. Corneal and Refractive Astigmatism in a Sample of 3- to 5-Year-Old Children with a High Prevalence of Astigmatism. *Optometry and Vision Science*; 76: 855–860.
- Zhao J, Pan X, Sui R, et al. Refractive Error Study in Children: results from Shunyi District, China. *Am J Ophthalmol* 2000; 129: 427–435. 2000/04/15. DOI: [https://doi.org/10.1016/s0002-9394\(99\)00452-3](https://doi.org/10.1016/s0002-9394(99)00452-3).
- Goh PP, Abqariyah Y, Pokharel GP, et al. Refractive error and visual impairment in school-age children in Gombak District, Malaysia.

- Ophthalmology 2005; 112: 678–685. 2005/04/06. DOI: <https://doi.org/10.1016/j.ophtha.2004.10.048>.
29. Dandona R, Dandona L, Srinivas M, et al. Refractive error in children in a rural population in India. *Invest Ophthalmol Vis Sci* 2002; 43: 615–622. 2002/02/28.
 30. Wu P-C, Tsai C-L, Wu H-L, et al. Outdoor activity during class recess reduces myopia onset and progression in school children. *Ophthalmology*. 2013;120:1080–5. <https://doi.org/10.1016/j.ophtha.2012.11.009>.
 31. Rose KA, Morgan IG, Smith W, et al. Myopia, lifestyle, and schooling in students of Chinese ethnicity in Singapore and Sydney. *Arch Ophthalmol* 2008; 126: 527–530. 2008/04/17. DOI: <https://doi.org/10.1001/archophth.126.4.527>.
 32. Gwiazda J and Weber C. Comparison of spherical equivalent refraction and astigmatism measured with three different models of autorefractors. *Optom Vis Sci* 2004; 81: 56–61. 2004/01/30. DOI: <https://doi.org/10.1097/00006324-200401000-00011>.
 33. Lu P, Chen X, Zhang W, et al. Prevalence of ocular disease in Tibetan primary school children. *Can J Ophthalmol*. 2008;43:95–9. <https://doi.org/10.3129/i07-194>.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

