

Ophthalmic Epidemiology



ISSN: 0928-6586 (Print) 1744-5086 (Online) Journal homepage: www.tandfonline.com/journals/iope20

Quality of Refractive Error Care in Cambodia: An Unannounced Standardized Patient Study

Anthea Burnett, Ngy Meng, Do Seiha, Neath Kong, Seila Chea, Malis Dean, Piseth Horm, Kim San Meas, Beatrice Varga, Suit May Ho, Myra McGuinness & Ling Lee

To cite this article: Anthea Burnett, Ngy Meng, Do Seiha, Neath Kong, Seila Chea, Malis Dean, Piseth Horm, Kim San Meas, Beatrice Varga, Suit May Ho, Myra McGuinness & Ling Lee (2025) Quality of Refractive Error Care in Cambodia: An Unannounced Standardized Patient Study, Ophthalmic Epidemiology, 32:1, 25-33, DOI: 10.1080/09286586.2024.2321890

To link to this article: https://doi.org/10.1080/09286586.2024.2321890

© 2024 The Author(s). Published with license by Taylor & Francis Group, LLC.	→ View supplementary material 🗹
Published online: 20 Mar 2024.	Submit your article to this journal 🗹
Article views: 917	View related articles 🗹
View Crossmark data 🗹	Citing articles: 1 View citing articles



ORIGINAL ARTICLE



Quality of Refractive Error Care in Cambodia: An Unannounced Standardized **Patient Study**

Anthea Burnett^{a,b}, Ngy Meng^c, Do Seiha^c, Neath Kong^d, Seila Chea^d, Malis Dean^d, Piseth Horm^e, Kim San Meas^d, Beatrice Varga^a, Suit May Ho^a, Myra McGuinness^{f,g}, and Ling Lee^{a,b}

^aInternational Programs Division, The Fred Hollows Foundation, Melbourne, Australia; ^bSchool of Optometry and Vision Science, The University of New South Wales, Sydney, Australia; 'National Program for Eye Health, Phnom Penh, Cambodia; 'International Programs Division, The Fred Hollows Foundation Cambodia, Phnom Penh, Cambodia; Eye Care Foundation, Phnom Penh, Cambodia; Melbourne School of Population and Global Health, University of Melbourne, Melbourne, Australia; ⁹Biostatistics, Centre for Eye Research Australia, Melbourne, Australia

ABSTRACT

Purpose: Quality-of-care in refractive error services is essential, as it directly affects vision outcomes, wellbeing, educational attainment, and workforce participation. In Cambodia, uncorrected refractive error is a leading cause of mild and moderate vision impairment in adults. We evaluated the quality of refractive error care in Cambodia by estimating the proportion of prescribed and dispensed spectacles appropriate for people's refractive error needs and factors associated with spectacle quality.

Methods: A cross-sectional protocol was employed with 18 Khmer-speaking adult participants observing testing procedures in 156 optical services across six provinces in 2022. A total of 496 dispensed spectacles were assessed against spectacle quality indicators.

Results: The analysis revealed that 35.1% of dispensed spectacles were of optimal quality. The most common error observed in sub-optimal spectacles was the presence of horizontal prism outside of tolerance limits. The study also found that 44.0% of emmetrope visits involved unnecessary prescription spectacle recommendations, and 18.3% of written prescriptions did not correspond with dispensed spectacles. Sex differences were observed, with men predominantly providing refractive error care and women more likely to be unnecessarily recommended prescription spectacles.

Conclusion: The findings highlight the importance of prioritizing quality-of-care in refractive error services. A key recommendation is to consider regulatory mechanisms to ensure optical services employ appropriately qualified staff. Additionally, efforts should be made to eliminate unnecessary prescriptions -- especially for emmetropes and females -- standardize written prescriptions, ensure consistent pupil distance measurements, reduce reliance on autorefraction, and address the gender imbalance in the refractive error workforce.

ARTICLE HISTORY

Received 11 June 2023 Revised 17 December 2023 Accepted 18 February 2024

KEYWORDS

Cambodia; optometry; public health; quality of care; refractive error

Introduction

Quality-of-care is a fundamental component within the context of universal health coverage (UHC), as it ensures that individuals receive effective, safe, and timely health care tailored to their needs, without experiencing financial hardship. In terms of refractive error care, the emphasis on quality-of-care is significant, as it directly impacts visual outcomes, as well as wellbeing, educational attainment, and workforce participation.² High-quality refractive error care involves accurate diagnosis, appropriate prescription, and precise spectacle dispensing, all of which contribute to optimal visual acuity (VA) and comfort. By prioritising quality-of-care in refractive error services, healthcare providers can fully address the needs of patients with varying degrees of visual impairment.

In Cambodia, uncorrected refractive error causes 61% of mild vision impairment (presenting VA < 6/12-6/18), and 17.5% of moderate vision impairment (presenting VA < 6/18-6/60) in adults over 50 years.³ In adults over 19 years, the estimated prevalence of myopia, hyperopia, and astigmatism is 49.1%, 4.45%, and 9.1% respectively,⁴ and 21.5% of adults over 50 years live with near vision impairment.⁵ In Takeo province, it was reported that more men than women (35% vs 10%) wear glasses, with the majority (68%) of glasses purchased from the market, 26%

3053, Australia

CONTACT Anthea Burnett a.burnett@unsw.edu.au International Programs Division, The Fred Hollows Foundation, 33 Lincoln Square S, Carlton, VIC

Supplemental data for this article can be accessed online at https://doi.org/10.1080/09286586.2024.2321890

from an optical shop, 16% given from a relative, and 10% from Takeo Eye Hospital.⁶

Effective refractive error care (eREC) is one of two global targets endorsed by Member States at the 74th World Health Assembly to measure progress towards achieving UHC.⁷ This target, which entails a 40-percentage point increase in eREC, is designed to monitor both the effective coverage and quality of refractive error care. Despite the absence of eREC estimates in Cambodia, it is anticipated that significant enhancements in the quantity and quality of refractive services will be necessary to meet this global target. Therefore, it is crucial to identify and address the specific challenges and gaps in refractive error care within Cambodia to ensure progress towards UHC and improved eye health for all.

Unannounced Standardized Patients (USPs), covertly impersonate patients, and are considered the gold standard for assessing quality in clinical practice. Their usage ensures an unbiased assessment of clinical techniques and services, as healthcare providers don't alter their behaviours, not knowing they're under observation. USPs have been instrumental in evaluating family planning, pharmaceutical dispensing, and clinical prescribing patterns in lower to middle income settings,⁹ and even in assessing refractive error outcomes. 10-12

Refractive error care encompasses identification of patient needs, a precise refraction, and dispensing spectacles that meet the patient's refraction. A Quality of Refractive Error Care (Q.REC) study leverages USPs to generate evidence on the standard of delivered refractive services, informing potential alterations to practices and policies.

The aim of this study was to evaluate the quality of refractive error care in Cambodia by using USPs to estimate the proportion of prescribed and dispensed spectacles appropriate for people's refractive error needs. Secondly, this study also aimed to assess the optical service and USP characteristics associated with spectacle quality and explore unnecessary prescribing practices.

Material and methods

This study employs a previously published cross-sectional protocol¹³ to recruit 18 fluent Khmer-speaking adult USPs with either:

- Myopia: Spherical equivalent < -.50 DS in at least
- Hyperopia: Spherical equivalent > +.50 DS in at least one eye
- Astigmatism: > .50 DC in at least one eye
- Emmetropia: Spherical equivalent ≥ -.50 DS and ≤ +.50 DS in both eyes

• Presbyopia: ≥ 1.00 DS above the best optical distance correction

USPs were excluded if they did not have good ocular health, had a history of refractive surgery, had manifest or intermittent strabismus or amblyopia, or any health conditions that could impact refractive error consistency.

The sampling frame for eligible optical services was compiled by collaborating with relevant Ministries of Health, Optical Councils, Optometry Associations, as well as cross-referencing with Google Maps. Optical services were deemed eligible if they provided refraction and dispensing services, were not previously known to USPs and did not have staff who worked across multiple services already selected. We excluded provinces with fewer than five optical services from the sample frame. Additionally, Banteay Meanchey province was not included, despite having an estimated seven optical services, due to the distance and extensive travel costs.

A total of 202 optical services in Phnom Penh, Kandal, Siem Reap, Battambang, Kampong Cham and Takeo provinces were selected. However, as some services had relocated or closed due to COVID-19, only 156 stores were visited. Service owners received written information about the study along with opportunities to opt-out verbally or in writing.

All USPs underwent an online training course in Khmer, which introduced them to the study and the equipment used during refraction. A standardized quiz was administered to confirm their understanding. The USPs then participated in a three-day in-person training course, during which a study optometrist briefed them on the tasks involved and provided sufficient time to practice acting as patients. The USPs were also provided with standard responses to use during optical service visits. Post-training, observations by a study optometrist were conducted to assess whether the USPs could accurately identify elements of refraction and dispensing techniques. If the USPs were unable to do so accurately, further training was provided. 13 USPs visited optical services, observed testing procedures and obtained one pair of single-vision spectacles, distance or near depending on the clinician's recommendation. USPs also recorded whether they felt the clinician communicated clearly with them during the eye test. Research optometrists then assessed VA, comfort and lens prescriptions by asking USPs if they experienced any eye strain or discomfort while wearing the spectacles. Non-presbyopic USPs with emmetropia did not purchase spectacles unless there was a risk of them being detected. The purchased



spectacles were categorized as distance or near based on the corresponding written prescription collected. This study follows the Checklist for Reporting Research Using Simulated Patient Methodology (CRiSP).¹⁴

Quality outcomes

Spectacle quality was categorized as optimal or suboptimal (Supplementary 1). Optimal quality singlevision spectacles met all quality components in both lenses, with the baseline distance sphere power plus any near addition for single vision near spectacles. Cylindrical axis quality was only assessed for USPs with cylinder power detected at baseline. Induced horizontal prism was derived using Prentice's rule, from the lens power at horizontal meridian and the amount of lens decentration from the averaged baseline pupillary distance. Vertical prism was derived from focimetry.

The written prescription obtained from each visit was classified as optimal or sub-optimal relative to the baseline refraction using the tolerance limits for spherical and cylinder power and cylinder axis components. Dispensed spectacles matched the written prescription if spherical and cylindrical power were within 0.25 D, and cylinder axis tolerance limits were met. Unnecessary prescribing occurred when USPs with emmetropia and no presbyopia were advised to purchase spectacles.

VA with dispensed spectacles was considered good if it was within 1.5 lines of monocular and binocular baseline best-corrected VA. USPs reported any discomfort or eye strain while wearing the spectacles.

Data analysis

Analyses were conducted using Stata/BE v16.0 (StataCorp LLC, College Station, TX). All dispensed spectacles were included in primary analysis. Secondary analyses examined associations between USP/optical service characteristics and spectacle quality.

Descriptive statistics were reported for baseline USP and optical service characteristics. Proportions were calculated for each outcome of interest (spectacle and prescription quality, comfort, spectacle-corrected VA, and unnecessary prescribing). Logit-transformed 95% confidence intervals (CI) were estimated using robust standard errors to account for intra-service correlation.

Univariable logistic regression was used to examine associations between USPs, optical service characteristics and spectacle quality. Multivariable models evaluated associations between service procedures (focimetry, autorefraction, distance and near subjective refraction) and spectacle quality, adjusting for USP baseline refraction type. Covariates were selected as potential confounders based on subject matter knowledge and the number of observations in each category. To account for within-optical service correlation, robust standard errors were used.

To account for the unequal numbers of spectacles dispensed in each province, weighting was applied so that pooled estimates could be generalized. Weights were assigned based on 2019 census population sizes, with Battambang, Kampong Cham, Ta Keo, Phnom Penh, and Siem Reap contributing 16.5%, 14.9%, 15.0%, 36.8%, and 16.7%, respectively. For each analysis set, each pair of spectacles within a province was weighted equally. As the primary aim of the study was to estimate the proportion of optimal-quality spectacles, comparative statistics were considered exploratory and no adjustment for multiple testing was applied.

Results

This study analyzed 16 USPs, consisting of 9 women and 7 men with ages ranging from 20 to 52 years. At baseline, all USPs had a best-corrected VA of logMAR .04 (~6/6 part) or better for distance and .2 (~6/9.5, N4) or better for near vision in each eye. The baseline spherical equivalent refraction for distance ranged from -5.31D to +1.04D, with a maximum of 1.25D cylinder power. The average interocular difference in spherical equivalent was 0.37D (SD .78, range 0.00 to 3.29D).

Each USP visited between 23 and 45 optical services, with each service being visited by up to five USPs (60% of services were visited by five USPs, 28% of services were visited by four USPs, 11% were visited by three USPs and the remaining services were visited by either one or two USPs). A total of 663 visits were made to 156 optical services, with 544 spectacles dispensed. After excluding ineligible visits (seven where the USP reported detection; eight visits to duplicate services; and 33 visits where a USP was later diagnosed with high Hba1c levels, which could have affected refractive error during data collection), 496 spectacles from 615 visits were included. Figure 1 depicts a flowchart of optical services, USPs, and spectacles included in the analysis.

Quality of spectacles and prescriptions

Of the 496 spectacles included, 259 (52.2%) were single-vision distance, and 237 (48.8%) were singlevision near, with 24 (4.8%) dispensed from public or non-governmental organization services and the remaining 472 from the private sector. Although




Figure 1. Flow chart of unannounced standardised patient (USP) visits to optical services and dispensed spectacles.

Table 1. Percentage of dispensed spectacles and written prescriptions with optimal quality^a.

	Optimal quality (95% CI) ^b					
	Dispens	Dispensed spectacles		Written prescriptions		
	Frequency	%	Frequency	%		
Battambang	6/16	37.5 (18.4–61.5)	9/16	56.3 (36.4–74.3)		
Kampong Cham	5/15	33.3 (11.8-65.1)	5/15	33.3 (33.3-33.3)		
Ta Keo	8/15	53.3 (31.5-74.0)	5/15	33.3 (33.3-33.3)		
Phnom Penh	131/386	33.9 (29.6-38.6)	110/386	28.5 (23.8-33.7)		
Siem Reap	13/64	20.3 (12.6-31.2)	35/64	54.7 (43.2-65.7)		
Pooled (Unweighted)	163/496	32.9 (29.0-36.9)	164/496	33.1 (28.8–37.7)		
Pooled (Weighted)		35.1 (29.9–40.7)		38.9 (34.4–43.6)		

^aOptimal quality for written prescriptions only included the spherical, cylindrical and axis components.

a higher proportion of spectacles from public or non-governmental services were optimal (41.7%) versus private services (32.4%), this difference was non-significant (p = 0.35). After weighting by population size, 35.1% (95%CI: 36.4–49.3%) of the

spectacles were optimal quality, ranging from 20.3% in Siem Reap to 53.3% in Takeo (Table 1). The most common error observed in sub-optimal spectacles was horizontal prism outside of tolerance limits (96/333, 28.8%).

^b95% Cls adjusted for intra-service correlation.

There was no significant difference in the proportion of distance and near spectacles that were of optimal quality (distance: 32.0% vs. near: 33.8%, p = 0.69). When comparing the individual spectacle components, a significantly lower proportion of distance spectacles were within tolerance limits for cylinder power (distance: 91.1% vs. near: 99.6%, p < 0.001) and axis (distance: 43.1% vs. near: 100.0%, p < 0.001) compared to near spectacles. However, no single-vision near spectacles were dispensed for those with astigmatism. A significantly lower proportion of near spectacles were within tolerance limits for horizontal prism compared to distance spectacles (distance: 90.3% vs. near: 48.1%, *p* < 0.001).

On average, dispensed spectacles exhibited wider lens centres than the baseline pupillary distance for both distance (mean: 4.7 mm, SD: 3.7 mm) and near spectacles (mean: 6.8 mm, SD: 4.4 mm). The range fluctuated from 16.7 mm wider to 2.3 mm narrower than baseline for distance spectacles and 19.7 mm wider to .3 mm narrower for near spectacles.

Of the dispensed spectacles, 81.7% (n = 405) matched the written prescription, and 38.9% (weighted, 95%CI: 34.4–43.6%) of the written prescriptions were within the tolerance limits for the sphere, cylinder, and axis components. Written prescriptions were inconsistent, particularly with the provision of pupillary distances. Over half of the written prescriptions (55.8%) did not provide any pupillary distance, and only 33 (6.7%) provided pupillary distance for both distance and near. For written prescriptions where only one pupillary distance was provided, sometimes it was unclear whether the value corresponded to distance or near.

Optical service characteristics and associations with spectacle quality

Most clinicians providing refractive error care were men (82.9%). Autorefraction was the most common procedure (92.5%), while distance subjective refraction was absent in 35.9% of visits (Table 2).

Table 2. Distribution of testing procedures and associations with spectacle quality.

	Visits (<i>N</i> = 615)	Dispensed spectacles (n = 496)					
Testing procedure	Frequency (%)	Optimal quality frequency (%)	OR	[95% CI]	<i>p</i> -value ^a		
Focimetry performed							
No/Did not take spectacles	342 (55.6)	77/22733 (33.9)	1.00				
Yes	273 (44.4)	86/269 (32.0)	0.92	[0.65,1.30]	0.618		
Distance VA checked at the beginning							
No	254 (41.3)	65/206 (31.6)	1.00				
Yes	361 (58.7)	98/290 (33.8)	1.11	[0.76,1.60]	0.590		
VA checked with pinhole							
No .	537 (87.3)	145/421 (34.4)	1.00				
Yes	78 (12.7)	18/75 (24.0)	0.60	[0.34,1.05]	0.073		
Near VA checked at the beginning		, ,		,			
No	388 (63.1)	95/297 (32.0)	1.00				
Yes	227 (36.9)	68/199 (34.2)	1.10	[0.77,1.58]	0.591		
Autorefraction performed	(,	,		,			
No	46 (7.5)	13/38 (34.2)	1.00				
Yes	569(92.5)	150/458 (32.8)	0.94	[0.46,1.90]	0.586		
Retinoscopy performed	,	,		,			
No	575 (93.5)	155/458 (33.8)	1.00				
Yes	40 (6.5)	8/38 (21.1)	0.52	[0.25,1.10]	0.087		
Distance subjective refraction	(****)	,		,			
Spherical/cylinder power not checked	318 (51.7)	68/221 (30.8)	1.00				
Only spherical component tested	100 (16.3)	43/89 (49.4)	2.20	[1.31,3.68]	0.003		
Only cylinder tested	5 (0.8)	0/5 (0.0)	NA	[
Sphere and cylinder tested	192 (31.2)	52/183 (28.4)	0.89	[0.57,1.41]	0.626		
Near VA with distance lenses checked	(/			[,]			
No	432 (70.2)	107/337 (31.8)	1.00				
Yes	183 (29.8)	55/159 (35.2)	1.17	[0.81,1.68]	0.400		
Near subjective refraction performed	(====)	22, 122 (22.2)		[,]			
No	345 (56.1)	70/243 (28.8)	1.00				
Yes	270 (43.9)	93/253 (36.8)	1.44	[1.00,2.07]	0.052		
Clinician used a phoropter	(,	11, 201 (0111)		[,]			
No	257 (41.8)	71/235 (30.2)	1.00				
Yes	137 (22.3)	45/127 (35.4)	1.27	[0.81,1.97]	0.293 ^b		
No distance subjective refraction	221 (35.9)	.57 .27 (551.)	,	[0.0.7.157]	0.275		
Clinician used a trial frame	22. (33.3)						
No	29 (4.7)	16/22 (72.7)	1.00				
Yes	365 (59.3)	100/340 (29.4)	0.16	[0.07,0.36]	<0.001 ^b		
No distance subjective refraction	221 (35.9)	. 56,5 15 (2511)	00	[0.07,0.50]	10.00		
Pupillary distance checked	22. (55.2)						
No	525 (85.4)	129/410 (31.5)	1.00				
Yes	90 (14.6)	34/86 (39.5)	1.42	[0.91,2.23]	0.122		

^aAssessed via univariable logistic regression, accounting for within-optical service correlation.

^bExcludes 134 visits at which distance refraction was not performed, yet spectacles dispensed.

Surprisingly, the proportion receiving optimal spectacles was greater after only the spherical component of distance refraction had been performed compared to both spherical and cylindrical components. Conversely, more sub-optimal spectacles were dispensed following a trial frame refraction compared to not using a trial frame (Table 2). For most visits, communication was perceived as clear during eye examinations (84.7%), outcomes (91.5%), and spectacle recommendations (76.9%). Clear communication during eye examinations (OR [95%CI]: 5.00 [2.29–10.91], p < 0.001) and outcomes (OR [95%CI]: 7.49 [1.63–34.31], p = 0.01) was associated with receiving optimal spectacles.

After adjustment for USP refraction and stratification by spectacle type (distance vs near), receiving optimal distance spectacles was associated with having either myopia or hyperopia compared to having astigmatism alone or myopia astigmatism, and having focimetry performed and having only the cylinder component of distance subjective refraction testing (Table 3). Receiving optimal near spectacles was associated with having emmetropia with presbyopia compared to having hyperopia with presbyopia. No testing procedures were associated with receiving optimal near spectacles (Table 3).

A total of 159 visits to 138 optical services were made by emmetropic individuals without presbyopia, with 44.0% (70/159) recommended spectacles and 46 dispensed (all in Phnom Penh). Men were less likely to receive unnecessary spectacle recommendations than women USPs (OR [95%CI]: .17 [0.06–0.489]; p < 0.001). Spectacles were less likely to be recommended after checking distance (OR [95%CI]: .09 [0.04–0.20], p < 0.001) and near VA (OR [95%CI]: .18 [0.06–0.49], p = 0.001) at the beginning of the eye examination. Conversely, both spherical and cylindrical distance subjective refraction (OR [95%CI]: 2.90 [1.03–8.19]; p = 0.04) and near subjective refraction (OR [95%CI]: 9.45 [4.23–21.14]; p < 0.001) was associated with unnecessary spectacle recommendations.

Vision and comfort

Good spectacle-corrected VA was achieved in 95.4% of distance spectacles (weighted percentage, 95% CI: 88.9–98.1%) for each eye separately, and 97.7% (weighted percentage, 95% CI: 94.4–99.1%) of near spectacles. More optimal distance spectacles achieved good distance

Table 3. Association between testing procedures and spectacle quality among spectacle types (Single-vision distance, n = 246; Single-vision near, n = 237).

	Single-vision distance			Single-vision near				
	Optimal quality frequency (%)	Odds Ratio	[95% CI]	<i>p</i> -value ^a	Optimal quality frequency (%)	Odds Ratio	[95% CI]	<i>p</i> -value ^a
Refractive status at baseline Emmetropic	-	-	-	-	71/165 (43.0)	1.00		
Astigmatism only/myopic astigmatism	10/119 (8.4)	1.00			-	-	-	-
Myopia only	11/14 (78.6)	101.48	[7.18,1434.88]	0.001	-	-	-	-
Hyperopia only	49/113 (43.4)	19.39	[7.44,50.48]	< 0.001	9/72 (12.5)	.17	0[.07,.40]	< 0.001
Focimetry performed								
No/USP did not take spectacles	16/65 (24.6)	1.00			48/149 (32.2)	1.00		
Yes	54/181 (29.8)	5.56	[2.25,13.75]	< 0.001	32/88 (36.4)	1.10	[0.62,1.94]	0.738
Autorefraction performed								
No	8/15 (53.3)	1.00			4/22 (18.2)	1.00		
Yes	62/231 (26.8)	0.58	[0.07, 5.06]	0.624	76/215 (35.3)	1.39	[0.30,6.43]	0.674
Distance VA checked at beginning								
No	29/111 (26.1)	1.00			-	-	-	-
Yes	41/135 (30.4)	0.82	[0.384,1.77]	0.610	-	-	-	-
Distance subjective refraction								
Spherical component not tested*	14/76 (18.4)	1.00			45/141 (31.9)	1.00		
Only spherical component tested	18/32 (56.3)	2.05	[0.59,7.08]	0.255	22/52 (42.3)	1.66	[0.77,3.55]	0.195
Sphere & cylinder tested	38/138 (27.5)	1.87	[0.73,4.81]	0.196	13/44 (29.5)	1.57	[0.75,3.31]	0.233
Near VA tested with distance lenses								
No	_	-	-	-	3/8 (37.5)	1.00		
Yes	-	-	-	-	77/229 (33.6)	.94	[.50,1.76]	0.848
Near subjective refraction performed								
No	-	-	-	-	66/203 (32.5)	1.00		
Yes	-	-	-	-	14/34 (41.2)	.83	[0.17,4.04]	0.817
Pupillary distance checked								
No No	51/195 (26.2)	1.00			66/203 (32.5)	1.00		
Yes	19/51 (37.3)	1.95	[0.93,4.09]	0.076	14/34 (41.2)	1.80	[0.85,3.83]	0.126

^{95%} CI = 95% confidence interval.

^aEach multivariable logistic regression model was assessed with robust standard errors to account for intra-service correlation.

^bIncludes 2 visits where only the cylindrical component was checked during distance subjective refraction.



VA compared to sub-optimal spectacles (100.0% vs 92.6%, p = 0.01). A similar proportion achieved good near VA between optimal and sub-optimal single-vision near spectacles (99.3% vs 98.0%, p = 0.44).

Discomfort or eyestrain was experienced in 47.3% (weighted, 95%CI: 38.8-56.0%) of distance spectacles, and at near in 35.1% of all spectacles (weighted, 95%CI: 31.1-39.3%). Fewer USPs experienced discomfort when wearing optimal spectacles compared to sub-optimal spectacles at distance (37.8% vs 55.7%, p = 0.01) and near (16.8% vs 41.1%, *p* < 0.001).

Discussion

In this Q.REC Cambodia study, 35.1% of dispensed spectacles were optimal (weighted), 44.0% of emmetrope visits involved unnecessary prescription spectacle recommendations, and 18.3% of written prescriptions did not correspond with dispensed spectacles. Optimal spectacles were associated with effective communication and refractive status rather than testing procedures. Unnecessary spectacle recommendations were linked to increased subjective refraction testing and the absence of VA testing. Sex differences were observed, with men predominantly providing refractive error care and women more likely to be unnecessarily recommended spectacles.

Although an adequate level of VA was achieved with most sub-optimal spectacles, discomfort or eyestrain was much more prevalent with sub-optimal spectacles. This is likely to impact the frequency and duration of spectacle use in the community.

Inadequate dispensing seems to significantly contribute to sub-optimal spectacles, with horizontal prism being the most common issue. Only 14.6% of visits involved observed pupil distance measurement, and the average lens centres were wider than patients' baseline pupil distance. This suggests that optical service staff may not be appropriately measuring pupil distance, or not accounting for convergence at near, which typically results in a narrower pupil distance. Additional training to enhance optical dispensers' or mechanics' skills to accurately measure pupil distances may reduce avoidable prismatic effects induced by spectacles.

The prevalence of astigmatism and hyperopia in Phnom Penh adults is estimated at 9.12% and 4.35% respectively. Individuals with astigmatism or hyperopia combined with presbyopia were less likely to receive optimal spectacles. This highlights the need for improved refraction and/or dispensing techniques, particularly for those with complex prescriptions who are in greater need of appropriate spectacles.

In Cambodia, autorefraction is more commonly used than subjective refraction. Autorefraction is mainly a preliminary step in determining lens prescriptions, similar to subjective refraction. Studies have reported the inadequacy of autorefractors for prescribing glasses, 15,16 except for non-presbyopic adults in lowresource areas. 17,18 Retinoscopy is a more accurate alternative to autorefraction, but still requires complementary subjective methods for adult patients.¹⁹

Written prescriptions exhibited many inconsistencies, particularly with respect to pupillary distance. Standardized approaches to written spectacle prescriptions are needed and should include all parameters necessary for dispensers to make optimal spectacles.

Patient-centredness is vital for quality eye care. Ormsby et al. found that most adults were satisfied with their eye care when they received glasses at a public hospital.²⁰ Though our study didn't measure satisfaction, it highlighted the importance of good communication for optimal spectacles. Communication skills include giving clear instructions, understanding patient needs, and informing them about their eye care.²¹ Moreover, quality care involves avoiding unnecessary prescriptions. In our study, 44% of USPs who didn't need glasses were prescribed them, with women being more likely to receive unnecessary prescriptions than men. This over-prescription could be due to insufficient refraction skills or an intent to increase sales -the gender discrepancy warrants further investigation.

The recent Cambodian National Strategic Plan for Blindness Prevention & Control 2021 - 2030, emphasizes strengthening integrated eye health service delivery.²² The focus on enhancing primary eye care for those most in need, aligns with the pursuit of improved refractive error quality. The Cambodian Health Workforce Development Plan 2006-2015 acknowledges optometry and spectacle suppliers as integral to the healthcare system, especially in the privatefor-profit sector. However, it notes a deficiency in regulation and registration in this sector, affecting service quality and workforce management. The plan mentions a seemingly balanced gender distribution in the health workforce, but lacks detailed breakdowns by sector or category. A new plan for 2021-2030 is in draft form.

This study has certain limitations that need to be acknowledged. Firstly, while visits were made to Kandal province, no spectacles were included in the primary analyses. The results may also not be representative of provinces that were not visited due to budgetary constraints. Secondly, the Q.REC indicator does not currently have a minimum threshold to denote 'high quality', although it highlights various aspects of refractive error care that could improve. Thirdly, the Q.REC



indicators cannot currently be used to assess the quality of bifocal or progressive addition lenses. They also do not include other vital aspects of spectacle quality, including frame quality and lens type, which could further affect the overall quality of the spectacles. Finally, all near spectacles achieved good VA (with both eyes open and each eye separately), which may be attributable to the absence of smaller steps in the near VA charts used.

Improving refractive error care in Cambodia is necessary, as only 35.1% of dispensed spectacles optimally address refractive error needs. Refraction and dispensing skills are currently limiting high-quality care in Cambodia, and there is a need for targeted clinical education, practice, and regulatory interventions. It is recommended that regulatory mandates are introduced to ensure optical services employ appropriately training and qualified staff. This measure would uphold professional standards and ultimately improve patient care. Additionally, there is a need to foster a diverse and inclusive workforce, with more opportunities for women to become integral members of the refractive error eye care. The evidence generated here can contribute to developing and implementing wellintegrated eye health service delivery systems in line with the National Strategic Plan's strategic priorities.

Acknowledgments

We would like to thank Tokyo Bak for assistance in USP recruitment, training and data collection.

Disclosure statement

LL and AB are consultants to The Fred Hollows Foundation.

Funding

This work was funded by The Fred Hollows Foundation with support from the Australia Government through the Australian NGO Cooperation Program (ANCP).

Data availability statement

Data are available upon reasonable request.

Ethics approval

This research adhered to the tenets of the Declaration of Helsinki, and was approved by the University of New South Wales Human Research Ethics Committee (HC210102), and the National Ethics Committee for Health Research in Cambodia (043 NECHR).

Patient consent for publication

Written informed consent was obtained from USPs. Optical services owners were informed of the study in writing and had the opportunity to opt-out verbally or in writing.

References

- 1. World Health Organization. Universal health coverage (UHC): what is UHC? Secondary universal health coverage (UHC): what is UHC? 1 April 2021. https://www. who.int/news-room/fact-sheets/detail/universalhealth-coverage-(uhc).
- 2. Burton MJ, Ramke J, Marques AP, et al. The lancet global health commission on global eye health: vision beyond 2020. Lancet Glob Health. 2021;9(4):e489-e551. doi:10.1016/S2214-109X(20)30488-5. published Online First: Epub Date.
- 3. National Program for Eye Health (NPEH). National rapid assessment of avoidable blindness in Cambodia 2018-2019. Ministry Of Health. 2019.
- 4. Shin H-M, Shin H-U, Jung J-H. Prevalence of refractive error in adults in Phnom Penh, Cambodia. J Korean Ophthalmic Opt Soc. 2018;23(4):301-307. doi:10.14479/ ikoos.2018.23.4.301.
- 5. Bourne R, Steinmetz JD, Flaxman S, et al. Trends in prevalence of blindness and distance and near vision impairment over 30 years: an analysis for the global burden of disease study. Lancet Glob Health. 2021;9(2):e130-e143. doi:10.1016/S2214-109X(20)30425-3.
- 6. Ormsby GM, Arnold AL, Busija L, Morchen M, Bonn TS, Keeffe JE. The impact of knowledge and attitudes on access to eye-care services in Cambodia. Asia Pac J Ophthalmol (Phila). 2012;1(6):331-5. doi:10. 1097/APO.0b013e31826d9e06. published Online First: Epub Date.
- 7. World Health Organization (WHO). Resolution WHA74(12): integrated people-centred eye care, including preventable vision impairment and blindness. In: Seventy-Fourth World Health Assembly. Geneva: World Health Organization; 2021. https:// apps.who.int/gb/ebwha/pdf_files/WHA74/A74(12)-en. pdf.
- 8. Rethans JJ, Gorter S, Bokken L, Morrison L. Unannounced standardised patients in real practice: a systematic literature review. Med Educ. 2007;41 (6):537-49. doi:10.1111/j.1365-2929.2006.02689.x. published Online First: Epub Date.
- 9. Wiseman V, Lagarde M, Kovacs R, et al. Using unannounced standardised patients to obtain data on quality of care in low-income and middle-income countries: key challenges and opportunities. BMJ Glob Health. 2019;4(5):e001908. doi:10.1136/bmjgh-2019-001908. published Online First: Epub Date.
- 10. Lee L, Burnett AM, D'Esposito F, et al. Indicators for assessing the quality of refractive error care. Optom Vis 2021;98(1):24-31. doi:10.1097/OPX. 000000000001629. published Online First: Epub Date.
- 11. Shah R, Ctori I, Edgar DF, Parker P. Use of standardised patients in optometry training. Clin Exp Optom.



- 2021;104(8):848-53. doi:10.1080/08164622.2021. 1896332. published Online First: Epub Date.
- 12. Nie J, Zhang L, Gao J, et al. Using incognito standardised patients to evaluate quality of eye care in China. Br J Ophthalmol. 2021;105(3):311-16. doi:10.1136/ bjophthalmol-2019-315103. published Online First: Epub Date.
- 13. Burnett AM, Lee L, McGuinness M, Varga B, Perez Hazel Y, Ho SM. Quality of refractive error care (Q. REC) in Cambodia, Malaysia and Pakistan: protocol for a cross-sectional unannounced standardised patient study. BMJ Open. 2022;12(3):e057594. doi:10.1136/bmjo pen-2021-057594. published Online First: Epub Date.
- 14. Amaratunge S, Harrison M, Clifford R, Seubert L, Page A, Bond C. Developing a checklist for reporting research using simulated patient methodology (CRiSP): a consensus study. Int I Pharm Pract. 2021;29 (3):218-227. doi:10.1093/ijpp/riaa002. published Online First: Epub Date.
- 15. Pesudovs K, Weisinger HS. A comparison of autorefractor performance. Optom Vis Sci. 2004;81(7):554-8. doi:10.1097/00006324-200407000-00018. published Online First: Epub Date.
- 16. Bennett JR, Stalboerger GM, Hodge Schornack MM. Comparison of refractive assessment by wavefront aberrometry, autorefraction, and subjective refraction. J Optom. 2015;8(2):109-115. doi:10. 1016/j.optom.2014.11.001. published Online First: Epub Date.

- 17. Durr NJ, Dave SR, Lim D, Joseph S, Ravilla TD, Lage E. Quality of eyeglass prescriptions from a low-cost wavefront autorefractor evaluated in rural India: results of a 708-participant field study. BMJ Open Ophthalmol. 2019;4(1):e000225. doi:10.1136/bmjophth-2018-000225. published Online First: Epub Date.
- 18. Joseph S, Varadaraj V, Dave SR, et al. Investigation of the accuracy of a low-cost, portable autorefractor to well-tolerated eyeglass prescriptions: a randomized crossover trial. Ophthalmology. 2021;128(12):1672-80. doi:10.1016/j.ophtha.2021.05. 030. published Online First: Epub Date.
- 19. Jorge J, Queiros A, Almeida JB, Parafita MA. Retinoscopy/Autorefraction: which is the best starting point for a noncycloplegic refraction? Optom Vis Sci. 2005;82(1):64-8 doi:10.1097/01.OPX.0000150182. 91410.97.
- 20. Ormsby GM, Grant-Skiba D, Naidoo K, Keeffe JE. Patient perspectives on acquiring spectacles: a cambodian experience. Asia Pac J Ophthalmol (Phila). 2016;5(5):339-43. doi:10.1097/APO. 000000000000176. published Online First: Epub Date.
- 21. World Health Organization. Eye Care Competency Framework. Geneva: World Health Organization; 2022.
- 22. The National Program for Eye Health, Ministry of Health. The national strategic plan for blindness prevention & control 2021 - 2030: access, coverage, quality, equity: Kingdom of Cambodia. 2022.